

Tutte polynomial

Chromatic polynomial $C_\Gamma(q)$.

A *coloring* of Γ with q colors is a map $c : V(\Gamma) \rightarrow \{1, \dots, q\}$. A coloring c is *proper* if for any edge e : $c(v_1) \neq c(v_2)$, where v_1 and v_2 are the endpoints of e .

Definition 1. $C_\Gamma(q) := \#$ of proper colorings of Γ in q colors.

Properties (Definition 2).

$$C_\Gamma = C_{\Gamma-e} - C_{\Gamma/e} ;$$

$$C_{\Gamma_1 \sqcup \Gamma_2} = C_{\Gamma_1} \cdot C_{\Gamma_2}, \quad \text{for a disjoint union } \Gamma_1 \sqcup \Gamma_2 ;$$

$$C_\bullet = 1 .$$

Tutte polynomial $T_\Gamma(x, y)$.

Definition 1.

$$T_\Gamma = T_{\Gamma-e} + T_{\Gamma/e} \quad \text{if } e \text{ is neither a bridge nor a loop ;}$$

$$T_\Gamma = xT_{\Gamma/e} \quad \text{if } e \text{ is a bridge ;}$$

$$T_\Gamma = yT_{\Gamma-e} \quad \text{if } e \text{ is a loop ;}$$

$$T_{\Gamma_1 \sqcup \Gamma_2} = T_{\Gamma_1, \Gamma_2} = T_{\Gamma_1} \cdot T_{\Gamma_2} \quad \text{for a disjoint union } \Gamma_1 \sqcup \Gamma_2 \\ \text{and a one-point join } \Gamma_1 \cdot \Gamma_2 ;$$

$$T_\bullet = 1 .$$

Properties.

$$T_\Gamma(1, 1) \quad \text{is the number of spanning trees of } \Gamma ;$$

$$T_\Gamma(2, 1) \quad \text{is the number of spanning forests of } \Gamma ;$$

$$T_\Gamma(1, 2) \quad \text{is the number of spanning connected subgraphs of } \Gamma ;$$

$$T_\Gamma(2, 2) = 2^{|E(\Gamma)|} \quad \text{is the number of spanning subgraphs of } \Gamma .$$

$$C_\Gamma(q) = q^{k(\Gamma)} (-1)^{r(\Gamma)} T_\Gamma(1 - q, 0) .$$

Definition 2.

Let \bullet F be a graph;

- $v(F)$ be the number of its vertices;
- $e(F)$ be the number of its edges;
- $k(F)$ be the number of components of F ;
- $r(F) := v(F) - k(F)$ be the *rank* of F ;
- $n(F) := e(F) - r(F)$ be the *nullity* of F ;

$$T_\Gamma(x, y) := \sum_{F \subseteq E(\Gamma)} (x - 1)^{r(\Gamma) - r(F)} (y - 1)^{n(F)}$$

Dichromatic polynomial $Z_\Gamma(q, v)$ (**Definition 3**).

Let $Col(\Gamma)$ denote the set of colorings of Γ with q colors, and let $D : E(\Gamma) \times Col(\Gamma) \rightarrow \{0, 1\}$ be defined by the formula $D(e, c) = 1$ is and only if $c(v_1) = c(v_2)$, where v_1 and v_2 are the endpoints of e .

$$Z_\Gamma(q, v) := \sum_{c \in Col(\Gamma)} \prod_{e \in E(\Gamma)} (1 + vD(e, c))$$

Properties .

$$\begin{aligned} Z_\Gamma &= Z_{\Gamma-e} + vZ_{\Gamma/e} ; \\ Z_{\Gamma_1 \sqcup \Gamma_2} &= Z_{\Gamma_1} \cdot Z_{\Gamma_2} , \quad \text{for a disjoint union } \Gamma_1 \sqcup \Gamma_2 ; \\ Z_\bullet &= q ; \end{aligned}$$

$$Z_\Gamma(q, v) = \sum_{F \subseteq E(\Gamma)} q^{k(F)} v^{e(F)} ;$$

$$\begin{aligned} C_\Gamma(q) &= Z_\Gamma(q, -1) ; \\ Z_\Gamma(q, v) &= q^{k(\Gamma)} v^{r(\Gamma)} T_\Gamma(1 + qv^{-1}, 1 + v) ; \\ T_\Gamma(x, y) &= (x - 1)^{-k(\Gamma)} (y - 1)^{-v(\Gamma)} Z_\Gamma((x - 1)(y - 1), y - 1) . \end{aligned}$$

Definition 4.

For a connected graph Γ fix an order of its edges: e_1, e_2, \dots, e_m . Let T be a spanning tree.

An edge $e_i \in E(T)$ is called *internally active (live)* if $i < j$ for any edge e_j connecting the two components of $T - e_i$

An edge $e_j \notin E(T)$ is called *externally active (live)* if $j < i$ for any edge e_i in the unique cycle of $T \cup e_j$.

Let $i(T)$ and $j(T)$ be the numbers of internally and externally active edges correspondingly.

$$T_\Gamma(x, y) := \sum_T x^{i(T)} y^{j(T)}$$