


Graphs

Definition. A graph G is a finite set of vertices $V(G)$ and a finite set $E(G)$ of unordered pairs (x, y) of vertices $x, y \in V(G)$ called edges.

A graph may have loops (x, x) and multiple edges when a pair (x, y) appears in $E(G)$ several times. Pictorially we represent the vertices by points and edges by lines connecting the corresponding points. Topologically a graph is a 1-dimensional cell complex with $V(G)$ as the set of 0-cells and $E(G)$ as the set of 1-cells. Here are two pictures representing the same graph.



Tutte polynomial

Chromatic polynomial $C_G(q)$.

A coloring of G with q colors is a map $c : V(G) \rightarrow \{1, \dots, q\}$. A coloring c is proper if for any edge $e: c(v_1) \neq c(v_2)$, where v_1 and v_2 are the endpoints of e .

Definition 1. $C_G(q) := \#$ of proper colorings of G in q colors.

Properties (Definition 2).

$$C_G = C_{G-e} - C_{G/e} ;$$

$$C_{G_1 \sqcup G_2} = C_{G_1} \cdot C_{G_2}, \quad \text{for a disjoint union } G_1 \sqcup G_2 ;$$

$$C_{\bullet} = q .$$

Flow polynomial $Q_G(q)$.

A q -flow on G is an assignment of a value $0, 1, \dots, q - 1$ to every edge of G with arbitrarily chosen orientation of its edges in such a way that the total flow entering and leaving each vertex is congruent modulo q .

Definition 1. $Q_G(q) := \#$ of nowhere-zero q -flows on G .

Properties (Definition 2).

$$Q_G(q) = 0 \quad \text{if } G \text{ has a bridge ;}$$

$$Q_G(q) = (q - 1)Q_{G-e}(q) \quad \text{if } e \text{ is a loop ;}$$

$$Q_G(q) = -Q_{G-e}(q) + Q_{G/e} \quad \text{if } e \text{ is neither a bridge nor a loop ;}$$

$$Q_G(q) = C_{G^*}(q)/q \quad \text{for dual planar graphs } G \text{ and } G^* .$$

Tutte polynomial $T_G(x, y)$.

Definition 1.


$$T_G = T_{G-e} + T_{G/e} \quad \text{if } e \text{ is neither a bridge nor a loop ;}$$

$$T_G = xT_{G/e} \quad \text{if } e \text{ is a bridge ;}$$

$$T_G = yT_{G-e} \quad \text{if } e \text{ is a loop ;}$$

$$T_{G_1 \sqcup G_2} = T_{G_1 \cdot G_2} = T_{G_1} \cdot T_{G_2} \quad \text{for a disjoint union } G_1 \sqcup G_2 \text{ and a one-point join } G_1 \cdot G_2 ;$$

$$T_{\bullet} = 1 .$$



Properties.

- $T_G(1, 1)$ is the number of spanning trees of G ;
- $T_G(2, 1)$ is the number of spanning forests of G ;
- $T_G(1, 2)$ is the number of spanning connected subgraphs of G ;
- $T_G(2, 2) = 2^{|E(G)|}$ is the number of spanning subgraphs of G ;
- $C_G(q) = q^{k(G)}(-1)^{r(G)}T_G(1 - q, 0)$;
- $Q_G(q) = (-1)^{n(G)}T_G(0, 1 - q)$.

Definition 2.

- Let F be a graph;
- $v(F)$ be the number of its vertices;
 - $e(F)$ be the number of its edges;
 - $k(F)$ be the number of connected components of F ;
 - $r(F) := v(F) - k(F)$ be the *rank* of F ;
 - $n(F) := e(F) - r(F)$ be the *nullity* of F ;

$$T_G(x, y) := \sum_{F \subseteq E(G)} (x - 1)^{r(G) - r(F)} (y - 1)^{n(F)}$$

Dichromatic polynomial $Z_G(q, v)$ (Definition 3).

Let $Col(G)$ denote the set of colorings of G with q colors.

$$Z_G(q, v) := \sum_{c \in Col(G)} (1 + v)^{\# \text{ edges colored not properly by } c}$$

Properties .

- $Z_G = Z_{G-e} + vZ_{G/e}$;
- $Z_{G_1 \sqcup G_2} = Z_{G_1} \cdot Z_{G_2}$, for a disjoint union $G_1 \sqcup G_2$;
- $Z_{\bullet} = q$;

$$Z_G(q, v) = \sum_{F \subseteq E(G)} q^{k(F)} v^{e(F)} ;$$

$$C_G(q) == Z_G(q, -1) ;$$

$$Z_G(q, v) = q^{k(G)} v^{r(G)} T_G(1 + qv^{-1}, 1 + v) ;$$


$$T_G(x, y) = (x - 1)^{-k(G)} (y - 1)^{-v(G)} Z_G((x - 1)(y - 1), y - 1) .$$

Potts model in statistical mechanics (Definition 4).

Potts model (C.Domb 1952); $q = 2$ the Ising model (W.Lenz, 1920)

Let G be a graph.

Particles are located at vertices of G . Each particle has a *spin*, which takes q different values . A *state*, $\sigma \in \mathcal{S}$, is an assignment of spins to all vertices of G . Neighboring particles interact with each other only if their spins are the same.



The energy of the interaction along an edge e is $-J_e$ (*coupling constant*). The model is called *ferromagnetic* if $J_e > 0$ and *antiferromagnetic* if $J_e < 0$.

Energy of a state σ (*Hamiltonian*),

$$H(\sigma) = - \sum_{(a,b)=e \in E(G)} J_e \delta(\sigma(a), \sigma(b)).$$

Boltzmann weight of σ :

$$e^{-\beta H(\sigma)} = \prod_{(a,b)=e \in E(G)} e^{J_e \beta \delta(\sigma(a), \sigma(b))} = \prod_{(a,b)=e \in E(G)} \left(1 + (e^{J_e \beta} - 1) \delta(\sigma(a), \sigma(b)) \right),$$

where the *inverse temperature* $\beta = \frac{1}{\kappa T}$, T is the temperature, $\kappa = 1.38 \times 10^{-23}$ joules/Kelvin is the *Boltzmann constant*.

The Potts partition function (for $x_e := e^{J_e \beta} - 1$)

$$Z_G(q, x_e) := \sum_{\sigma \in \mathcal{S}} e^{-\beta H(\sigma)} = \sum_{\sigma \in \mathcal{S}} \prod_{e \in E(G)} (1 + x_e \delta(\sigma(a), \sigma(b)))$$

Properties of the Potts model Probability of a state σ : $P(\sigma) := e^{-\beta H(\sigma)} / Z_G$.

Expected value of a function $f(\sigma)$:

$$\langle f \rangle := \sum_{\sigma} f(\sigma) P(\sigma) = \sum_{\sigma} f(\sigma) e^{-\beta H(\sigma)} / Z_G.$$

Expected energy: $\langle H \rangle = \sum_{\sigma} H(\sigma) e^{-\beta H(\sigma)} / Z_G = -\frac{d}{d\beta} \ln Z_G$.

Fortuin—Kasteleyn'1972: $Z_G(q, x_e) = \sum_{F \subseteq E(G)} q^{k(F)} \prod_{e \in F} x_e$,

where $k(F)$ is the number of connected components of the spanning subgraph F .

$$Z_G = Z_{G \setminus e} + x_e Z_{G/e}.$$

Spanning tree generating function (Definition 5).

For a connected graph G fix an order of its edges: e_1, e_2, \dots, e_m . Let T be a spanning tree.

An edge $e_i \in E(T)$ is called *internally active (live)* if $i < j$ for any edge e_j connecting the two components of $T - e_i$.

An edge $e_j \notin E(T)$ is called *externally active (live)* if $j < i$ for any edge e_i in the unique cycle of $T \cup e_j$.

Let $i(T)$ and $j(T)$ be the numbers of internally and externally active edges correspondingly.

$$T_G(x, y) := \sum_T x^{i(T)} y^{j(T)}$$