

CONTENTS

1. Review: Complex numbers and functions of a complex variable	5
2. Convergent power series	7
2.1. Series	7
2.2. Power series	7
3. Continuity and differentiability	8
3.1. Differentiability of power series	9
3.2. Some basic functions	11
3.3. Operations with power series	13
3.4. The Cauchy–Riemann equations	14
3.5. Analyticity at infinity	14
4. Integrals	15
5. Cauchy’s formula	19
5.1. Homotopic curves	19
5.2. Independence of the integral on the path	19
5.3. Cauchy’s Formula	21
6. Taylor series of analytic functions	22
7. The Fundamental Theorem of Algebra	24
8. More properties of analytic functions	25
9. Harmonic functions	26
9.1. Potential and Hamiltonian flows	28
10. The maximum modulus principle	28
Steepest ascent and descent lines	29
10.1. Application: soap films	30
10.2. Cauchy principal value integrals (PV)	31
10.3. Bi-analytic bijections (automorphisms) of $\mathbb{D}$	32
11. Poisson’s formula	32
11.1. The Dirichlet problem for the Laplacian in $\mathbb{D}$	34
12. Isolated singularities, Laurent series	35
13. Laurent series and Fourier series	37
14. Calculating the Taylor series of simple functions	38
15. Residues and integrals	39
16. Calculating the Taylor series of simple functions	40
17. Residues and integrals	41
18. Integrals of trigonometric functions	42
19. Counting zeros and poles	43
19.1. Constancy of the number of roots of $f(s) - \zeta$	43
19.2. Some further properties of the logarithm	44
19.3. The argument principle	45
19.4. Hurwitz’s theorem	46

19.5.	Rouché's Theorem	47
20.	The inverse function theorem	48
21.	Analytic continuation	50
22.	The Schwarz reflection principle	53
23.	Multi-valued functions	55
23.1.	Generalization: log of a function	57
23.2.	General powers of z	57
24.	Conformal Mapping	58
24.1.	Uniqueness	59
24.2.	Preservation of angles and small shapes: heuristics	60
24.3.	Preservation of angles	61
24.4.	Rescaling of arc length	62
24.5.	Transformation of areas	62
24.6.	Puiseux series	63
25.	Evaluation of definite integrals	65
25.1.	Fourier-type integrals	67
25.2.	Further examples	68
26.	Integrals of branched functions	70
26.1.	Automorphisms	72
26.2.	Automorphisms of the plane	73
26.3.	The Riemann sphere	74
26.4.	Linear fractional transformations (LFTs): a first view	74
26.5.	Automorphisms of the Riemann sphere	75
27.	Linear fractional transformations (Möbius transformations)	76
27.1.	Finding specific LFTs	77
27.2.	Mappings of regions	77
27.3.	The Schwarz lemma	78
27.4.	Automorphisms of the unit disk	79
28.	Some special biholomorphic transformations	80
	Application: Ideal flow above a semicircular bump (normalized case $U = 1$)	83
29.	The Riemann Mapping Theorem	89
29.1.	Equicontinuity and Normal Families	92
29.2.	The Riemann Mapping Theorem	94
29.3.	The Riemann Mapping Theorem	94
29.4.	Behavior at the boundary of biholomorphisms: a general but weaker result	97
30.	Boundary behavior.	98
30.1.	Behavior at the boundary of biholomorphisms: a general but weaker result	98
30.2.	A reflection principle for harmonic functions	99
30.3.	Analytic arcs	100

30.4.	An extension of the Schwarz reflection principle	101
30.5.	Behavior at the boundary, a stronger result	104
31.	Conformal mappings of polygons and the Schwarz-Christoffel formulas	106
31.1.	Heuristics	107
31.2.	Conformal map of the UHP onto the interior of a triangle of angles $\pi\alpha, \pi\beta, \pi\gamma$	113
31.3.	Schwarz triangle functions and hypergeometric functions.	114
32.	Optional material: curvilinear triangles	115
33.	Two other important examples of Schwarz-Christoffel transformations	120
33.1.	Another look at the sine function	120
34.	Mapping of a rectangle: Elliptic functions	121
34.1.	Continuation to the whole of $\mathbb{C}$. Double periodicity	122
35.	Riemann surfaces: a first view	123
36.	The modular group	128
36.1.	Bases of lattices	129
36.2.	The fundamental region of $PSL(2, \mathbb{Z})$	130
36.3.	The generators of the modular group	132
36.4.	The hyperbolic plane	133
36.5.	Schwarz reflection of domains about a circle	133
37.	Entire and Meromorphic functions	134
37.1.	A historical context	134
37.2.	Partial fraction decompositions	135
37.3.	The Mittag-Leffler theorem	137
37.4.	Further examples	138
38.	Infinite products	139
38.1.	Uniform convergence of products	140
38.2.	Example: the sin function	141
38.3.	Canonical products	142
38.4.	Counting zeros of analytic functions. Jensen's formula	144
38.5.	Entire functions of finite order	145
38.6.	Estimating analytic functions by their real part	147
39.	Hadamard's theorem	148
39.1.	Canonical products	149
40.	The minimum modulus theorem; end of proof of Theorem 39.1	150
40.1.	Some applications	153
41.	The Phragmén-Lindelöf Theorem	154
41.1.	An application to Laplace transforms	155
41.2.	A Laplace inversion formula	157
41.3.	Abstract Stokes phenomena	158

42.	Elliptic functions	158
42.1.	The period module	159
42.2.	General properties of elliptic functions	159
42.3.	The Weierstrass elliptic functions	160
42.4.	The modular function λ	163
42.5.	The action of the $\mathrm{PSL}(2, \mathbb{Z})$ on λ	163
42.6.	The conformal mapping of λ	165
43.	The uniformization theorem	167
43.1.	An example: M , the universal cover of $\hat{\mathbb{C}} \setminus \{0, 1, \infty\}$	167
44.	The little Picard theorem	169
45.	Riemann-Hilbert problems: an introduction	171
45.1.	A simple Riemann-Hilbert problem	171
46.	Cauchy type integrals	172
46.1.	Asymptotic behavior of $\Phi(z)$ for large z	172
46.2.	Regularity and singularities	172
46.3.	Examples	175
47.	More general scalar R-H problems	177
47.1.	Scalar homogeneous R-H problems	177
47.2.	Applications	178
48.	Asymptotic series	180
48.1.	More general asymptotic series	182
48.2.	Asymptotic power series	184
48.3.	Integration and differentiation of asymptotic power series.	186
48.4.	Watson's Lemma	187
48.5.	The Gamma function	188
48.6.	Lebesgue's Dominated Convergence Theorem	190
48.7.	Appendix to Chapter 8	190
48.8.	Some facts about the topology of $\mathbb{C}$	190
49.	Soap films and the minimal surface equation	192
	References	194

It is hard to overstate the importance of complex analysis in virtually every area of mathematics. Its power stems from several sources; one of them is that it represents the “ultimate” extension of the real numbers: it satisfies all the axioms of $\mathbb{R}$ except for the ordering property. Furthermore, the Gelfand–Mazur theorem shows that the only normed division algebras over $\mathbb{R}$ are $\mathbb{R}$, $\mathbb{C}$, and the quaternions (the latter being non-commutative).

Functions that extend from $\mathbb{R}$ to $\mathbb{C}$ —the analytic functions—preserve all their local properties, by the so-called *principle of permanence of relations* (to be discussed later). Hence, to understand such functions, it is essential to consider their full complex domain of definition.

A simple example is the function

$$\frac{1}{1+x^2}.$$

It admits a convergent power series at $x = 0$ of radius 1. There is nothing remarkable about the points $x = \pm 1$ on the real axis, but as soon as we extend the function to $\mathbb{C}$, we see that $\pm i$ are singular points, being the roots of the denominator.

More generally, the fundamental theorem of algebra holds in $\mathbb{C}$ but not in $\mathbb{R}$.

As another illustration, the exponential decay of Fourier transforms of real-analytic functions cannot be shown by purely real methods; one “needs” complex-analytic arguments.

1. REVIEW: COMPLEX NUMBERS AND FUNCTIONS OF A COMPLEX VARIABLE

- Complex numbers $\mathbb{C}$ form a field. Addition and multiplication of complex numbers obey the same rules as their real counterparts.
- There is no “good” order relation on $\mathbb{C}$. Aside from this, we operate with complex numbers in essentially the same way as with real numbers.
- A function f of a complex variable is a function defined on some subset of $\mathbb{C}$ with values in $\mathbb{C}$. Equivalently, we can view it as a pair of real-valued functions of two real variables. Writing $z = x + iy$ with $x, y \in \mathbb{R}$ and $i^2 = -1$, we have

$$f(z) = f(x + iy) = u(x, y) + iv(x, y).$$

- Note that $i^2 = (-i)^2 = -1$. There is no intrinsic distinction between i and $-i$. This entails a fundamental symmetry of the theory: invariance under complex conjugation.¹
- Using only the algebraic properties of complex numbers we can already define several elementary functions: z , $1/z$, and more generally, for $m \in \mathbb{Z}$, the power z^m , as well as any polynomial

$$\sum_{m=0}^K c_m (z - z_0)^m.$$

- To define and study more general functions we must introduce continuity, derivatives, and limits. Writing $z = x + iy$, the modulus

$$|z| = \sqrt{x^2 + y^2}$$

provides a natural notion of length (or smallness). It induces a norm under which $\mathbb{C}$ becomes a **complete metric space**. Convergence then reduces to convergence of real numbers:

$$(1.2) \quad z_n \rightarrow z \quad \Leftrightarrow \quad |z - z_n| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

- The *topology* of $\mathbb{C}$ coincides with that of $\mathbb{R}^2$, under the identification $z = x + iy \leftrightarrow (x, y) \in \mathbb{R}^2$. Some basic topological facts are reviewed in Appendix §48.7.
- In what follows, a *domain* in $\mathbb{C}$ means an open connected subset of $\mathbb{C}$. Examples include disks of radius $r > 0$ centered at $z_0 \in \mathbb{C}$:

$$(1.3) \quad \mathbb{D}(z_0, r) := \{z \in \mathbb{C} : |z - z_0| < r\}.$$

The special cases $r = 0$ (the empty set, $\emptyset$) and $r = \infty$ (the whole of $\mathbb{C}$) are also open sets. The unit disk

$$\mathbb{D} := \mathbb{D}(0, 1)$$

will play a special role as a canonical choice of a disk.

Exercise 1.1. Show that $z_n \rightarrow z$ if and only if $\operatorname{Re}(z_n) \rightarrow \operatorname{Re}(z)$ and $\operatorname{Im}(z_n) \rightarrow \operatorname{Im}(z)$. Using the completeness of $\mathbb{R}$, show that $\mathbb{C}$ is a complete normed space.

Definition 1.2. For functions, limits reduce to the real case: we write

$$\lim_{z \rightarrow z_0} f(z) = a \quad \Longleftrightarrow \quad |f(z) - a| \rightarrow 0 \quad \text{as } z \rightarrow z_0.$$

¹More precisely, the map $z = x + iy \mapsto \bar{z} = x - iy$ is an involution and a field automorphism of $\mathbb{C}$.

2. CONVERGENT POWER SERIES

2.1. Series. A complex series is an infinite sum

$$(2.2) \quad \sum_{k=0}^{\infty} a_k,$$

where $a_k \in \mathbb{C}$ for $k \geq 0$. It is said to *converge* if the sequence of *partial sums*

$$(2.3) \quad S_N := \sum_{k=0}^N a_k$$

converges as $N \rightarrow \infty$.

The series is said to *converge absolutely* if the real series

$$(2.4) \quad \sum_{k=0}^{\infty} |a_k|$$

converges.

Exercise 2.3. (a) Show that a necessary condition for convergence is $a_k \rightarrow 0$ as $k \rightarrow \infty$.

(b) Show that absolute convergence implies convergence.

(c) Verify that the standard convergence tests from real analysis (ratio test, n -th root test, etc. which do not involve ordering) apply equally to complex series.

2.2. Power series. A *power series* centered at $z_0 \in \mathbb{C}$ is a series of the form

$$(2.5) \quad \sum_{k=0}^{\infty} c_k (z - z_0)^k,$$

where $c_k, z, z_0 \in \mathbb{C}$.

Theorem 2.4 (Abel). *If for some $z_1 \neq z_0$ the series*

$$(2.6) \quad \sum_{k=0}^{\infty} c_k (z_1 - z_0)^k$$

converges, then

$$(2.7) \quad \sum_{k=0}^{\infty} c_k (z - z_0)^k$$

converges absolutely and uniformly in any disk $\mathbb{D}(z_0, r)$ with $r < |z_1 - z_0|$.

Exercise 2.5. Prove Theorem 2.4 by reducing it to the corresponding fact about real series and using the completeness of $\mathbb{C}$.

Abel's theorem shows that the domain of convergence of a power series is always a disk; convergence may also extend to parts of the boundary. The largest r such that the series (2.7) converges for all $z \in \mathbb{D}(z_0, r)$ is called the *radius of convergence*.

The disk of convergence can be degenerate: in one extreme case, it reduces to the point $z = z_0$ (radius of convergence 0), while in the other extreme it covers all of $\mathbb{C}$ ("infinite radius of convergence").

3. CONTINUITY AND DIFFERENTIABILITY

Definition 3.6. A complex function f is said to be continuous at z_0 if

$$f(z) \rightarrow f(z_0) \quad \text{as } z \rightarrow z_0.$$

Exercise 3.7. Show that polynomials are continuous in $\mathbb{C}$.

We can now define differentiability.

Definition 3.8. A function f is differentiable at z_0 if there exists a number $f'(z_0)$ such that

$$\frac{f(z) - f(z_0)}{z - z_0} \rightarrow f'(z_0) \quad \text{as } z \rightarrow z_0.$$

Exercise 3.9. Show that differentiation in $\mathbb{C}$ satisfies the familiar rules from real analysis. In particular, the sum rule, product rule, and chain rule all hold. (Prove this by mimicking the real-variable arguments.)

Also prove that differentiability implies continuity.

Differentiability in $\mathbb{C}$ is much more restrictive than in $\mathbb{R}$. For the same reason, complex differentiable functions enjoy far stronger regularity properties than merely real-differentiable functions.

Exercise 3.10. Show that the function $f(z) = |z|$ is **nowhere differentiable** in $\mathbb{C}$.

Definition 3.11. Let $\mathcal{D}$ be a domain in $\mathbb{C}$. A function f is said to be analytic in $\mathcal{D}$ if it is differentiable at every point $z \in \mathcal{D}$.

Functions analytic in a domain are also called holomorphic. The two terms are synonymous in this course.

If A is a set that contains a domain $\mathcal{D}$ together with parts of its boundary, we say that f is analytic in A if there exists a domain $\mathcal{D}' \supset A$ such that f is analytic in $\mathcal{D}'$.

We will see that if f is analytic in some domain $\mathcal{D}$, then f' is also analytic there. In particular, f has continuous derivatives of all orders in $\mathcal{D}$. This stems from the crucial fact that in complex analysis the derivative has an integral representation. We will also later establish that analyticity in a domain $\mathcal{D}$ is equivalent to the convergence of the Taylor series of f at every $z_0 \in \mathcal{D}$.

Clearly, in $\mathbb{R}$ differentiability does not imply C^∞ (infinite differentiability).

3.1. Differentiability of power series.

Theorem 3.12. *Let*

$$(3.2) \quad S(z) = \sum_{k=0}^{\infty} c_k (z - z_0)^k$$

be a power series convergent in the open disk $\mathbb{D}(z_0, r)$ with $r > 0$. Then S has derivatives of all orders in $\mathbb{D}(z_0, r)$. In particular,

$$(3.3) \quad S'(z) = \sum_{k=0}^{\infty} k c_k (z - z_0)^{k-1},$$

$$(3.4) \quad S''(z) = \sum_{k=0}^{\infty} k(k-1) c_k (z - z_0)^{k-2},$$

$\vdots$

$$(3.5) \quad S^{(p)}(z) = \sum_{k=0}^{\infty} k(k-1) \cdots (k-p+1) c_k (z - z_0)^{k-p},$$

$\vdots$

and each of these series converges in $\mathbb{D}(z_0, r)$ to the corresponding derivative of S .

Remark 3.13. *The derivative of the sum of a convergent power series S is obtained by differentiating each term individually. In other words, S can be differentiated term by term.*

Proof. It suffices to prove the case $p = 1$, since higher derivatives then follow by induction. By translation, assume without loss of generality that $z_0 = 0$.

Fix $|z| < \rho < r$, and choose h small enough so that $|z| + |h| < \rho$. Consider the partial sums

$$S_N(z) = \sum_{k=0}^N c_k z^k, \quad \frac{S_N(z+h) - S_N(z)}{h}.$$

For **real, positive** z, h , the binomial formula and Taylor's theorem with remainder give

$$(3.6) \quad (z+h)^n - z^n - nz^{n-1}h = \frac{n(n-1)}{2}z^{n-2}h^2 + \sum_{k=3}^n \binom{n}{k} z^{n-k}h^k$$

$$\leq \frac{n(n-1)}{2}(z+a)^{n-2}h^2 \leq \frac{n(n-1)}{2}\rho^{n-2}h^2,$$

for some $a \in (0, h)$. The **equality** remains valid for complex z, h . Thus,

$$\left| \frac{(z+h)^n - z^n}{h} - nz^{n-1} \right| = \left| \frac{n(n-1)}{2}z^{n-2}h + \sum_{k=3}^n \binom{n}{k} z^{n-k}h^k \right|$$

$$\leq \frac{n(n-1)}{2}|z|^{n-2}|h| + \sum_{k=3}^n \binom{n}{k}|z|^{n-k}|h|^k$$

$$\leq \frac{n(n-1)}{2}\rho^{n-2}|h|.$$

Therefore,

$$\left| \frac{S_N(z+h) - S_N(z)}{h} - S'_N(z) \right| \leq |h| \sum_{k=0}^N \frac{k(k-1)}{2} |c_k| \rho^k.$$

The series

$$C = \sum_{k=0}^{\infty} \frac{k(k-1)}{2} |c_k| \rho^k$$

converges because $\rho < r$. Indeed, since the original power series $\sum c_k z^k$ has radius of convergence r , the series $\sum |c_k| \rho^k$ converges absolutely for every $\rho < r$. Multiplying the coefficients by the polynomial factor $k(k-1)/2$ does not affect convergence (apply, for instance, the n th root test).

Hence the right-hand side is bounded by $C|h|$, which tends to 0 as $h \rightarrow 0$.

Thus $\lim_{h \rightarrow 0} \frac{S(z+h) - S(z)}{h}$ exists and equals $S'(z) = \sum_{k=0}^{\infty} k c_k z^{k-1}$, as claimed. ■

Exercise 3.14. Show that

$$\frac{(z+h)^n - z^n}{h} = \int_0^1 (z+th)^{n-1} dt.$$

Use the dominated convergence theorem, the triangle inequality, and the fact that $\lim_{n \rightarrow \infty} n^{1/n} = 1$ to derive a shorter proof of Theorem 3.12.

Corollary 3.15. *Assume S is a power series convergent in the disk $\mathbb{D}(z_0, r)$ with $r > 0$. Then S is analytic (holomorphic) in $\mathbb{D}(z_0, r)$.*

Corollary 3.16. *For the series (3.2), we have*

$$S^{(k)}(z_0) = k! c_k.$$

Hence (3.2) is precisely the Taylor series of S centered at z_0 .

Corollary 3.17. *Suppose $S(z)$ converges in a disk $\mathbb{D}(z_0, r)$, and there exists a sequence $\{z_n\}_{n \in \mathbb{N}}$ with an accumulation point at z_0 such that $S(z_n) = 0$ for all n . Then $S(z) \equiv 0$.*

Remark 3.18. *Corollary 3.17 is a crucial ingredient in the important principle of permanence of relations.*

Proof. Without loss of generality, assume $z_0 = 0$. Write

$$S(z) = c_0 + zT(z),$$

where $T(z)$ is a power series convergent in $\mathbb{D}(0, r)$.

By assumption,

$$0 = S(z_n) = c_0 + z_n T(z_n), \quad n \in \mathbb{N}.$$

Letting $n \rightarrow \infty$ shows that $c_0 = 0$.

Now divide by z to obtain $S(z)/z = T(z) = c_1 + zT_1(z)$, where $T_1(z)$ is again a power series convergent in $\mathbb{D}(0, r)$. Since $T(z_n) = 0$ for infinitely many $z_n \rightarrow 0$, the same reasoning shows $c_1 = 0$.

Repeating the argument inductively, we conclude that all coefficients c_k vanish. Hence $S \equiv 0$. ■

3.2. Some basic functions. ◦ *The exponential.* We define

$$(3.7) \quad e^z = \sum_{k=0}^{\infty} \frac{z^k}{k!}.$$

This series converges for every $z \in \mathbb{C}$, hence e^z is analytic in $\mathbb{C}$ by Theorem 3.12. Differentiating term by term (see (3.5)) yields

$$(3.8) \quad (e^z)' = e^z.$$

In particular,

$$(3.9) \quad (e^z e^{-z})' = 0,$$

so $e^z e^{-z}$ is constant throughout $\mathbb{C}$. Evaluating at $z = 0$ shows $e^0 = 1$, hence

$$(3.10) \quad e^z e^{-z} = 1 \quad \Leftrightarrow \quad e^{-z} = \frac{1}{e^z}.$$

Similarly, for any $a \in \mathbb{C}$ we compute

$$(3.11) \quad (e^{z+a}e^{-z})' = (e^{z+a})'e^{-z} + e^{z+a}(e^{-z})' = 0 \\ \implies e^{z+a}e^{-z} = e^ae^0 = e^a \implies e^{z+a} = e^ze^a,$$

which is the fundamental exponential identity.

For real $\phi \in \mathbb{R}$, one immediately checks Euler's formula:

$$(3.12) \quad e^{i\phi} = \cos \phi + i \sin \phi.$$

Remark 3.19. Euler's formula shows that the mapping $\phi \mapsto e^{i\phi}$ parametrizes the unit circle in the complex plane. As ϕ varies over $\mathbb{R}$, the point $e^{i\phi}$ traces the circle $|z| = 1$ counterclockwise, starting at 1 when $\phi = 0$. Thus the exponential provides a natural bridge between trigonometry and complex geometry.

Exercise 3.20. (a) In the above arguments we tacitly used an additional fact. Identify it and fill in the missing details using results already established.

(b) Define $\cos$ and $\sin$ in $\mathbb{C}$ by their power series and prove that (3.12) extends to all $z \in \mathbb{C}$:

$$e^{iz} = \cos z + i \sin z.$$

Deduce the important exponential representations

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i}, \quad \cos z = \frac{e^{iz} + e^{-iz}}{2}.$$

Use these to show that $\sin^2 z + \cos^2 z = 1$ for all $z \in \mathbb{C}$.

Exercise 3.21 (A first form of permanence of analytic relations). (a) Let f be analytic in a domain $\mathcal{D}$, and let F be analytic in a domain $\mathcal{D}'$ with $\mathcal{D}' \supset f(\mathcal{D})$. Suppose $\mathcal{D}$ contains a real interval and that

$$F(f(x)) = 0 \quad \text{for all real } x \text{ in that interval.}$$

Show that $F(f(z)) = 0$ for all $z \in \mathcal{D}$.

(b) Use part (a) to give an alternative proof of the fundamental exponential identities, relying on Corollary 3.17.

(c) Use part (a), together with the power series extensions of $\sin$ and $\cos$, to prove that

$$\sin^2 z + \cos^2 z = 1 \quad \text{for all } z \in \mathbb{C}.$$

Exercise 3.22. Show that

$$(3.13) \quad e^s = 1 \iff s = 2N\pi i, \quad N \in \mathbb{Z}$$

◦ *The logarithm.* In the complex domain, the logarithm is a more delicate function. For the moment, we consider the simpler case of defining $\log(1+z)$ for $|z| < 1$. This is done via the convergent Taylor series

$$(3.14) \quad \log(1+z) = z - \frac{z^2}{2} + \frac{z^3}{3} - \frac{z^4}{4} + \cdots.$$

By Theorem 3.12, term-by-term differentiation yields

$$(3.15) \quad \frac{d}{dz} \log(1+z) = 1 - z + z^2 - z^3 + \cdots = \frac{1}{1+z}, \quad |z| < 1.$$

Exercise 3.23. Show that if $|s|$ is small then

$$\log(e^s) = s, \quad e^{\log(1+s)} = 1 + s.$$

We will return later to the more general problem of defining $\log z$ for arbitrary $z \neq 0$. Its natural maximal domain of definition is not a subset of $\mathbb{C}$ but rather a *Riemann surface*, which we will study in detail later. The logarithm is one of the fundamental examples of a *branched function*, and in fact many other multivalued functions in complex analysis inherit their branching behavior from it.

3.3. Operations with power series. If S and T are power series convergent in a neighborhood of $z_0 = 0$, then the following combinations are also convergent in some (possibly smaller) neighborhood of 0:

$$S+T, \quad S \cdot T, \quad \frac{S}{T} \text{ (provided } T(0) \neq 0), \quad S \circ T \text{ (provided } T(0) = 0).$$

If S and T are power series convergent in a neighborhood of $z_0 = 0$, then the combinations

$$S + T, \quad S \cdot T, \quad \frac{S}{T} \text{ (if } T(0) \neq 0), \quad S \circ T \text{ (if } T(0) = 0)$$

are also convergent in some neighborhood of 0. Formulas for the coefficients of these new series are obtained by observing that power series are uniform limits of polynomials and that the operations agree with those for polynomials.

For example,

$$(3.16) \quad ST = s_0t_0 + (s_1t_0 + s_0t_1)z + (s_2t_0 + s_1t_1 + s_0t_2)z^2 + \cdots$$

Exercise 3.24. (a) If S and T are two power series with radius of convergence r , show that ST has radius of convergence at least r .

(b) Write three terms of the series S/T if $T(0) \neq 0$.

(c)* Under the assumptions above, show that S/T has nonzero radius of convergence.

(d)* Replace the assumption in (b) by $T(0) = 0$. Show that in this case $S \circ T$ has nonzero radius of convergence.

Exercise 3.25. Formulate these statements for points other than $z_0 = 0$.

3.4. The Cauchy–Riemann equations. Analytic functions admit several equivalent characterizations. So far, we have defined them by requiring differentiability at every point of a domain (Definition 3.8). An alternative, equally fundamental characterization is given by the Cauchy–Riemann equations:

Theorem 3.26 (Cauchy–Riemann equations). (1) If $f = u + iv$ is analytic in a domain $\mathcal{D} \subset \mathbb{C}$, then the partial derivatives exist and satisfy

$$(3.17) \quad \begin{aligned} u_x &= v_y, \\ u_y &= -v_x, \end{aligned}$$

throughout $\mathcal{D}$.

(2) Conversely, if u, v are differentiable functions in $\mathcal{D}$ that satisfy (3.17), then $f = u + iv$ is analytic in $\mathcal{D}$.

Proof. (1) Let $f(z) = u(x, y) + iv(x, y)$ and suppose $f'(z_0) = a + ib$. By translation we may assume $z_0 = 0$. Then

$$(3.18) \quad \begin{aligned} f(z) - f(0) &= u(x, y) - u(0, 0) + i(v(x, y) - v(0, 0)) \\ &= [f'(0) + \varepsilon(z)]z = (a + ib)(x + iy) + \varepsilon(z)z, \end{aligned}$$

where $\varepsilon(z) \rightarrow 0$ as $z \rightarrow 0$. This implies the existence of u_x, u_y, v_x, v_y at $(0, 0)$ and the validity of the Cauchy–Riemann equations there.

(2) Conversely, if u, v are differentiable at $(0, 0)$, then

$$(3.19) \quad \begin{aligned} f(z) - f(0) &= u(x, y) - u(0, 0) + i(v(x, y) - v(0, 0)) \\ &= u_x(0, 0)x + u_y(0, 0)y + i(v_x(0, 0)x + v_y(0, 0)y) + \varepsilon(x, y)x + i\eta(x, y)y, \end{aligned}$$

where $\varepsilon, \eta \rightarrow 0$ as $z \rightarrow 0$. Comparing (3.18) and (3.19) shows that f is differentiable iff (3.17) holds. ■

Exercise 3.27. Show that (3.18) and (3.19) are compatible if and only if (3.17) holds. (Hint: equate the real and imaginary parts and note that x and y are independent variables.)

3.5. Analyticity at infinity. As $|z| \rightarrow \infty$, we have $1/z \rightarrow 0$. By definition, a function f is said to be *analytic at infinity* if the function $f(1/z)$ is analytic at $z = 0$.

4. INTEGRALS

Integration plays an important role in complex analysis. As we shall see, the derivative of a function can be written as an integral, and many of the nice properties of analytic functions originate in this fact.

If $f(t) = u(t) + iv(t)$ is a complex-valued function of one real variable t , then

$$(4.2) \quad \int_a^b f(t) dt = \int_a^b u(t) dt + i \int_a^b v(t) dt.$$

Thus questions of complex integration reduce to familiar real integration.

Note 4.28. *In the following, unless otherwise specified, all curves are assumed to be piecewise differentiable.*

Let $\gamma(t) = x(t) + iy(t)$, $t \in [a, b]$, be a piecewise C^1 parametrized curve. We define

$$\int_{\gamma} f(z) dz := \int_a^b f(\gamma(t)) \gamma'(t) dt,$$

in the same way as in (4.2), by breaking γ , γ' , and f into their real and imaginary parts. (Do this calculation explicitly.)

The value of the integral depends on the *orientation* of γ , specified by stating that t increases from a to b . It is natural to say that $\gamma(a)$ is the *starting point* and $\gamma(b)$ the *endpoint*.

The same geometric curve with the opposite orientation is denoted $-\gamma$. A convenient formula is

$$(-\gamma)(t) = x(a + b - t) + i y(a + b - t), \quad t \in [a, b].$$

It follows that

$$(4.3) \quad \int_{-\gamma} f(z) dz = \int_b^a f(\gamma(t)) \gamma'(t) dt = - \int_{\gamma} f(z) dz.$$

If $\gamma(a) = \gamma(b)$ the curve is called *closed*. Unless otherwise specified, positive orientation (counterclockwise) is assumed, in which case we write

$$\oint_{\gamma} := \int_{\gamma}.$$

Exercise 4.29. *Show that*

$$(4.4) \quad \int_{\gamma} f(z) dz = \int_{\gamma} (u dx - v dy) + i \int_{\gamma} (u dy + v dx).$$

Positively oriented γ (counterclockwise) Reversed orientation $-\gamma$ (clockwise)




FIGURE 1. Orientation of a closed curve: γ counterclockwise (left) and $-\gamma$ clockwise (right).

Exercise 4.30. Show that the integral along a given geometric curve depends on its parametrization only through the orientation (i.e. reparametrizations preserving orientation leave the integral unchanged, while reversing orientation multiplies it by -1).

A curve is called *simple* if it has no self-intersections. For example, the circle is simple, but the figure “8” is not.

A domain $\mathcal{D}$ is called *simply connected* if any simple closed curve γ contained in $\mathcal{D}$ can be deformed continuously to a point through curves completely contained in $\mathcal{D}$. Formally, this means:

There exists a continuous function of two variables

$$F(t, s) = x(t, s) + iy(t, s), \quad (t, s) \in [a, b] \times [0, 1],$$

with values in $\mathcal{D}$ such that $F(t, 0) = \gamma(t)$ and $F(t, 1) = p$ for some point $p \in \mathcal{D}$.

Intuitively, a simply connected domain has no holes. For example, a disk is simply connected, but a punctured disk (such as $\mathbb{D} \setminus \{0\}$) or an *annulus*

$$\{z \in \mathbb{C} : r < |z| < R\}$$

are not simply connected.

Theorem 4.31 (Cauchy). Assume $\mathcal{D}$ is a simply connected domain and that f is continuously differentiable in $\mathcal{D}$. If γ is a piecewise differentiable simple closed curve contained in $\mathcal{D}$, then

$$(4.5) \quad \oint_{\gamma} f(z) dz = 0.$$

Proof. Start from the decomposition (4.4) and apply Green's theorem:

$$(4.6) \quad \int_{\gamma} (u \, dx - v \, dy) = - \iint_{\text{Int}(\gamma)} \left(\frac{\partial v}{\partial x} + \frac{\partial u}{\partial y} \right) dx \, dy.$$

This vanishes by the Cauchy–Riemann equations (3.17). The second integral in (4.4) is handled in the same way. ■

Cauchy originally assumed continuity of f' was required. Later, Montel and others weakened this hypothesis, requiring only boundedness, L^1 , or other mild conditions. Finally, Goursat (1884) observed that mere differentiability suffices.

Theorem 4.32 (Cauchy–Goursat). *Assume $\mathcal{D}$ is a simply connected domain and that f is differentiable in $\mathcal{D}$. If γ is a piecewise differentiable simple closed curve contained in $\mathcal{D}$, then*

$$(4.7) \quad \oint_{\gamma} f(z) \, dz = 0.$$

Proof. By continuity of f and piecewise differentiability of γ , the integral $\int_{\gamma} f(s) \, ds$ can be approximated with arbitrary accuracy by integrals along polygonal curves P , i.e. $\int_P f(s) \, ds$. Decomposing the polygon into triangles, it suffices to show that

$$\int_T f(s) \, ds = 0$$

for any triangle $T \subset \mathcal{D}$.




FIGURE 2. Orientation of integration after the first subdivision (left); further partitions (right).

Split T into four smaller triangles and write $\int_T = \sum_{j=1}^4 \int_{T_j}$, choosing orientations so that integrals along inner sides cancel. Let T_{11} denote

the subtriangle such that

$$\left| \int_{T_{11}} f(s) ds \right| = \max_j \left\{ \left| \int_{T_j} f(s) ds \right| \right\}.$$

Then

$$\left| \int_T f(s) ds \right| \leq 4 \left| \int_{T_{11}} f(s) ds \right|.$$

Repeating inductively, we obtain nested closed triangles T_{nn} with

$$\left| \int_T f(s) ds \right| \leq 4^n \left| \int_{T_{nn}} f(s) ds \right|, \quad n \in \mathbb{N}.$$

Since $\{T_{nn}\}$ is a nested sequence of compact sets, their intersection reduces to a point $z_0 \in T$.

Now for any fixed n and $z \in \partial T_{nn}$, by differentiability, we have

$$f(z) = f(z_0) + f'(z_0)(z - z_0) + \varepsilon(z; z_0)(z - z_0),$$

with $\varepsilon(z)$ as $z \rightarrow z_0$. Since $\oint_{T_{nn}} f(z_0) dz = \oint_{T_{nn}} f'(z_0)(z - z_0) dz = 0$ (by Cauchy's theorem), it follows that

$$\int_{T_{nn}} f(s) ds = \int_{T_{nn}} \varepsilon(s; z_0)(s - z_0) ds.$$

Estimating the diameter of T_{nn} by $C2^{-n}$ for some constant C (why can we choose C independent of n ?), we conclude

$$\left| \int_T f(s) ds \right| \leq C^2 4^n 2^{-n} 2^{-n} \sup_{T_{nn}} |\varepsilon(s, t)| \rightarrow 0 \quad (n \rightarrow \infty).$$

■

Exercise 4.33. *Why is the supremum above tending to zero?*

It is sometimes useful to integrate analytic functions along the boundary of their domain of analyticity. For instance, if f is continuous up to the boundary, Cauchy's theorem still holds:

Exercise 4.34. *Assume that f is analytic in $\mathcal{D}$, continuous on $\overline{\mathcal{D}}$, and that $\partial\mathcal{D}$ is a rectifiable curve of winding number one. Show that (4.5) holds if γ is a simple closed curve in $\overline{\mathcal{D}}$.*

5. CAUCHY'S FORMULA

5.1. Homotopic curves. Let $\mathcal{D}$ be a domain in $\mathbb{C}$. Two curves in $\mathcal{D}$ are said to be *homotopic in $\mathcal{D}$* if one can be continuously deformed into the other through curves lying entirely in $\mathcal{D}$ (cf. the discussion preceding Theorem 4.31).

For example, if $\mathcal{D}$ is simply connected then any simple closed curve is homotopic to a point (see Fig. 3). As another example, if $\mathcal{D}$ is the annulus

$$\{z \in \mathbb{C} : 1 < |z| < 2\},$$

then all circles $|z| = r$ with $1 < r < 2$ are homotopic to each other, but not to a point. On the other hand, any simple closed curve that does not wind around 0 is homotopic to a point.




FIGURE 3. All dotted curves inside γ are homotopic to each other and to the central point.

It is also useful to consider curves γ_1, γ_2 in $\mathcal{D}$, given by functions $\gamma_1(t), \gamma_2(t)$ for $t \in [a, b]$, which have the same endpoints, $\gamma_1(a) = \gamma_2(a)$ and $\gamma_1(b) = \gamma_2(b)$. Two such curves are called *homotopic with fixed endpoints* if they can be continuously deformed into each other through a transformation that preserves the endpoints and remains entirely within $\mathcal{D}$ (see Fig. 4).

5.2. Independence of the integral on the path. Line integrals are additive with respect to concatenation of curves. Let γ_1 be an oriented curve, and let γ_2 start at the final point of γ_1 . If $\gamma_1(t)$ is defined for $t \in [a, b]$ and $\gamma_2(t)$ for $t \in [b, c]$ with $\gamma_1(b) = \gamma_2(b)$, then the concatenated curve $\gamma_1 + \gamma_2$ runs from $t = a$ to $t = c$. By definition,

$$(5.2) \quad \int_{\gamma_1 + \gamma_2} f(z) dz = \int_{\gamma_1} f(z) dz + \int_{\gamma_2} f(z) dz.$$




FIGURE 4. Two curves γ_1 and γ_2 homotopic with fixed endpoints.

(i) Suppose f is analytic in a simply connected domain $\mathcal{D}$, and let γ_1, γ_2 be two oriented curves in $\mathcal{D}$ with the same endpoints. Then $\gamma = \gamma_1 - \gamma_2$ is a closed curve. If there exists a subdomain $\mathcal{D}' \subset \mathcal{D}$ containing γ such that γ is homotopic to a point in $\mathcal{D}'$, then by (4.5), (5.2), and (4.3) we obtain

$$0 = \int_{\gamma_1 - \gamma_2} f(z) dz = \int_{\gamma_1} f(z) dz - \int_{\gamma_2} f(z) dz.$$

Therefore, *the integral of an analytic function on a simply connected domain depends only on the endpoints of the path, provided the paths are homotopic:*

$$(5.3) \quad \int_{\gamma_1} f(z) dz = \int_{\gamma_2} f(z) dz.$$




FIGURE 5. In the yellow domain, γ_1 and γ_2 are not homotopic, while γ_3 and γ_4 are.

5.3. Cauchy's Formula. Let $\mathcal{D}$ be a domain in $\mathbb{C}$ and $z_0 \in \mathcal{D}$. The functions $(z - z_0)^{-n}$, $n = 1, 2, \dots$ are analytic in $\mathcal{D} \setminus \{z_0\}$. Thus, if γ_1 and γ_2 are two closed curves in $\mathcal{D}$ not passing through z_0 , and homotopic to each-other in $\mathcal{D} \setminus \{z_0\}$ then

$$(5.4) \quad \int_{\gamma_1} (z - z_0)^{-n} dz = \int_{\gamma_2} (z - z_0)^{-n} dz$$

Clearly, these integrals are zero if γ_i does not contain z_0 inside. To calculate the integrals on a simple closed curve encircling z_0 it suffices, by (5.4), to do the calculation when the curve is a circle, which can be done explicitly. Indeed, a circle centered at z_0 with radius ρ is parametrized by $z = z_0 + \rho e^{it}$, $t \in [0, 2\pi]$ (where we used Euler's formula), and we get

$$(5.5) \quad \oint \frac{dz}{(z - z_0)^n} = \frac{i}{\rho^{n-1}} \int_0^{2\pi} e^{-i(n-1)t} dt = \begin{cases} 2\pi i & \text{if } n = 1 \\ 0 & \text{otherwise} \end{cases}$$

Definition 5.35. If γ is a curve and $z_0 \notin \gamma$, then the index of z_0 w.r.t. γ is defined as

$$\text{Ind}_\gamma(z_0) = \frac{1}{2\pi i} \oint_\gamma \frac{d\zeta}{\zeta - z_0}$$

and it represents the “number of times γ winds around z_0 ”.

Theorem 5.36 (Cauchy's formula). *If f is analytic in a simply connected domain $\mathcal{D}$ and γ is a piecewise differentiable simple closed curve in $\mathcal{D}$ enclosing z , then*

$$(5.6) \quad f(z) = \frac{1}{2\pi i} \oint_\gamma \frac{f(s)}{s - z} ds.$$

Proof. Write

$$f(s) - f(z) = f'(z)(s - z) + \varepsilon(s)(s - z),$$

where $\varepsilon(s) \rightarrow 0$ as $s \rightarrow z$. Note that ε is continuous in $\mathcal{D} \setminus \{z\}$. Choose $\rho > 0$ small enough that $\mathbb{D} := \mathbb{D}_\rho(z) \subset \mathcal{D}$, and fix $\delta > 0$ with $|\varepsilon(s)| < \delta/(2\pi)$ for $s \in \partial\mathbb{D}$. Then

$$(5.7) \quad \begin{aligned} \frac{1}{2\pi i} \oint_\gamma \frac{f(s)}{s - z} ds &= \frac{1}{2\pi i} \oint_{\partial\mathbb{D}} \frac{f(s)}{s - z} ds \\ &= \frac{f(z)}{2\pi i} \oint_{\partial\mathbb{D}} \frac{ds}{s - z} + \frac{f'(z)}{2\pi i} \oint_{\partial\mathbb{D}} ds + \frac{1}{2\pi i} \oint_{\partial\mathbb{D}} \varepsilon(s) ds. \end{aligned}$$

By (5.5), the first integral equals $f(z)$, the second vanishes, and the last has absolute value $\leq \delta$. Letting $\delta \rightarrow 0$ gives the result. ■

6. TAYLOR SERIES OF ANALYTIC FUNCTIONS

Theorem 6.37 (Taylor series; Cauchy's formula for higher derivatives). *If f is analytic in a domain $\mathcal{D}$ and $z_0 \in \mathcal{D}$, then there exists $\rho > 0$ such that for all $z \in \mathbb{D}(z_0; \rho)$*

$$(6.2) \quad f(z) = \sum_{k=0}^{\infty} c_k (z - z_0)^k,$$

where

$$(6.3) \quad c_k = \frac{1}{2\pi i} \oint_{\partial \mathbb{D}_\rho(z_0)} \frac{f(s)}{(s - z_0)^{k+1}} ds = \frac{f^{(k)}(z_0)}{k!}.$$

Proof. Assume f is analytic in $\mathcal{D}$ and fix $z_0 \in \mathcal{D}$. Choose $\rho > 0$ small enough so that $\overline{\mathbb{D}_\rho(z_0)} \subset \mathcal{D}$. By Theorem 5.36, for $z \in \mathbb{D}_\rho(z_0)$

$$(6.4) \quad f(z) = \frac{1}{2\pi i} \oint_{\partial \mathbb{D}_\rho(z_0)} \frac{f(s)}{s - z} ds.$$

Expand the kernel:

$$(6.5) \quad \frac{1}{s - z} = \frac{1}{s - z_0 - (z - z_0)} = \frac{1}{s - z_0} \sum_{k=0}^{\infty} \left(\frac{z - z_0}{s - z_0} \right)^k,$$

which converges absolutely for $|z - z_0| < |s - z_0| = \rho$. Since the series is bounded by

$$\frac{1}{|s - z_0|} \sum_{k=0}^{\infty} \left| \frac{z - z_0}{s - z_0} \right|^k,$$

the dominated convergence theorem allows us to interchange sum and integral, yielding

$$(6.6) \quad f(z) = \sum_{k=0}^{\infty} c_k (z - z_0)^k, \quad c_k = \frac{1}{2\pi i} \oint_{\partial \mathbb{D}_\rho(z_0)} \frac{f(s)}{(s - z_0)^{k+1}} ds.$$

■

Remark 6.38. *This theorem shows the central equivalence of complex analysis: analytic functions are exactly those that admit a convergent power series expansion in a neighborhood of every point of their domain. In contrast with real analysis, where differentiability need not imply such an expansion, holomorphic functions are “rigid”—their local behavior is completely determined by their power series.*

Exercise 6.39 (Cauchy remainder). *Show that*

$$(6.7) \quad f(z) = \sum_{k=0}^n c_k (z - z_0)^k + E(z, z_0, n),$$

where

$$(6.8) \quad E(z, z_0, n) = \frac{1}{2\pi i} \oint_{\partial \mathbb{D}_\rho(z_0)} \frac{f(s)}{s - z} \frac{(z - z_0)^{n+1}}{(s - z_0)^{n+1}} ds.$$

Prove that, if $\overline{\mathbb{D}_\rho(z_0)} \subset \mathcal{D}$, then

$$(6.9) \quad |E(z, z_0, n)| \leq \max_{\mathbb{D}_\rho(z_0)} |f| \frac{|z - z_0|^{n+1}}{\rho - |z - z_0|}.$$

If $\mathbb{D}_\rho(z_0)$ is the maximal disk centered at z_0 contained in $\mathcal{D}$, then the optimal estimate is

$$(6.10) \quad |E(z, z_0, n)| \leq \inf_{\rho' \leq \rho} \sup_{\mathbb{D}_{\rho'}(z_0)} |f| \frac{|z - z_0|^{n+1}}{\rho' - |z - z_0|}.$$

Theorem 6.40. If f is analytic in a domain $\mathcal{D}$ and $z_0 \in \mathcal{D}$, then f has derivatives of all orders at z_0 . In particular, $f', f'', \dots$ are analytic in $\mathcal{D}$.

Remark 6.41. The disk of convergence of the Taylor series of an analytic function cannot be *zero*. In fact, its radius of convergence equals the distance from z_0 to the nearest singularity of f (see Fig. 6).




FIGURE 6. Disk of convergence of a Taylor series. The yellow region is a domain of analyticity and the red dot a singularity.

Indeed, in any smaller disk Theorem 6.37 applies. If the radius of convergence were larger than the distance to the nearest singularity, f would extend analytically beyond its given domain (*why?*), contradicting maximality.

Example. Consider $f(z) = \frac{1}{1+z^2}$. Its Taylor series at $z = 0$ is

$$\frac{1}{1+z^2} = \sum_{k=0}^{\infty} (-1)^k z^{2k}, \quad |z| < 1.$$

The singularities at $z = \pm i$ lie exactly on the boundary of the disk of convergence.

Theorem 6.42 (Liouville). *An entire function (analytic in all of $\mathbb{C}$) that is bounded on $\mathbb{C}$ must be constant.*

Proof. Let $M = \sup_{\mathbb{C}} |f|$. By Theorem 6.37, for any $\rho > 0$,

$$(6.11) \quad f'(z) = \frac{1}{2\pi i} \oint_{\partial \mathbb{D}_{\rho}(0)} \frac{f(s)}{(s-z)^2} ds.$$

Therefore

$$(6.12) \quad |f'(z)| \leq \frac{1}{2\pi} M \frac{1}{\rho^2} (2\pi\rho) = \frac{M}{\rho}.$$

Letting $\rho \rightarrow \infty$ yields $f'(z) \equiv 0$, so f is constant. ■

Exercise 6.43. Show that an entire function that is not a polynomial must grow faster than any power of $|z|$ along some path as $z \rightarrow \infty$.

7. THE FUNDAMENTAL THEOREM OF ALGEBRA

Exercise 7.44. Let P be a polynomial of degree $n \geq 1$. Show that there exists $R > 0$ such that $1/P$ is analytic in the domain $\{z : |z| > R\}$ and that for some nonzero c , $z^n/P(z) \rightarrow c$ as $|z| \rightarrow \infty$.

One classical application of Liouville's Theorem is the Fundamental Theorem of Algebra:

Theorem 7.45 (Fundamental Theorem of Algebra). A polynomial $P_n(z)$ of degree $n \geq 1$ has exactly n roots in $\mathbb{C}$, counting multiplicity.

Proof. It suffices to show the existence of one root when $n \geq 1$, since the general statement then follows inductively by factoring.

By Exercise 7.44, if P_n had no zeros, then $1/P_n$ would be entire and vanish at infinity. But then $1/P_n$ would be bounded (why?) and entire, hence constant by Liouville's theorem. This is impossible, so P_n must have at least one zero. ■

8. MORE PROPERTIES OF ANALYTIC FUNCTIONS

Assume f is analytic in $\mathbb{D}_\rho(z_0)$ and that all derivatives of f vanish at z_0 . Then $f \equiv 0$ in $\mathbb{D}_\rho(z_0)$ (check!). In fact, more is true:

Proposition 8.46. *If f is analytic in a domain $\mathcal{D}$ and all derivatives at some $z_0 \in \mathcal{D}$ vanish, then f is identically zero in $\mathcal{D}$.*

Proof. For any $z \in \mathcal{D}$ there is a polygonal path P from z_0 to z , composed of segments $[z_{j-1}, z_j]$, $j = 1, \dots, n$, with disks $\mathbb{D}_{r_j}(z_j) \subset \mathcal{D}$ covering each step (see Proposition 48.19).

By simple geometry, we may choose $\rho > 0$ such that every point along P is contained in a disk of radius ρ lying in $\mathcal{D}$. Cover P by a finite chain of such overlapping disks, starting with one centered at z_0 .

Since f vanishes identically in the first disk, all its derivatives vanish at the center of the second disk, so f vanishes there too. Repeating this argument along the chain of disks, we conclude $f(z) = 0$ for all $z \in \mathcal{D}$. ■

Theorem 8.47 (Morera). *Let f be continuous in a simply connected domain $\mathcal{D}$ and assume that*

$$\oint_{\gamma} f(z) dz = 0$$

for every simple piecewise C^1 closed curve $\gamma \subset \mathcal{D}$. Then f is analytic in $\mathcal{D}$. (The same conclusion holds when the curves γ are boundaries of triangles; see Exercise 8.48 below.)

Proof. Fix $z_0 \in \mathcal{D}$ and define

$$F(z) = \int_{z_0}^z f(s) ds,$$

where the path of integration is any piecewise C^1 path in $\mathcal{D}$. By the assumption $\oint_{\gamma} f = 0$, this definition is path-independent.

Let δ be small and choose a path γ' from z_0 to $z + \delta$. Then

$$\int_{\gamma} f(s) ds - \int_{\gamma'} f(s) ds = \int_z^{z+\delta} f(s) ds,$$

where the last integral can be taken along a straight line. Writing $f(s) = f(z) + \varepsilon(s)$ with $\varepsilon(s) \rightarrow 0$ as $s \rightarrow z$, we obtain

$$\int_z^{z+\delta} f(s) ds = f(z)\delta + o(\delta).$$

Thus F is differentiable with $F'(z) = f(z)$. Since F is analytic, so is f . The restriction to triangles follows from polygonal connectivity and is left as an exercise. ■

Exercise 8.48 (Restriction to triangles). (a) Prove Morera's theorem assuming $\oint_{\gamma} f = 0$ only for triangular paths, using polygonal-path connectedness. (b) Give an alternative proof based on approximation of curves by polygonal lines.

We now have three equivalent viewpoints on analytic functions in a domain $\mathcal{D}$:

- as functions differentiable with respect in $\mathcal{D}$,
- as sums of convergent power series centered at any point in $\mathcal{D}$,
- as continuous functions with vanishing integrals along closed loops.

Each characterization is fundamental and useful.

Theorem 8.49 (Weierstrass). Let f_n be analytic in a domain Ω and suppose $f_n \rightarrow f$ uniformly on compact subsets of Ω . Then f is analytic in Ω . Moreover, $f'_n \rightarrow f'$ uniformly on compact subsets of Ω .

Note 8.50. By iterating, it follows that $f_n^{(k)} \rightarrow f^{(k)}$ uniformly on compact sets for each $k \in \mathbb{N}$.

Proof. Let T be a triangle contained in a compact $K \subset \Omega$. Since f_n is analytic,

$$\int_T f_n(z) dz = 0.$$

By uniform convergence, f is continuous on K and dominated convergence applies, giving

$$\int_T f(z) dz = 0.$$

By Morera's theorem, f is analytic. The convergence of the derivatives follows directly from Cauchy's integral formula and dominated convergence. ■

9. HARMONIC FUNCTIONS

A real-valued C^2 function $u(x, y)$ satisfying Laplace's equation

$$(9.2) \quad u_{xx} + u_{yy} = 0$$

in a domain U is called *harmonic* in U .

Theorem 9.51. *A function u is harmonic in a simply connected domain $\mathcal{D} \subset \mathbb{C}$ if and only if it is the real (or imaginary) part of some analytic function f in $\mathcal{D}$. The function f is unique up to an additive purely imaginary constant.*

In other words: for each harmonic u there exists a harmonic conjugate v (unique up to a constant) such that $u + iv$ is analytic.

Proof. If $u = \operatorname{Re}(f)$ for some analytic f , then $f = u + iv$ satisfies the Cauchy–Riemann equations

$$u_x = v_y, \quad u_y = -v_x.$$

Differentiating once more, we obtain

$$u_{xx} = v_{yx}, \quad u_{yy} = -v_{xy}.$$

Since $v_{xy} = v_{yx}$, it follows that $u_{xx} + u_{yy} = 0$, i.e. u is harmonic.

Conversely, suppose u is harmonic in $\mathcal{D}$. Define the vector field

$$\mathbf{E} = (-u_y, u_x).$$

By the harmonicity condition $u_{xx} + u_{yy} = 0$, one checks that

$$\frac{\partial}{\partial x}(u_x) = \frac{\partial}{\partial y}(-u_y),$$

which shows that $\mathbf{E}$ is curl-free. Since $\mathcal{D}$ is simply connected, $\mathbf{E}$ is conservative, so there exists a potential v (unique up to an additive constant) such that

$$v_x = -u_y, \quad v_y = u_x.$$

These are exactly the Cauchy–Riemann equations for $u + iv$, so $f = u + iv$ is analytic in $\mathcal{D}$. ■

Exercise 9.52. *Assume f is analytic in a domain $\mathcal{D}$ and write $f = u + iv$ with $u = \operatorname{Re} f$ and $v = \operatorname{Im} f$ on $\mathcal{D}$. Show that*

$$(9.3) \quad \begin{pmatrix} u_x & u_y \\ v_x & v_y \end{pmatrix} = \begin{pmatrix} \operatorname{Re} f'(z) & -\operatorname{Im} f'(z) \\ \operatorname{Im} f'(z) & \operatorname{Re} f'(z) \end{pmatrix}.$$

Hint: See the section on the C-R equations.

Lemma 9.53. *Let $f = u + iv$ be analytic near z_0 with $f'(z_0) \neq 0$. Then the constant level curves $u(x, y) = u(x_0, y_0)$ and $v(x, y) = v(x_0, y_0)$ exist near z_0 , are smooth, and intersect orthogonally.*

Proof. Left as an exercise, since the result follows immediately from (9.3). ■

9.1. Potential and Hamiltonian flows. Consider an autonomous system of ODEs in $\mathcal{D}$:

$$\dot{x} = E_1(x, p), \quad \dot{p} = E_2(x, p).$$

The system is *Hamiltonian* if there exists $H \in C^1(\mathcal{D})$ such that

$$E_1 = \frac{\partial H}{\partial p}, \quad E_2 = -\frac{\partial H}{\partial x}.$$

It is a *potential system* if there exists $V \in C^1(\mathcal{D})$ with

$$E_1 = \frac{\partial V}{\partial x}, \quad E_2 = \frac{\partial V}{\partial p}.$$

A system is simultaneously potential and Hamiltonian precisely when there exist H and V such that

$$\frac{\partial H}{\partial p} = \frac{\partial V}{\partial x}, \quad \frac{\partial H}{\partial x} = -\frac{\partial V}{\partial p}.$$

Thus H and V must be harmonic conjugates, i.e. the real and imaginary parts of an analytic function.

In Hamiltonian systems, H is conserved:

$$\frac{d}{dt}H(x(t), p(t)) = H_x \dot{x} + H_p \dot{p} = 0,$$

so H remains constant along trajectories. In gradient systems, $(\dot{x}, \dot{p})$ points in the direction of steepest ascent of V .

Note 9.54. *The steepest descent/ascent lines of a harmonic function u satisfy the ODE system*

$$\dot{x} = v_x, \quad \dot{y} = v_y,$$

where v is the harmonic conjugate. The solution curves are smooth wherever $v_x^2 + v_y^2 \neq 0$. This viewpoint will be important in the method of steepest descent.

10. THE MAXIMUM MODULUS PRINCIPLE

One of the most striking features of analytic functions is that their absolute value cannot attain an interior maximum unless the function is constant:

Theorem 10.55 (Maximum Modulus Principle). *If f is analytic and nonconstant in a domain $\mathcal{D}$, then $|f|$ cannot achieve a maximum at any point of $\mathcal{D}$.*

In other words, any maximum of $|f|$, if achieved, must occur on the boundary of $\mathcal{D}$.

Proof. Assume, for contradiction, that $z_0 \in \mathcal{D}$ is a point where $|f|$ attains its maximum. Since f is nonconstant, it cannot be identically zero. Let $M = \max_{\mathcal{D}} |f| = |f(z_0)| > 0$. By scaling, we may assume without loss of generality that $M = 1$ and $z_0 = 0$.

If $f \equiv 1$ identically, this contradicts the assumption that f is non-constant. Otherwise, in the Taylor expansion of f about 0,

$$f(z) = 1 + c_k z^k + c_{k+1} z^{k+1} + \cdots,$$

let $k \geq 1$ be the smallest index with $c_k = |c_k|e^{i\theta} \neq 0$. Factor

$$(10.2) \quad f(z) = 1 + c_k z^k (1 + d_k z E_1(z)),$$

where E_1 is analytic near 0 and bounded.

Choose $z_1 = |z_1|e^{-i\theta}$ small enough so that $|z_1 d_k E_1(z_1)| < 1$. Then $c_k z_1^k = |c_k||z_1|^k \in \mathbb{R}^+$. By the triangle inequality,

$$|f(z_1)| = |1 + c_k z_1^k (1 + d_k z_1 E_1(z_1))| > 1,$$

contradicting the maximality of $|f|$ at 0.

Therefore f must be constant. ■

Exercise 10.56. Show that if $|f|$ has a minimum in $\mathcal{D}$, then this minimum must be 0.

Exercise 10.57. * Find the maximum and minimum values of $|\sin z|$ inside the closed unit disk.

Steepest ascent and descent lines. Let f be analytic near z_0 , and let $k \geq 1$ be the smallest index such that $c_k \neq 0$ in the Taylor expansion

$$f(z) = f(z_0) + c_k (z - z_0)^k + \cdots.$$

Definition 10.58. Let $f(z) = \sum_{j=0}^{\infty} c_j (z - z_0)^j$ with $k \geq 1$ the smallest index such that $c_k \neq 0$. The direction of $d \in \mathbb{C}$ is called a steepest ascent direction for $|f|$ at z_0 if $c_k d^k \in \mathbb{R}^+$, and a steepest descent direction if $c_k d^k \in \mathbb{R}^-$.

A piecewise differentiable oriented curve γ is called a steepest ascent (descent) line for f if, at every point of differentiability $z \in \gamma$, its tangent vector is a steepest ascent (descent) direction for $|f|$ at z .

Note 10.59. By Note 9.54, the lines of steepest ascent and descent are smooth and unique through any z_0 such that

$$f(z) = f(z_0) + f'(z_0)(z - z_0) + \cdots, \quad f'(z_0) \neq 0.$$

A **saddle point** is a point where $f'(z_0) = 0$. Then f is locally of the form $z^k + \cdots$ with $k \geq 2$. One checks that there are exactly k




FIGURE 7. A soap film (image from the google images, <https://math.hmc.edu/~jacobsen/demolab/soap-film-2/>)

distinct directions of steepest ascent at such a point. Consequently, lines of steepest descent can be continued, in k possible ways, through a saddle point. These notions are central in the *steepest descent method* (also called the *saddle point method*).

Theorem 10.60. *Assume u is harmonic and nonconstant in a domain $\mathcal{D}$. Then u has no maximum and no minimum in $\mathcal{D}$.*

Proof. Let $u = \operatorname{Re}(f)$ and define $g = e^f$, which is analytic in $\mathcal{D}$. Writing $e^f = e^u e^{iv}$, we see that $|e^f| = e^u$. Thus u has a maximum if and only if $|g|$ has a maximum. But by the maximum modulus principle, $|g|$ cannot attain a maximum strictly inside $\mathcal{D}$. For a minimum, note that $\min(u) = -\max(-u)$, which reduces to the previous case. ■

10.1. Application: soap films. A soap film picked up by a thin closed wire assumes the surface of *least area* compatible with the constraint that it is bounded by the wire (since the potential energy is proportional to the surface area).

If the wire is close to planar (say lying near the (x, y) -plane), the local height $u(x, y)$ of the film satisfies Laplace's equation. (This can be shown by elementary differential geometry.)²

If the wire is exactly planar, then $u = 0$ on the boundary, and by Theorem 10.60, the minimal surface is flat. This is unsurprising.

²If the film can be written locally as a graph $z = u(x, y)$ over a planar domain $\Omega \subset \mathbb{R}^2$, then u solves the *minimal surface equation*

$$(10.3) \quad \operatorname{div} \left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right) = 0 \quad \text{in } \Omega, \quad u|_{\partial\Omega} = u_{\partial\Omega},$$

where the boundary data $u_{\partial\Omega}$ encodes the wire (after projection to the plane).

More generally, we will be able to solve Laplace's equation with arbitrary boundary conditions, yielding insight into minimal surfaces. Conversely, the intuition we have about soap films translates directly to solutions of Laplace's equation: for example, a soap film has no local extremum, since any local flattening would reduce the surface area.

10.2. Cauchy principal value integrals (PV). Suppose f is analytic in a domain containing the simple closed, piecewise differentiable curve C . By Cauchy's theorem we have

$$(10.4) \quad \frac{1}{2\pi i} \oint_C \frac{f(s)}{s-z} ds = \begin{cases} f(z), & z \text{ inside } C, \\ 0, & z \text{ outside } C. \end{cases}$$

What if z lies *on* C ? Then the integral requires interpretation. One definition is to excise from C a symmetric segment of length ε centered at z , and then let $\varepsilon \rightarrow 0$. This yields the *Cauchy principal value integral*, denoted $\text{PV} \oint$. Another equivalent definition is to take the half-sum of the integrals along curves circumventing z from the inside and from the outside.

Exercise 10.61. Show that if C is a smooth closed curve and f is analytic in a neighborhood of C , then

$$(10.5) \quad \frac{1}{2\pi i} \text{PV} \oint_C \frac{f(s)}{s-z} ds = \frac{1}{2} f(z),$$

and that the two definitions above coincide. In particular, if $f(s)/(s-z)$ extends analytically across z , then the PV integral agrees with the usual one.

Definition 10.62 (Cauchy PV on the line). If for every small $\varepsilon > 0$ one has $f \in L^1((a, L-\varepsilon)) \cap L^1((L+\varepsilon, b))$, then the principal value is defined by

$$(10.6) \quad \text{PV} \int_a^b f(s) ds := \lim_{\varepsilon \downarrow 0} \left(\int_a^{L-\varepsilon} f(s) ds + \int_{L+\varepsilon}^b f(s) ds \right),$$

provided the limit exists.

Example 10.63. The *Hilbert transform*, an important operator in analysis and applications, is defined by

$$(10.7) \quad H(u)(t) = \frac{1}{\pi} \text{PV} \int_{-\infty}^{+\infty} \frac{u(\tau)}{t-\tau} d\tau.$$

It is bounded on $L^p(\mathbb{R})$ for $1 < p < \infty$, where the limit exists pointwise almost everywhere, and maps L^1 to weak L^1 . With some Hölder

regularity, the limit exists everywhere. The Hilbert transform is a central tool in solving Riemann–Hilbert problems. *to which we will return later.*

Exercise 10.64. *Show that if $f \in L^1(a, b)$ then*

$$(10.8) \quad \text{PV} \int_a^b f(s) ds = \int_a^b f(s) ds.$$

10.3. Bi-analytic bijections (automorphisms) of $\mathbb{D}$.

Exercise 10.65. *Let $a \in \mathbb{C}$ with $|a| < 1$ and $\theta \in \mathbb{R}$. Show that*

$$(10.9) \quad A(z) := e^{i\theta} \frac{a + z}{1 + \bar{a}z}$$

is an automorphism of $\mathbb{D}$.

Remark 10.66. *We will prove later (using Schwarz’s Lemma) that every automorphism of $\mathbb{D}$ is of the form (10.9) for some $a \in \mathbb{C}$ with $|a| < 1$ and $\theta \in \mathbb{R}$.*

11. POISSON’S FORMULA

A function u is said to be harmonic in an open set $\Omega \subset \mathbb{R}^n$ if it is of class C^2 in Ω and satisfies Laplace’s equation

$$\Delta u = 0 \quad \text{in } \Omega.$$

The **mean value property** for harmonic functions states that if $B(x, r)$ is a ball with center x and radius r completely contained in Ω , then the value of u at the center of the ball is given by the average value of u :

$$u(x) = \frac{1}{n\omega_n r^{n-1}} \int_{\partial B(x, r)} u d\sigma = \frac{1}{\omega_n r^n} \int_{B(x, r)} u dV,$$

where ω_n is the volume of the unit ball in $\mathbb{R}^n$ and $d\sigma$ denotes the $(n - 1)$ -dimensional surface measure.

We shall first prove this property in the case $n = 2$. Clearly, we can change variables so that $x = 0$ and $r = 1$. In complex notation, the statement becomes:

Proposition 11.67 (Mean value property for harmonic functions). *Assume u is harmonic in the open unit disk and continuous on the closed unit disk. Then*

$$(11.2) \quad u(0) = \frac{1}{2\pi} \int_0^{2\pi} u(e^{it}) dt.$$

Proof. If v is a harmonic conjugate of u , then $f := u + iv$ is analytic in the open unit disk. By Cauchy's integral formula, for any $\rho < 1$ we have

$$(11.3) \quad \begin{aligned} u(0) + iv(0) &= f(0) = \frac{1}{2\pi i} \oint_{\partial \mathbb{D}_\rho(0)} \frac{f(s)}{s} ds = \frac{1}{2\pi} \int_0^{2\pi} f(\rho e^{it}) dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} u(\rho e^{it}) dt + i \frac{1}{2\pi} \int_0^{2\pi} v(\rho e^{it}) dt. \end{aligned}$$

Taking the real part and letting $\rho \rightarrow 1^-$ yields (11.2). $\blacksquare$

Exercise 11.68. (i) Let u be as in Proposition 11.67 and let A be an automorphism of the disk, as in Exercise 10.65. Show that

$$U(z) = u(A(z))$$

is harmonic in the open unit disk and continuous on the closed unit disk. (*Hint: Write the Laplacian as $\Delta = 4\partial_z\partial_{\bar{z}}$.*)

(ii) Show that, if $z_0 = re^{i\theta}$, then

$$(11.4) \quad u(z_0) = \frac{1}{2\pi} \int_0^{2\pi} u\left(e^{i\theta} \frac{r + e^{is}}{1 + re^{is}}\right) ds = \frac{1}{2\pi} \int_{-\pi}^{\pi} u\left(e^{i\theta} \frac{r + e^{is}}{1 + re^{is}}\right) ds.$$

(*Hint for (ii): Use (i) with $A(w) = e^{i\theta} \frac{r + w}{1 + rw}$, so that $A(0) = z_0$, and apply Proposition 11.67 to $U = u \circ A$.)*

Proposition 11.69 (Poisson's formula). Let $f(e^{it})$ be continuous on $[-\pi, \pi)$. Define

$$(11.5) \quad u(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \frac{1 - r^2}{1 - 2r \cos(t - \theta) + r^2} f(e^{it}) dt.$$

Then u is harmonic in $\mathbb{D}$, extends continuously to $\bar{\mathbb{D}}$, and satisfies $u(e^{i\theta}) = f(e^{i\theta})$.

Moreover, the **Poisson kernel**

$$P_r(\alpha) := \frac{1 - r^2}{1 - 2r \cos \alpha + r^2}$$

satisfies $P_r(\alpha) \geq 0$ and

$$\frac{1}{2\pi} \int_0^{2\pi} P_r(\alpha) d\alpha = 1.$$

Proof. Harmonicity follows either from a direct computation $\Delta u = 0$.

To conclude the proof of the first part, it remains to show that u extends continuously to $\partial\mathbb{D}$ and that

$$\lim_{r \rightarrow 1^-} u(re^{i\theta}) = f(e^{i\theta}).$$

Fix θ and, by rotation, assume $\theta = 0$. Then

$$u(r) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{1-r^2}{1-2r\cos t+r^2} f(e^{it}) dt.$$

Split the integral at $|t| = \delta$. For $|t| > \delta$,

$$0 \leq \frac{1-r^2}{1-2r\cos t+r^2} \leq \frac{(1+r)(1-r)}{2r(1-\cos\delta)} \rightarrow 0 \quad (r \rightarrow 1^-),$$

so by dominated convergence this part tends to 0.

For $|t| \leq \delta$, write $\cos t = 1 - \frac{t^2}{2}\phi(t)$ with $\phi(t) \rightarrow 1$. Then

$$\frac{1-r^2}{1-2r\cos t+r^2} = \frac{(1+r)(1-r)}{(1-r)^2 + rt^2\phi(t)}.$$

Setting $t = (1-r)\tau$ gives

$$u(r) = \frac{1}{2\pi} \int_{-\frac{\delta}{1-r}}^{\frac{\delta}{1-r}} \frac{1+r}{1+r\tau^2\phi((1-r)\tau)} f(e^{i(1-r)\tau}) d\tau,$$

and dominated convergence (check the details here) yields

$$\lim_{r \rightarrow 1^-} u(r) = \frac{2}{2\pi} f(1) \int_{-\infty}^{\infty} \frac{d\tau}{1+\tau^2} = f(1).$$

Thus u extends continuously to $\partial\mathbb{D}$ with boundary values f . The positivity and normalization properties of the Poisson kernel are straightforward to verify. ■

11.1. The Dirichlet problem for the Laplacian in $\mathbb{D}$. Formula (11.5) provides the solution of Laplace's equation in two dimensions with Dirichlet boundary conditions, namely with u prescribed on the boundary, when the domain is $\mathbb{D}$. A simple change of variables adapts this formula to any disk. More generally, the formula can be extended to apply to a general simply connected domain bounded by a simple, closed, piecewise differentiable curve. This follows from the *Riemann mapping theorem*.

Example 11.70 (The Faraday cage). “A region surrounded by a conductor does not feel the electrical influence of static outside charges.” We illustrate this phenomenon in two dimensions.

At equilibrium, the electric potential along a conductor is constant. Indeed, if two points of the conductor were at different potentials, there would exist a potential difference, hence an electric current $I = V/R$ (with R the resistivity). This would contradict equilibrium. This is, of course, a standard physics argument, though not a mathematically rigorous one. A full mathematical treatment would require setting up the precise PDE model and proving existence and uniqueness results, which lies beyond the scope of our course.

Thus we consider Laplace's equation (24.2) with boundary condition $V = C$ on $\partial\mathcal{D}$. Since the constant function $V = C$ is a solution, uniqueness of solutions implies that it is *the* solution. Hence

$$\mathbf{E} = -\nabla V = 0,$$

which is the desired conclusion.

12. ISOLATED SINGULARITIES, LAURENT SERIES

Definition 12.71. A function f has an isolated singularity at z_0 if f is analytic in a punctured disk $\mathbb{D}_\rho(z_0) \setminus \{z_0\}$ for some $\rho > 0$.

More generally, we will analyze functions analytic in annuli

$$\mathbb{D}_\rho(z_0) \setminus \mathbb{D}_{\rho'}(z_0), \quad 0 < \rho' < \rho.$$

For example, the functions $e^{1/z}$ and $1/\sin z$ have isolated singularities at $z = 0$, whereas the singularity of $\ln z$ at 0 is not isolated (we will see that $\ln$ is not well defined in $\mathbb{D} \setminus \{0\}$).

Definition 12.72. An isolated singularity z_0 of f is classified as follows:

- (1) a **removable singularity** if f extends to a function $\tilde{f}$ analytic in the whole disk. By a slight abuse of notation, we typically do not distinguish $\tilde{f}$ from f itself;
- (2) a **pole of order** $M \geq 1$ if $(z - z_0)^M f$ has a removable singularity at z_0 , but $(z - z_0)^{M-1} f$ does not;
- (3) an **essential singularity** if it is neither removable nor a pole.

A function all of whose singularities are poles is called **meromorphic**.

In the following, we denote the punctured disk $\mathbb{D}_\rho(z_0) \setminus \{z_0\}$ by $\mathbb{D}_\rho^*(z_0)$.

Proposition 12.73 (Laurent series). If f is analytic in $\mathbb{D}_\rho(z_0) \setminus \mathbb{D}_{\rho'}(z_0)$, where $0 < \rho' < \rho$, then f has a convergent representation

$$(12.2) \quad f(z) = \sum_{k \in \mathbb{Z}} a_k (z - z_0)^k, \quad z \in \mathbb{D}_\rho(z_0) \setminus \mathbb{D}_{\rho'}(z_0),$$




FIGURE 8. Circles of integration and cut in the proof of the Laurent series expansion.

where

$$a_k = \frac{1}{2\pi i} \oint \frac{f(s)}{(s - z_0)^{k+1}} ds, \quad k \in \mathbb{Z},$$

with the integral taken along any circle centered at z_0 with radius between ρ' and ρ .

Proof. Without loss of generality, take $z_0 = 0$. Consider the annulus between two circles C_o and C_i in $A = \mathbb{D}_\rho \setminus \mathbb{D}_{\rho'}$, as in Fig. 8 (C_o is the outer boundary and C_i the inner one). Introduce a cut L in the annulus as shown, so that $A \setminus L$ is simply connected. By Cauchy's integral formula, for $z \in A \setminus L$ we have

$$(12.3) \quad f(z) = \frac{1}{2\pi i} \oint_{C_o} \frac{f(s)}{s - z} ds - \frac{1}{2\pi i} \oint_{C_i} \frac{f(s)}{s - z} ds.$$

Here we used the fact that the boundary of the cut annulus is the closed path composed of C_o (counterclockwise), followed by the cut segment ℓ , then C_i (clockwise), and finally $-\ell$. The rest of the argument parallels the proof of Taylor's theorem and is left as an exercise. ■

Exercise 12.74. Complete the proof of formula (12.2) by expanding the integrands in (12.3) into power series in z/s and s/z , respectively, and estimating the remainders as we did in deriving formula (6.4).

Note. Check that (12.3) provides a decomposition of f into two parts: a function f_1 , analytic in the disk enclosed by C_o , and a function f_2 ,

analytic in $1/z$ in the exterior of the disk bounded by C_i . In [3] this decomposition is used to give an elegant proof of (12.2). For example, $e^{1/z}$ has an essential singularity at $z = 0$. Application of (12.2) yields

$$(12.4) \quad e^{1/z} = \sum_{k=0}^{\infty} \frac{1}{k!} z^{-k}.$$

(Of course, this formula could also be obtained directly from the power series of the exponential by replacing z with z^{-1} .)

Note. The part of the Laurent series consisting of the terms with negative powers is called the *principal part* of the series.

Note. Laurent series are of great theoretical value. However, they are not practical for *computing* the values of functions near an essential singularity, since the convergence becomes slower as the singularity is approached.

Theorem 12.75 (Casorati–Weierstrass; also known as the “little Picard theorem”). *Assume f is analytic in $\mathbb{D}_{\rho}^*(z_0)$ and has an essential singularity at z_0 . Then, for any $\rho_1 < \rho$, the set $f(\mathbb{D}_{\rho_1}^*(z_0))$ is dense in $\mathbb{C}$.*

Proof. Without loss of generality, assume $z_0 = 0$. Suppose, for contradiction, that there exists $\zeta \in \mathbb{C}$ and $\rho_1 > 0$ such that

$$|f(z) - \zeta| > c > 0 \quad \text{for all } z \in \mathbb{D}_{\rho_1}^*(0).$$

Then

$$h(z) = \frac{1}{f(z) - \zeta}$$

is bounded in $\mathbb{D}_{\rho_1}^*(0)$, and hence 0 is a removable singularity of h (why?). But then

$$f(z) = \zeta + \frac{1}{h(z)}$$

would have either a removable singularity or a pole at 0 (why?), contradicting the assumption that 0 is an essential singularity. ■

13. LAURENT SERIES AND FOURIER SERIES

Let f be 2π -periodic and assume:

Assumption. The Fourier coefficients $\{c_k\}_{k \in \mathbb{Z}}$ of f satisfy $\{c_k\} \in \ell^1(\mathbb{Z})^3$.

³ $\ell^1(\mathbb{Z})$ denotes the space of sequences $\{c_k\}_{k \in \mathbb{Z}}$ which are absolutely summable, i.e. $\sum_{k \in \mathbb{Z}} |c_k| < \infty$.

Then

$$g(z) := \sum_{k=0}^{\infty} c_k z^k$$

converges absolutely and uniformly in $\overline{\mathbb{D}}$ (why?), hence g is analytic in $\mathbb{D}$ and continuous up to $\partial\mathbb{D}$ (why?). For all $k \leq -1$ we then have

$$\oint_{\partial\mathbb{D}} \frac{g(s)}{s^{k+1}} ds = 0.$$

Similarly, define

$$H(\zeta) := \sum_{k=1}^{\infty} c_{-k} \zeta^k.$$

Then H is analytic in $\mathbb{D}$ and continuous up to $\overline{\mathbb{D}}$. Thus

$$h(z) := H\left(\frac{1}{z}\right)$$

is analytic in $\text{ext}(\overline{\mathbb{D}})$ and continuous down to $\partial\mathbb{D}$. For all $k \geq 0$ one checks that

$$\oint_{\partial\mathbb{D}} \frac{h(s)}{s^{k+1}} ds = 0.$$

Hence, for $s \in \mathbb{R}$,

$$f(s) = g(e^{is}) + h(e^{is}).$$

We have therefore proved the following result.

Proposition 13.76. *If f is 2π -periodic and its Fourier coefficients satisfy the assumption above, then f decomposes as the sum of two functions: one analytic in $\mathbb{D}$ and continuous in $\overline{\mathbb{D}}$, and another analytic in $\text{ext}(\overline{\mathbb{D}})$ and continuous in $\text{ext}(\mathbb{D})$.*

14. CALCULATING THE TAYLOR SERIES OF SIMPLE FUNCTIONS

One easy way to calculate Taylor series is to use operations with series; see §3.3.

Example. (1) The Taylor series of the function $z^{-1} \sin z$ is

$$(14.2) \quad \frac{\sin z}{z} = 1 - \frac{z^2}{6} + \frac{z^4}{120} - \cdots$$




FIGURE 9. Multiply connected domains

(2) The Taylor series of the function $z/\sin z$ is

$$\begin{aligned}
 (14.3) \quad \frac{z}{\sin z} &= \frac{1}{1 - \left(\frac{z^2}{6} - \frac{z^4}{120} + \cdots \right)} = 1 + \left(\frac{z^2}{6} - \frac{z^4}{120} + \cdots \right) \\
 &\quad + \left(\frac{z^2}{6} - \frac{z^4}{120} + \cdots \right)^2 + \cdots \\
 &= 1 + \frac{z^2}{6} - \frac{z^4}{120} + \frac{z^4}{36} + \cdots = 1 + \frac{z^2}{6} + \frac{7z^4}{360} + \cdots
 \end{aligned}$$

The first function is entire, while the second one is not. *Question:* What is the radius of convergence of the second series?

Exercise 14.77. Find the integral of $1/\cos z$ on a circle of radius $1/2$ centered at $z_0 = \pi/2$.

15. RESIDUES AND INTEGRALS

Definition 15.78. A function f is meromorphic in a neighborhood of a point z_0 if it is analytic in some punctured domain $\mathcal{D} \setminus \{z_0\}$ and there exists $k \geq 0$ such that $(z - z_0)^k f(z)$ has a removable singularity at z_0 .

Note 15.79. Meromorphic functions have one-sided convergent Laurent expansions in $\mathcal{D} \setminus \{z_0\}$, in the sense that only finitely many Laurent coefficients of negative order are nonzero.

Definition 15.80 (Residues). Let f be meromorphic in $\mathcal{D} \setminus \{z_0\}$. The coefficient c_{-1} in its Laurent expansion at z_0 is called the **residue** of f at z_0 , denoted $\text{Res}(f, z_0)$. If $\overline{\mathbb{D}}_\rho(z_0) \subset \mathcal{D}$, then

$$\frac{1}{2\pi i} \oint_{\partial \overline{\mathbb{D}}_\rho(z_0)} f(s) ds = c_{-1},$$

hence the terminology “residue.”

Proposition 15.81. *Let $\mathcal{D}$ be a simply connected domain. Suppose f is analytic in $\mathcal{D} \setminus \bigcup_{k=1}^n \mathbb{D}_k$, where $\mathbb{D}_k$ are disjoint disks centered at $z_k \in \mathcal{D}$. Let γ be a simple closed, piecewise differentiable curve which encircles each $\mathbb{D}_k$ once (see Definition 19.96 and Figure 15). Then*

$$(15.2) \quad \oint_{\gamma} f(s) ds = 2\pi i \sum_{k=1}^n \text{Res}(f, z_k).$$

Exercise 15.82. *Prove Proposition 17.89 by deforming and cutting the curve of integration appropriately.*

Example. Calculate

$$\oint \frac{dz}{\sin^3 z}$$

on a circle of radius $1/2$ around the origin.

Solution. In $\mathbb{D}(0, 1/2)$ we have

$$(15.3) \quad \frac{1}{\sin^3 z} = \frac{1}{\left(z - \frac{z^3}{6} + \frac{z^5}{120} - \dots\right)^3} = \frac{1}{z^3} \frac{1}{\left(1 - \frac{z^2}{6} + \frac{z^4}{120} - \dots\right)^3} \\ = \frac{1}{z^3} \left(1 + \frac{z^2}{2} + \frac{17z^4}{120} + \dots\right).$$

Thus the residue of $\sin^{-3}(z)$ at $z = 0$ is $1/2$, and the integral equals πi .

Exercise 15.83. *Show that if f has a pole of order m at $z = z_i$, then*

$$(15.4) \quad \text{Res}(f, z_i) = \frac{1}{(m-1)!} \left[\frac{d^{m-1}}{dz^{m-1}} \left((z - z_i)^m f(z) \right) \right]_{z=z_i}$$

by applying Laurent's formula near $z = z_i$.

Note 15.84 (Computational tool). *Formula (17.4) is the standard method for computing residues at poles of finite order. It reduces the problem to taking a finite number of derivatives of an analytic function. In practice, this formula, together with the method used in 15.3, are often the main “go-to” tools for residue computations; for low order poles the latter method is usually faster.*

16. CALCULATING THE TAYLOR SERIES OF SIMPLE FUNCTIONS

One easy way to calculate Taylor series is to use operations with series; see §3.3.

Example.

(1) The Taylor series of the function $z^{-1} \sin z$ is

$$(16.2) \quad \frac{\sin z}{z} = 1 - \frac{z^2}{6} + \frac{z^4}{120} - \dots$$

(2) The Taylor series of the function $z/\sin z$ is

$$\begin{aligned}
 (16.3) \quad \frac{z}{\sin z} &= \frac{1}{1 - (z^2/6 - z^4/120 + \cdots)} \\
 &= 1 + (z^2/6 - z^4/120 + \cdots) + (z^2/6 - z^4/120 + \cdots)^2 + \cdots \\
 &= 1 + \frac{z^2}{6} - \frac{z^4}{120} + \frac{z^4}{36} + \cdots = 1 + \frac{z^2}{6} + \frac{7z^4}{360} + \cdots
 \end{aligned}$$

The first function is entire; the second one is not. *Question:* What is the radius of convergence of the second series?⁴

Exercise 16.85. Find the integral of $1/\cos z$ on a circle of radius $1/2$ centered at $z_0 = \pi/2$.




FIGURE 10. Multiply connected domains

17. RESIDUES AND INTEGRALS

Definition 17.86. A function f is meromorphic in a neighborhood of a point z_0 if it is analytic in some punctured domain $\mathcal{D} \setminus \{z_0\}$ and there is a $k \geq 0$ so that $(z - z_0)^k f$ has a removable singularity at z_0 .

Note 17.87. Meromorphic functions have one-sided convergent Laurent expansions in $\mathcal{D} \setminus \{z_0\}$, in the sense that only a finite number of Laurent coefficients of negative order are nonzero (why?).

Definition 17.88 (Residues). Let f be meromorphic in $\mathcal{D} \setminus \{z_0\}$. The coefficient c_{-1} in its Laurent expansion at z_0 is called the **residue** of f at z_0 , denoted $\text{Res}(f, z_0)$. If $\overline{\mathbb{D}}_\rho(z_0) \subset \mathcal{D}$, then

$$\frac{1}{2\pi i} \oint_{\partial \mathbb{D}_\rho(z_0)} f(s) ds = c_{-1},$$

hence the name “residue.”

⁴Hint: the radius of convergence is determined by the distance from 0 to the nearest pole of $z/\sin z$.

Proposition 17.89. *Let $\mathcal{D}$ be a simply connected domain. Suppose f is analytic in $\mathcal{D} \setminus \bigcup_{k=1}^n \mathbb{D}_k$, where $\mathbb{D}_k$ are disks centered at $z_k \in \mathcal{D}$. Let γ be a simple closed, piecewise differentiable curve which encircles each $\mathbb{D}_k$ once (see Definition 19.96 and Figure 15). Then*

$$(17.2) \quad \oint_{\gamma} f(s) ds = 2\pi i \sum_{k=1}^n \text{Res}(f, z_k).$$

Exercise 17.90. *Prove Proposition 17.89 by deforming and cutting the curve of integration appropriately.*

Example. Calculate

$$\oint \frac{dz}{\sin^3 z}$$

on a circle of radius $1/2$ around the origin.

Solution. In $\mathbb{D}(0, 1/2)$ $\sin$ vanishes only at the origin. We have

$$(17.3) \quad \frac{1}{\sin^3 z} = \frac{1}{\left(z - \frac{z^3}{6} + \frac{z^5}{120} - \dots\right)^3} = \frac{1}{z^3} \frac{1}{\left(1 - \frac{z^2}{6} + \frac{z^4}{120} - \dots\right)^3} \\ = \frac{1}{z^3} \left(1 + \frac{z^2}{2} + \frac{17z^4}{120} + \dots\right).$$

Thus the residue of $\sin^{-3}(z)$ at $z = 0$ is $1/2$, and the integral equals πi .

Exercise 17.91. *Show that if f has a pole of order m at $z = z_i$, then*

$$(17.4) \quad \text{Res}(f, z_i) = \frac{1}{(m-1)!} \left[\frac{d^{m-1}}{dz^{m-1}} \left((z - z_i)^m f(z) \right) \right]_{z=z_i}$$

by applying Laurent's formula near $z = z_i$.

18. INTEGRALS OF TRIGONOMETRIC FUNCTIONS

Contour integration is very useful in calculating or estimating Fourier coefficients of periodic functions. Consider the integral

$$(18.2) \quad I = \int_0^{2\pi} \frac{\cos(nt)}{2 + \cos t} dt = \text{Re}(J), \quad \text{where} \quad J := \int_0^{2\pi} \frac{e^{int}}{2 + \cos t} dt.$$

Let $z = e^{it}$. Then

$$(18.3) \quad J = -i \oint_C \frac{z^{n-1}}{2 + (z + 1/z)/2} dz = -2i \oint_C \frac{z^n}{z^2 + 4z + 1} dz,$$

where C is the unit circle. The roots of $z^2 + 4z + 1$ are $-2 \pm \sqrt{3}$, and only one, $z_0 = -2 + \sqrt{3}$, lies inside the unit disk. Thus (check)

$$(18.4) \quad J = -2i \cdot 2\pi i \frac{z_0^n}{2z_0 + 4} \Rightarrow I = \frac{4\pi z_0^n}{2z_0 + 4} = \frac{2\pi z_0^n}{\sqrt{3}}.$$

19. COUNTING ZEROS AND POLES

Notations and definitions. (1) Assume f is analytic in a disk $\mathbb{D}_\rho(z_0)$ and $f(z_0) = 0$. Then, in $\mathbb{D}_\rho(z_0)$,

$$(19.2) \quad f(z) = \sum_{k=1}^{\infty} c_k (z - z_0)^k.$$

If $f \not\equiv 0$, there exists some k_0 such that $c_{k_0} \neq 0$ (see Proposition 8.46). The smallest such k_0 is called the **order** (or **multiplicity**) of the zero z_0 . For a meromorphic function g (see Definition 12.72), the order of a pole at z_0 is the multiplicity of the zero of $1/g$ at z_0 .

Exercise 19.92 (The zeros of an analytic function are isolated). Assume $f \not\equiv 0$ is analytic near z_0 and $f(z_0) = 0$. Use the Taylor series expansion to show that there is some disk around z_0 in which

$$f(z) = 0 \Rightarrow z = z_0.$$

Now assume f is meromorphic in $\mathcal{D}$, and let γ be a simple, piecewise C^1 closed curve contained in $\mathcal{D}$ together with its interior Γ . Since $\gamma \cup \Gamma$ is a closed subset of $\mathcal{D}$, by definition, the region of analyticity of f strictly contains Γ . For the purpose of the next theorem, the assumptions can be relaxed, allowing γ to be the boundary of the analyticity domain of f , provided suitable continuity conditions are imposed on f and f' .

Theorem 19.93 (Counting zeros and poles). Let f be as above, and assume f has no zeros on γ . Let N be the total number of zeros of f in Γ , counting multiplicities, and let P be the total number of poles of f in Γ , each counted with its order. Then

$$(19.3) \quad \frac{1}{2\pi i} \oint_{\gamma} \frac{f'(s)}{f(s)} ds = N - P.$$

Proof. We leave some details to the reader. The function f'/f is meromorphic. At a zero of f of order n_i , f'/f has a simple pole with residue n_i ; at a pole of f of order p_i , f'/f has a simple pole with residue $-p_i$. The formula then follows immediately from the residue theorem (Proposition 17.89). ■

19.1. Constancy of the number of roots of $f(s) - \zeta$. Let γ and Γ be as before. Assume that f is analytic in $\mathcal{D} \supset \Gamma$.

Let V be the set of values of f attained in Γ but not on γ , that is,

$$V = \{ \zeta \in \mathbb{C} : \zeta \in f(\Gamma) \setminus f(\gamma) \}.$$

Theorem 19.94. Assume f is not constant and for each $\zeta \in V$, let $N(\zeta)$ be the number of roots of $f(z) - \zeta = 0$, counting multiplicity. Then $N(\zeta)$ is independent of ζ .

Hence, every value that f takes in Γ is attained the same finite number of times. This integer is called the **order** of f in Γ .

Proof. ■

For $\zeta \in V$ define

$$g(\zeta) = \frac{1}{2\pi i} \oint_{\gamma} \frac{(f(s) - \zeta)'}{f(s) - \zeta} ds = \frac{1}{2\pi i} \oint_{\gamma} \frac{f'(s)}{f(s) - \zeta} ds.$$

By Theorem 19.93, $g(\zeta)$ equals the number of zeros of $f(s) - \zeta$ in Γ (since there are no poles).

Moreover, g is continuous in $\zeta \in V$. But an integer-valued function is continuous on a connected set if and only if it is constant.

Remark 19.95. The integer $N(\zeta)$, independent of $\zeta \in V$, is what we called the *order* of f in Γ . In other contexts, especially in topology and the theory of Riemann surfaces, the same invariant is referred to as the *degree* of the holomorphic map f . It measures how many times f covers its image, counting multiplicity, and anticipates the notion of branched coverings.

Definition 19.96. The **winding number** of a point z_0 with respect to a closed, piecewise C^1 curve γ not passing through z_0 is defined by

$$(19.4) \quad \frac{1}{2\pi i} \oint_{\gamma} \frac{dz}{z - z_0}.$$

Exercise 19.97. Show that the winding number defined above is always an integer. (Hint: see §19.1.)

Remark 19.98. Geometrically, the winding number measures how many times the curve γ winds around the point z_0 , counted with orientation. A positively oriented simple closed curve enclosing z_0 has winding number 1, while if z_0 lies outside γ the winding number is 0. More complicated curves may yield larger positive or negative integers depending on how many times they encircle z_0 and in which direction.

19.2. Some further properties of the logarithm. If γ is a curve without self-intersections joining 1 to z and not passing through 0, then

$$\int_{\gamma} \frac{ds}{s} =: \log_{\gamma} z$$




FIGURE 11. Images under $(z - 1)(z - 1/2)$ of circles centered at 0 with radii $1/4$ (innermost), $3/4$ (middle), and $5/4$ (outer).

defines an analytic function in a neighborhood of γ . We may homotopically deform γ so that it begins with the interval $[1, 1 + \varepsilon]$ for some small $\varepsilon > 0$. Since for positive real x we have $e^{\log x} = x$, analyticity in a neighborhood of γ allows us to apply the permanence of relations. It follows that $e^{\log_\gamma s} = s$ for s near γ , and in particular $e^{\log_\gamma z} = z$. If we take another curve without self-intersections, $\tilde{\gamma}$ joining 1 to z , the same argument yields $e^{\log_{\tilde{\gamma}} z} = z$. We thus have

$$(19.5) \quad e^{\log_\gamma z} = e^{\log_{\tilde{\gamma}} z} \Rightarrow e^{\log_\gamma z - \log_{\tilde{\gamma}} z} = 1 \\ \Rightarrow \log_\gamma z - \log_{\tilde{\gamma}} z = 2N\pi i \quad \text{for some } N \in \mathbb{Z}$$

Thus, logarithms defined along different curves may differ only by an integer multiple of $2\pi i$. This multiple determines a particular *branch* of the log. Writing $z = \rho e^{i\phi}$ and $\log z = u + iv$, we obtain $e^u e^{iv} = \rho e^{i\phi}$, whence

$$\log z = \log |\rho| + i\psi,$$

where $\log |\rho|$ is the usual real log of a real positive number, and the angle $\psi = \phi + 2N\pi$ for some $N \in \mathbb{Z}$ is the *argument of z denoted by $\arg(z)$* .

19.3. The argument principle. Consider again the setting of Theorem 19.93. Take $g = \ln f$ for any particular branch of the log. We

have $g' = f'/f$ and then (19.3) shows that the change in $g = \ln f$ as we traverse positively γ is $N - P$. Since $\ln f = \ln |f| + 2\pi i \arg f$, and f is single-valued, $f(z)$ and $|f(z)|$ do not change as we traverse the curve. In other words,

$$(19.6) \quad \frac{1}{2\pi i} \oint_{\gamma} \frac{f'(s)}{f(s)} ds = \frac{1}{2\pi i} \oint_{f \circ \gamma} \frac{d\zeta}{\zeta} = N - P = \frac{1}{2\pi i} \Delta \log f = \frac{1}{2\pi i} \Delta \arg f.$$

Thus $2\pi i(N - P)$ is the **variation of the argument of f along γ** . Equivalently, it is the winding number of 0 with respect to $f \circ \gamma$ (see Definition 19.96), that is, the number of times — positive or negative — that $f \circ \gamma$ winds around 0.

Finally, this says that if we take the image of a parametrization of γ under f , then, $N - P$ counts the number of times the image $f(\gamma)$ turns around zero. More precisely, since f is not zero on the contour, we can define the log of f on γ as follows. Since the zeros of f' are isolated, there will be only finitely many of them on γ . We break the integral at the points where f' vanishes. Take a $z_0 \in \gamma$ such that $f'(z_0) \neq 0$ and choose any smooth curve γ' joining 1 and $f(z_0)$, which avoids zero. Define $\log f(z_0) = \int_{\gamma'} ds/s$. As discussed, the value of this integral depends on the path through a multiple of $2\pi i$ which will turn out to be immaterial. We take a small disk $\mathbb{D}_{\rho}(z_0)$ where $f' \neq 0$, and we note that, for $z \in \mathbb{D}_{\rho}(z_0)$, $\int_{f(z_0)}^{f(z)} ds/s = \int_{z_0}^z f'(u)/f(u) du := \log(f(z)) - \log(f(z_0))$. We note that these integrals are equal piecewise between any two successive points where $f' \neq 0$, thus everywhere along γ and you can check that this $\log f$ is well defined and analytic in the union of the disks. We then have

Exercise 19.99. Let $f(z) = \exp(1/z) - 1$. Clearly 0 is an essential singularity and Proposition 19.93 does not apply. Determine, as a function of $\varepsilon > 0$, the number of times the curve

$$\{f(\varepsilon e^{it}) : t \in [0, 2\pi)\}$$

winds around the origin.

19.4. Hurwitz's theorem. This theorem shows that a uniform limit of analytic functions with no zeros in a domain is either identically zero, or else an analytic function with no zeros in the domain:

Theorem 19.100 (Hurwitz). *Let $\{f_n\}_{n \in \mathbb{N}}$ be analytic and nonvanishing in a domain $\Omega \subset \mathbb{C}$, and assume that $\{f_n\}_{n \in \mathbb{N}}$ converges uniformly on compact sets to a function $f \not\equiv 0$. Then f has no zeros in Ω .*

Proof. By standard results, the uniform convergence of $\{f_n\}$ on compacta implies that f is analytic. Consider the neighborhood of a point

$z_0 \in \Omega$; by translation we may take $z_0 = 0$. Since $f \not\equiv 0$, there exists $\delta > 0$ such that $f(z) \neq 0$ for all $z \in \partial\mathbb{D}_\delta(0)$ (by Corollary 3.17).

Moreover, by Weierstrass's theorem (Theorem 8.49), the sequence $\{f'_n\}$ converges uniformly to f' . Hence $\{f'_n/f_n\}_{n \in \mathbb{N}}$ converges uniformly to f'/f on $\partial\mathbb{D}_{\delta/2}(0)$. Applying Theorem 19.93 to each f_n and passing to the limit shows that f has no zeros in $\mathbb{D}_{\delta/2}(0)$. Since z_0 was arbitrary, the conclusion holds throughout Ω . ■

19.5. Rouché's Theorem.

Theorem 19.101 (Rouché). *Assume f and h are analytic in a domain $\mathcal{D}$ containing the piecewise differentiable simple closed curve γ and its interior Γ . Suppose that on γ we have $|h| < |f|$. Then the number of zeros of f and of $f + h$ in Γ is the same (counting multiplicities).*

Note 19.102. *Often h is viewed as a perturbation of f . The theorem is quantitative: it does not merely assert that a “small enough” perturbation preserves the number of zeros, but gives the explicit sufficient condition $|h| < |f|$ on γ . Whenever this inequality holds, f and $f + h$ have the same number of zeros inside Γ .*

Proof. Since $|h| < |f|$ on γ , there is a neighborhood $\mathcal{N}$ of γ in which f has no zeros. In $\mathcal{N}$ we can write

$$(19.7) \quad f + h = f \cdot (1 + h/f) =: fQ,$$

where $Q = 1 + g$ with $g = h/f$. On $\mathcal{N}$ we have $|g| < 1$, so the series

$$q(z) = \log(1 + g(z)) = \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} g(z)^k$$

converges uniformly, and hence q is analytic in $\mathcal{N}$ by Theorem 8.49.

Differentiating gives

$$q' = \frac{g'}{1 + g} = \frac{Q'}{Q}.$$

Thus

$$\frac{(fQ)'}{fQ} = \frac{f'}{f} + \frac{Q'}{Q}.$$

But since Q is analytic and nonvanishing in $\mathcal{N}$, we have

$$\oint_{\gamma} \frac{Q'}{Q} dz = 0.$$

Therefore

$$\oint_{\gamma} \frac{(f + h)'}{f + h} dz = \oint_{\gamma} \frac{f'}{f} dz,$$

and the result follows from Theorem 19.93. ■

Definition 19.103. A mapping $F : \mathcal{D} \rightarrow \mathbb{C}$ is called **open** if for every open set $\mathcal{O} \subset \mathcal{D}$, the image $F(\mathcal{O})$ is open in $\mathbb{C}$.

Theorem 19.104 (Open Mapping Theorem). *Let f be analytic and non-constant in a domain $\mathcal{D}$. Then f is an open mapping.*

Proof. This is left as an exercise. (Hint: factor out the first nonzero term in the Taylor expansion of f at z_0 , and use the continuity of $\arg$ near 1.)

Namely, if $w_0 \in f(\mathcal{D})$, then there exists $z_0 \in \mathcal{D}$ such that $f(z_0) = w_0$. Consider the Taylor expansion

$$f(z) = w_0 + \sum_{k=1}^{\infty} f_k(z - z_0)^k.$$

Since f is non-constant, there is a smallest $k \geq 1$ with $f_k \neq 0$. Then we may factor

$$f(z) - w_0 = f_k(z - z_0)^k(1 + g(z)),$$

where g is analytic and $g(z) \rightarrow 0$ as $z \rightarrow z_0$. A key point is that $1 + g(z)$ remains close to 1 near z_0 , so $\log(1 + g(z))$ is analytic there and hence $\arg(1 + g(z)) = \text{Im}(\log(1 + g(z)))$ varies continuously (as does $\arg(z - z_0)^k$). From this, it follows that along every ray from w_0 there is a nearby point in $f(\mathcal{D})$ (check!). Finally, continuity of f implies that $f(\mathcal{D})$ contains a neighborhood of w_0 , proving that f is open. ■

20. THE INVERSE FUNCTION THEOREM

Theorem 20.105. *Assume f is analytic at z_0 and $f'(z_0) = a \neq 0$. Then there exists a disk $\mathbb{D}_\varepsilon(z_0)$ such that f is injective on $\mathbb{D}_\varepsilon(z_0)$ and maps it biholomorphically onto the open set $f(\mathbb{D}_\varepsilon(z_0))$; in particular, the inverse is analytic.*

In this situation we also say that f is a *conformal map* between $\mathbb{D}_\varepsilon(z_0)$ and $f(\mathbb{D}_\varepsilon(z_0))$. (We will return to this notion in more detail later.)

Proof. Without loss of generality, assume $z_0 = 0$ and $f(0) = 0$. Then

$$f(z) = az + z^2g(z),$$

where g is analytic near 0. For z, z_1 near 0,

$$\begin{aligned}
 f(z) - f(z_1) &= a(z - z_1) + (z^2g(z) - z_1^2g(z_1)) \\
 &= a(z - z_1) + (z - z_1)(z + z_1)g(z) + z_1^2(g(z) - g(z_1)) \\
 (20.2) \quad &= (z - z_1) \left[a + (z + z_1)g(z) + z_1^2 \frac{g(z) - g(z_1)}{z - z_1} \right].
 \end{aligned}$$

By analyticity of g , for z, z_1 in a small disk $\mathbb{D}_\rho(0)$,

$$\frac{g(z) - g(z_1)}{z - z_1} = \int_0^1 g'(z_1 + t(z - z_1)) dt,$$

so the bracket in (20.2) has a removable singularity at z_1 . In particular, it extends continuously and tends to a as $z, z_1 \rightarrow 0$. Hence we can choose $\rho > 0$ so small that

$$\left| (z + z_1)g(z) + z_1^2 \frac{g(z) - g(z_1)}{z - z_1} \right| < |a| \quad \text{for all } z, z_1 \in \mathbb{D}_\rho(0).$$

Then the bracket in (20.2) never vanishes on $\mathbb{D}_\rho(0) \times \mathbb{D}_\rho(0)$, so $f(z) = f(z_1)$ implies $z = z_1$. Therefore f is injective on $\mathbb{D}_\rho(0)$.

Now fix $w_0 = f(z_0)$ with $z_0 \in \mathbb{D}_\rho(0)$. From (20.2) we have, as $z \rightarrow z_0$,

$$\frac{f(z) - f(z_0)}{z - z_0} \longrightarrow a \neq 0.$$

Thus

$$\lim_{z \rightarrow z_0} \frac{z - z_0}{f(z) - f(z_0)} = \frac{1}{a},$$

which shows that f^{-1} is differentiable at w_0 with derivative $1/f'(z_0)$. Since z_0 was arbitrary, f^{-1} is analytic on $f(\mathbb{D}_\rho(0))$. ■

Exercise 20.106 (Important generalization). (a) Assume f is analytic in a neighborhood of z_0 (say $z_0 = 0$) and

$$f(0) = f'(0) = \cdots = f^{(k-1)}(0) = 0, \quad f^{(k)}(0) = a \neq 0.$$

Show that there exists $\varepsilon > 0$ such that, for every $t \in \mathbb{D}_\varepsilon(0)$, the equation $f(z) = t$ has *exactly* k roots near 0 (counted with multiplicity).

Hint: factor $f(z) = a(z^k + \text{higher order terms})$ and argue as in the proof of Theorem 20.105, or apply Rouché's theorem.

(b) Deduce from (a) an alternative proof of the open mapping theorem.

21. ANALYTIC CONTINUATION

Assume f is analytic in $\mathcal{D}$ and f_1 is analytic in $\mathcal{D}_1 \supsetneq \mathcal{D}$ with $f = f_1$ in $\mathcal{D}$. Then f_1 is called an *analytic extension* of f . We also say that f_1 is obtained from f by *analytic continuation*.

Weierstrass's point of view was to regard analytic functions as properly defined *chains of Taylor series*, up to a natural equivalence (see below), each Taylor series being the analytic continuation of the adjacent ones.

If f is analytic at z_0 , then there exists $\varepsilon > 0$ such that f is given by its Taylor series in $\mathbb{D}_\varepsilon(z_0)$. Let ε_0 be the maximal such radius. If $z_1 \in \mathbb{D}_{\varepsilon_0}(z_0)$, then f is also analytic at z_1 , hence represented by a Taylor series centered at z_1 . The radius of convergence of this series is at least $d(z_1, \partial\mathbb{D}_{\varepsilon_0}(z_0))$, but possibly larger. (Example: the function $1/(1+z)$ with disks centered at $z = 0$ and $z = 1/2$.)

In the latter case, we obtain a function f_1 , piecewise given by the two Taylor series, which is analytic in $\mathbb{D}_{\varepsilon_0}(z_0) \cup \mathbb{D}_{\varepsilon_1}(z_1)$. Continuing in this way, we construct chains $z_0, z_1, \dots$ with corresponding disks $\mathbb{D}_{\varepsilon_i}(z_i)$. There is then a natural equivalence relation: f_α and f_β are equivalent if they are analytic continuations of each other.

Definition 21.107 (Global analytic functions (Weierstrass)). *An analytic function in the sense of Weierstrass is an equivalence class under this relation.*

Uniqueness. If there is an analytic continuation of f to $\mathbb{D}_\varepsilon(z_0) \cup \mathbb{D}_{\varepsilon_1}(z_1)$, then it is unique (use Proposition 8.46).

Ahlfors makes this more precise as follows.

Definition 21.108 (Function elements and global analytic functions). *An analytic function defined on an open set $U \subset \mathbb{C}$ is called a function element and denoted (f, U) . Two function elements (f_1, U_1) and (f_2, U_2) are said to be analytic continuations of one another if $U_1 \cap U_2 \neq \emptyset$ and $f_1 = f_2$ on $U_1 \cap U_2$.*

A chain of analytic continuations is a finite sequence of function elements

$$(f_1, U_1), (f_2, U_2), \dots, (f_n, U_n)$$

such that each consecutive pair are analytic continuations of one another.

A global analytic function is a family $\mathcal{F}$ of function elements such that, for any $(f, U), (g, V) \in \mathcal{F}$, there exists a chain of analytic continuations in $\mathcal{F}$ beginning at (f, U) and ending at (g, V) .

A complete global analytic function is a global analytic function $\mathcal{F}$ that contains every analytic continuation of each of its elements.

But the *values of the continuations* may differ. For instance, starting from (f, U_0) with $f = \log$ defined near 1, we can build two chains of analytic continuations, one passing along the upper semicircle from 1 to -1 , the other along the lower semicircle. Both chains end in function elements defined on the same domain U near -1 , yet the resulting functions differ by $2\pi i$. Thus the same open set U may occur in two different function elements with distinct values, reflecting the multivalued nature of functions like $\log$.

This phenomenon leads naturally to the notion of *Riemann surfaces*, which provide the correct setting for treating multivalued analytic functions (such as $\log z$, $z^{1/n}$, or $\arcsin z$) as genuinely single-valued. We will return to this perspective later.

Cuts. One way to restore single-valuedness is to define $\log$ in $\mathbb{C} \setminus (-\infty, 0]$ where $(-\infty, 0]$ is called a **cut** and its function is to prevent curves to circle around 0.

We note that the cut $(-\infty, 0]$ is **highly arbitrary**. It can be replaced by any ray $\{te^{i\phi} : \phi \in [0, 2\pi)\}$ or any curve that has the same functionality, namely to ensure there are no paths of continuation that encircle 0. This is worth emphasizing this, because many books tend to give the wrong impression that the cut $(-\infty, 0]$, giving the "principal branch of the log", is somehow set in stone. Notably too, the **value of the log depends on the cut**.

Exercise 21.109. What is the value of $\log e^{3\pi i/4}$ if the cut is $(-\infty, 0]$ and what is it if the cut is $\{it : t \geq 0\}$?

Remark. The choice of cut is a way to make a multivalued function (like $\log z$) single-valued on a simply connected domain. Different choices of cuts give different branches of the function. From the perspective of Riemann surfaces, these branches are just different "sheets" of a single global object. We will return to this viewpoint later when discussing Riemann surfaces and branch points.

Natural boundaries. For general analytic functions, it might happen that one encounters lines, or more generally closed regions, through which no analytic continuation exists.

Such a boundary is called a *natural boundary*. It represents a set of "terminus" points for a global analytic function in the sense of Weierstrass. Yet, *some* natural boundaries can be traversed in a precise mathematical sense. This idea was pioneered by Borel (see Borel, *Leçons sur les fonctions monogènes*), who developed methods of analytic continuation through natural boundaries. We will return to this

question later. Separately, Borel also introduced methods of resumming divergent (zero radius of convergence) series, which later evolved into the modern theory of Borel summation and resurgent analysis.

The standard example of such a function is

$$f(z) = \sum_{k=0}^{\infty} z^{2^k}.$$

We have $f(z) \rightarrow \infty$ as $z \rightarrow 1$, and also as $z \rightarrow -1, i$, and $-i$, and more generally as $z \rightarrow e^{2\pi i N/2^M}$ with $(N, M) \in \mathbb{N}^2$. The points $N/2^M$ with $(N, M) \in \mathbb{N}^2$ are the *binary rationals*, i.e. the numbers whose base-2 expansion has only finitely many digits. They form a dense subset of $\mathbb{R}$, hence the points $e^{2\pi i N/2^M}$ are dense on the unit circle. This shows that **no point** of the unit circle is a point of analyticity (why?).

Exercise 21.110 (Another example of natural boundary). Consider the lacunary series

$$f(z) = \sum_{k=0}^{\infty} z^{n!}.$$

Show that $\partial\mathbb{D}$ is a natural boundary for f .

Hint: For each n , the $n!$ -th roots of unity are singular points of f . Observe that these roots are dense on $\partial\mathbb{D}$, and argue as in the case of $\sum z^{2^k}$ that no boundary point can be a point of analyticity.

This is a special case of a lacunary series: if the sequence of natural numbers p_k has the property that $\liminf_{n \rightarrow \infty} p_{n+1}/p_n = 1 + \delta$, then the series $\sum_{k=0}^{\infty} c_k z^{p_k}$ is called **lacunary**. There is a whole literature about them. The following theorem is due to Hadamard, with a proof in [8]:

Theorem 21.111 (Hadamard–Mandelbrojt). *If the exponents p_n satisfy $\liminf_{n \rightarrow \infty} p_{n+1}/p_n > 1$, and $\sum_{n=0}^{\infty} c_n z^{p_n}$ has radius of convergence 1, then $\partial\mathbb{D}$ is a natural boundary for f .*

Hadamard proved this under the stronger hypothesis $p_{n+1}/p_n > 1 + 1/n$, and Mandelbrojt later weakened it to the general form above.

Remark 21.112. Mandelbrojt’s condition is essentially sharp.

If $\liminf_{n \rightarrow \infty} \frac{p_{n+1}}{p_n} > 1$, then for *any* coefficients c_n , the unit circle is a natural boundary of f . On the other hand, if $\liminf_{n \rightarrow \infty} \frac{p_{n+1}}{p_n} = 1$, then there exist lacunary series of radius of convergence 1 which admit analytic continuation across parts of $\partial\mathbb{D}$. Thus Mandelbrojt’s criterion gives a necessary and sufficient condition for natural boundaries in this class. (see [16]).

Remark 21.113 (Genericity of natural boundaries). The examples above (such as $\sum z^{2^n}$ or $\sum z^{n!}$) show that highly sparse (“lacunary”) series typically have the unit circle as a natural boundary. Even more striking is the fact that natural boundaries are in fact *generic*: if one considers power series

$$f(z) = \sum_{k=0}^{\infty} c_k z^k, \quad c_k \in \{0, 1\},$$

and interprets the sequence $\{c_k\}_k$ as the binary expansion of a number in $[0, 1]$, then for a set of full measure (with respect to Lebesgue measure on $[0, 1]$), the function f has the unit circle as a natural boundary.

This phenomenon was first observed in work of Steinhaus (1929), and later developed in the probabilistic theory of random power series (see Kahane, *Some Random Series of Functions*, 2nd ed., 1985). In this sense, having the unit circle as a natural boundary is the “typical” behavior for analytic functions defined by such series.

As we shall see later, natural boundaries occur in the *uniformization of nontrivial Riemann surfaces*. For example, let $\mathcal{R}$ be the set of curves in $\mathbb{C} \setminus \{0, 1\}$ modulo homotopies. A uniformization ψ is a bi-analytic bijection between $\mathcal{R}$ and $\mathbb{D}$. Then $\partial\mathbb{D}$ is a natural boundary for ψ^{-1} .

Exercise 21.114 (Example of natural boundary due to Poincaré). Consider the rational numbers $r = p/q$ (with p, q relatively prime) and associate to r $N_{pq} = 7^{|p|}5^{|q|}$ (other primes would work too) (check that this is injective as a function from $\mathbb{Q}^+$ to $\mathbb{N}$). Similarly, find an isomorphism between $\mathbb{Q}^-$ and $\mathbb{Z}^-$. Define

$$(21.2) \quad f(z) = \sum_{N_{pq}}^{\infty} \frac{2^{-N_{pq}}}{z - p/q}.$$

Show that the series converges for $z \in \mathbb{C} \setminus \mathbb{R}$ and that $\mathbb{R}$ is a natural boundary for f (the poles accumulate densely on $\mathbb{R}$).

How can this example be modified to obtain an analytic function f in any domain bounded by a simple closed curve γ , such that γ is a natural boundary of f ?

22. THE SCHWARZ REFLECTION PRINCIPLE

Theorem 22.115. Assume f is analytic in two domains $\mathcal{D}_1$ and $\mathcal{D}_2$ which share a common boundary portion given by a piecewise differentiable curve γ . Suppose further that f is continuous on $\mathcal{D}_1 \cup \mathcal{D}_2 \cup \gamma$. Then f is analytic in

$$\mathcal{D} = \mathcal{D}_1 \cup \mathcal{D}_2.$$




FIGURE 12. Analytic continuation through a line along which a function is continuous.

Proof. By Morera's theorem, it suffices to show that the integral of f around any triangular contour in $\mathcal{D}$ vanishes. If the triangle lies entirely inside $\mathcal{D}_1$ or $\mathcal{D}_2$, this follows immediately from analyticity there. If the triangle intersects γ , split it along γ into two smaller triangles (see Fig. 14). Each piece lies in $\mathcal{D}_1$ or $\mathcal{D}_2$, so their integrals vanish. The contributions along γ cancel due to continuity of f , giving zero for the whole integral. ■

If all is known about a function f is analyticity in $\mathcal{D}_1$, and the fact that a piece of the boundary is an *analytic curve* (meaning, there is a parametrization by a function $\gamma : [0, 1] \rightarrow \mathbb{C}$ which extends analytically in a complex neighborhood of $[0, 1]$) up to which f is continuous, and a further condition on the values of f on γ holds, then it is possible to extend f analytically past γ . We start with the simplest such case.

Theorem 22.116 (The Schwarz reflection principle). Assume f is analytic in a domain $\mathcal{D}$ in the upper half plane (UHP, also denoted in these notes by $\mathbb{H}^u$) whose boundary contains an interval $I \subset \mathbb{R}$ and assume f is continuous on $\mathcal{D} \cup I$ and real valued on I . Then f has analytic continuation across I , in a domain $\mathcal{D} \cup \mathcal{D}^* \cup I$ where $\mathcal{D}^* = \{\bar{z} : z \in \mathcal{D}\}$.

Note 22.117. see §30.4 for a generalization of this result.

Proof. Define

$$F(z) = \begin{cases} f(z), & z \in \mathcal{D} \cup I, \\ \overline{f(\bar{z})}, & z \in \mathcal{D}^* \cup I. \end{cases}$$




FIGURE 13. Schwarz's reflection principle.

Then F is continuous on $\mathcal{D} \cup I \cup \mathcal{D}^*$ (check). It is analytic in $\mathcal{D}^*$ by a Taylor expansion argument. For Morera's theorem: integrals over closed curves lying entirely in $\mathcal{D}$ or in $\mathcal{D}^*$ vanish. If a curve crosses I , it can be split into two parts symmetric across I , which cancel because f is real on I (see again Fig. 14). Hence F is analytic in $\mathcal{D} \cup \mathcal{D}^*$, as desired. ■

Example 22.118. The square root $\sqrt{z} = \rho^{1/2}e^{i\phi/2}$, $z = \rho e^{i\phi}$ with $\phi \in [0, \pi)$, is analytic in the upper half-plane, continuous down to $[0, \infty)$, and real there. By Schwarz reflection, it continues analytically to the lower half-plane. For $z = \rho e^{-i\theta} \in \mathbb{H}_\ell$, $\theta \in (0, \pi)$, we have $\bar{z} = \rho e^{i\theta} \in \mathbb{H}^u$, so

$$\sqrt{z} = \overline{\sqrt{\bar{z}}} = \rho^{1/2}e^{-i\theta/2}.$$

23. MULTI-VALUED FUNCTIONS

As we discussed, as a result of analytic continuation in the complex plane we may get a *global analytic function* which is not necessarily a function **defined on** $\mathbb{C}$ since the definition is path-dependent; the function is thus defined on a space of paths or curves, modulo homotopies.

As long as the domain of continuation is simply connected, we still get a function in the usual sense:

Theorem 23.119 (The monodromy theorem). Assume $\mathcal{D}$ is a simply connected domain, $\mathbb{D}_\varepsilon(z_0) \subset \mathcal{D}$ that f is analytic in $\mathbb{D}_\varepsilon(z_0)$. Assume further that for any $z_1 \in \mathcal{D}$ there is a curve γ joining z_0 to z_1 such that f




FIGURE 14. The monodromy theorem.

has analytic continuation along γ ⁵. Then, f has analytic continuation in $\mathcal{D}$.

Proof. Take $z_1 \in \mathcal{D}$ and let $\mathcal{D}' \subset \mathcal{D}$ be a simply connected subdomain containing $\{z_0, z_1\}$ whose boundary decomposes as $\gamma_1 \cup \gamma_2$, where γ_1 and γ_2 are smooth curves with common endpoints.

Choose a curve γ_3 sufficiently close to γ_1 so that: (a) γ_3 can be covered by disks $U'_1 = U_1 = \mathbb{D}_\varepsilon(z_0), \dots, U'_k = U_n = \mathbb{D}_\delta(z_1)$; (b) the domain $\mathcal{D}_3 = U_1 \cup U_2 \cup \dots \cup U_n$ is connected and simply connected; and (c) $(f, U_1), (g_2, U_2), \dots, (g_k, U_k)$ forms a chain of analytic continuations.

Thus f is analytic in each U_j , and by construction the restrictions agree on overlaps. By uniqueness of analytic continuation (Proposition 8.46), the local definitions glue together to a single analytic function on $\mathcal{D}_3$.

Repeating this deformation step, one can deform γ_1 continuously into γ_2 without losing analyticity. Hence f admits analytic continuation in $\mathcal{D}'$. Since $\mathcal{D}'$ was arbitrary, f extends analytically to the whole of $\mathcal{D}$.

■

In a nutshell: if f admits analytic continuation along every path in a simply connected domain $\mathcal{D}$, then f extends to an analytic function on $\mathcal{D}$. Equivalently, for analyticity to fail in $\mathcal{D}$, there must exist singular points of f within $\mathcal{D}$ that obstruct continuation.

Exercise 23.120 (Alternative proof of monodromy). Let $\overline{\mathcal{D}_1} \subsetneq \mathcal{D}$ be a connected, simply connected compact domain with smooth boundary

⁵More precisely, since γ is compact, γ is contained in a finite union of open disks in $\mathcal{D}$ such that $U_1 = \mathbb{D}_\varepsilon(z_0), \dots, U_n = \mathbb{D}_\delta(z_1)$... such that $(f, U_1), (f_2, U_2), \dots, (f_n, U_n)$ form a chain of analytic continuations.

that contains z_0 . Cover $\overline{\mathcal{D}_1}$ by finitely many disks $\{U_j\}_{j=1}^N$, with $U_1 = \mathbb{D}_\rho(z_0)$, arranged so that consecutive disks intersect.

Since analytic continuation exists along any polygonal arc contained in $\overline{\mathcal{D}_1}$ that joins adjacent disks, define F on $\overline{\mathcal{D}_1}$ by continuing f from U_1 to any $p \in \overline{\mathcal{D}_1}$ along such a polygonal chain of overlapping disks.

Show that F is *well-defined* (path-independent): if two polygonal chains join z_0 to p , together they form a closed loop in $\overline{\mathcal{D}_1}$; since $\overline{\mathcal{D}_1}$ is simply connected, the loop is null-homotopic. By uniqueness of analytic continuation on overlaps, the two continuations agree. Conclude that F is single-valued and analytic on $\mathcal{D}_1$.

Because $\overline{\mathcal{D}_1} \subsetneq \mathcal{D}$ was arbitrary, exhaust $\mathcal{D}$ by an increasing sequence of such compact, simply connected domains $\overline{\mathcal{D}_k}$ with $\bigcup_{k \in \mathbb{N}} \mathcal{D}_k = \mathcal{D}$. Construct a sequence $(\mathcal{D}_k, f_k)_{k \in \mathbb{N}}$ where f_k is the analytic continuation on $\mathcal{D}_k$ and $f_k|_{\mathcal{D}_{k-1}} = f_{k-1}$. Then $f := \bigcup_{k \in \mathbb{N}} f_k$ defines the desired analytic continuation on all of $\mathcal{D}$.

23.1. Generalization: log of a function. If g is a function defined in a region in $\mathbb{C}$, we can define $\ln g$ by

$$(23.2) \quad \ln g(z) = \int_a^z \frac{g'(s)}{g(s)} ds.$$

Now, depending on the properties of g , the homotopy classes will in general be more complicated than those of $\log z$.

If, for instance, g is a meromorphic function in $\mathbb{C}$, then the set S consisting of all the zeros and poles of g are points where the integral is undefined. The logarithm of g is defined on homotopy classes of curves over $\mathbb{C} \setminus S$.

It is sometimes convenient to define a branch of $\ln g$ by cutting the plane along rays originating at the points in S . (Check that such cuts exist for any meromorphic g —hint: join each singularity to infinity with non-intersecting rays.) See also Fig. 15.

23.2. General powers of z . Once we have defined the log, it is natural to define

$$(23.3) \quad z^\alpha = e^{\alpha \ln z}$$

Since $\ln z$ is defined on the homotopy classes of curves over $\mathbb{C} \setminus \{0\}$, so is z^α . For a general $\alpha \in \mathbb{C}$, the multivaluedness of z^α is inherited from the multivaluedness of the log: $e^{\alpha(\ln z + 2N\pi i)} = e^{\alpha \ln z} e^{2N\pi i \alpha}$. Note however that if $\alpha \in \mathbb{Z}$ then the value does not depend on the homotopy class and the definition (23.3) defines a function in $\mathbb{C} \setminus \{0\}$, with a pole




FIGURE 15. Cuts defining the log of a rational function.

at zero if $\alpha < 0$ and a removable singularity if $\alpha \geq 0$. Check that this definition coincides with the usual power on $\mathbb{R}^+$.

If $\alpha = p/q$ with p, q relatively prime, then $z^{p/q}$ admits exactly q distinct values corresponding to the q -th roots of unity.

Note. Beware of possible pitfalls.

$$(23.4) \quad e^{\ln z_1 + \ln z_2} = e^{\ln z_1} e^{\ln z_2} = z_1 z_2$$

However, this does not mean $\ln z_1 + \ln z_2 = \ln z_1 z_2$, but just that

$$(23.5) \quad \ln z_1 + \ln z_2 = \ln z_1 z_2 + 2N\pi i, N \in \mathbb{Z}.$$

For the same reason, $z^{\alpha_1} z^{\alpha_2}$ is not necessarily $z^{\alpha_1 + \alpha_2}$. Note the fallacious calculation⁶

$$(23.6) \quad 1 = \sqrt{1} = \sqrt{(-1)(-1)} = \sqrt{-1}\sqrt{-1} = i \cdot i = -1 \quad (!)$$

24. CONFORMAL MAPPING

Laplace's equation in two dimensions,

$$(24.2) \quad \Delta f = f_{xx} + f_{yy} = 0,$$

arises in many physical problems.

In electrostatics, for instance, the equation $\Delta V = 0$ describes the potential V in a charge-free region, following from two of Maxwell's

⁶This calculation was done as presented by an early version of a computer algebra program.

equations:

$$\nabla \cdot \mathbf{E} = \varepsilon_0^{-1} \rho = 0, \quad \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} = 0,$$

which imply $\mathbf{E} = -\nabla V$ and $\Delta V = 0$. Boundary conditions are imposed by the distribution of charges, and the physical problem becomes: solve (24.2) with V prescribed on $\partial\mathcal{D}$ (the *Dirichlet problem*).

In fluid dynamics, consider a two-dimensional incompressible velocity field $\mathbf{V} = (v, u)$. Incompressibility gives

$$(24.3) \quad \operatorname{div}(v, u) = v_x + u_y = 0,$$

while irrotationality implies

$$(24.4) \quad \nabla \times \mathbf{V} = 0 \Rightarrow u_x - v_y = 0.$$

Equations (24.3)–(24.4) show that u and v are harmonic conjugates. Thus, in any simply connected domain, there exists a *velocity potential* φ with $\mathbf{V} = \nabla\varphi$ and $\Delta\varphi = 0$. Its harmonic conjugate ψ is called the *stream function*, and the curves $\psi = \text{const.}$ coincide with the streamlines of the flow. Physically, the natural boundary condition is impermeability of the boundary $\partial\mathcal{D}$:

$$(v, u) \cdot (n_1, n_2) = 0,$$

the so-called *no-penetration condition*. Laplace's equation with prescribed normal derivative on $\partial\mathcal{D}$ is known as the *Neumann problem*.

We already know the explicit solution of the Dirichlet problem for the special case $\mathcal{D}$ a disk: Poisson's formula (11.5). More generally, existence and uniqueness of solutions hold for any bounded domain $\mathcal{D}$ with sufficiently smooth boundary and continuous boundary data.

24.1. Uniqueness. Uniqueness follows from the maximum principle: if u_1, u_2 are two solutions of (24.2) with the same boundary conditions, then $u = u_1 - u_2$ satisfies (24.2) with $u = 0$ on $\partial\mathcal{D}$. But a harmonic function attains both its maximum and its minimum on the boundary; hence $u \equiv 0$. Similarly, in the Neumann problem, u is determined uniquely up to an arbitrary additive constant.

Often, a PDE possesses a symmetry group, which can be exploited both for solving the equation and for understanding its qualitative behavior. For Laplace's equation in $\mathbb{C}$, the symmetry group is especially large: it is invariant under conformal maps (analytic homeomorphisms). More precisely, if $f : \mathcal{D}_1 \rightarrow \mathcal{D}_2$ is a biholomorphism between simply connected domains, then solutions in $\mathcal{D}_2$ pull back to solutions in $\mathcal{D}_1$. In this sense, Laplace's equation is said to be *conformally invariant*.

Proposition 24.121. *If u solves (24.2) ($\Delta u = 0$) in a simply connected domain $\mathcal{D}_2$, and $f = f_1 + if_2 : \mathcal{D}_1 \rightarrow \mathcal{D}_2$ is analytic, then*

$$(x, y) \mapsto u(f_1(x, y), f_2(x, y))$$

is a solution of (24.2) in $\mathcal{D}_1$.

Proof. Since u is harmonic in $\mathcal{D}_2$, it has a harmonic conjugate v (unique up to an additive constant). Thus $g = u + iv$ is analytic in $\mathcal{D}_2$. Then $g \circ f$ is analytic in $\mathcal{D}_1$, and hence its real and imaginary parts, $u \circ f$ and $v \circ f$, satisfy the Cauchy–Riemann equations in $\mathcal{D}_1$. It follows that $u \circ f$ is harmonic in $\mathcal{D}_1$. ■

We will be especially interested in analytic *homeomorphisms*, which preserve the topological structure of regions. Two regions related by such a map are said to be *conformally equivalent*.

The celebrated **Riemann mapping theorem** (to be proved later) states that any simply connected domain other than $\mathbb{C}$ itself is conformally equivalent to the unit disk. In this correspondence, the boundary of the region is mapped onto the unit circle. Thus

Remark 24.122. *The orbit of the unit disk under all conformal transformations contains every simply connected region (except $\mathbb{C}$ itself).*

Because of this invariance, the conformal group is powerful enough that, in principle, Laplace’s equation in any simply connected domain (with sufficiently smooth boundary, say $C^{1,\alpha}$) can be reduced to the disk case. This is one of the key motivations for studying conformal mappings carefully.

24.2. Preservation of angles and small shapes: heuristics. Let f be analytic at z_0 with $f'(z_0) = a \neq 0$. (Without loss of generality we may assume $z_0 = 0$ and $f(0) = 0$.) Consider a tiny neighborhood $\mathcal{N}$ of 0. If $z_\beta \in \mathcal{N}$ then

$$(24.5) \quad f(z_\beta) \approx az_\beta.$$

Thus all points near 0 are multiplied by the *same* complex number a . Multiplication by a rescales vectors by $|a|$ and rotates them by $\arg(a)$. If we think of the collection $\{z_\beta\}$ as forming a small figure, then $\{f(z_\beta)\}$ is essentially the same figure, uniformly rotated and rescaled. In particular, *angles are preserved*, and the local shape is unchanged up to similarity. This is precisely what it means for f to be *conformal*.

Since a tiny square of side ε is mapped to a square of side $|a|\varepsilon$, areas are multiplied by the factor $|a|^2$.

We make this rigorous in what follows.




FIGURE 16. Conformal kitty. Angles are preserved, shapes are locally similar, but scales and orientations are changed.

24.3. Preservation of angles. Recall that additivity of arguments holds up to an integer multiple of 2π . In the calculations below, this ambiguity will be immaterial.

Assume f is analytic in a disk D , that $z_0 \in D$, and that $f'(z_0) \neq 0$. Let $\gamma(t)$ and $\Gamma(t)$ be two smooth curves crossing at z_0 , and without loss of generality assume $\gamma(0) = \Gamma(0) = z_0$. The angle between the curves is defined as the difference of the arguments of their tangent vectors:

$$\theta = \arg \gamma'(0) - \arg \Gamma'(0),$$

provided these derivatives are nonzero.

Now consider the images of these curves under f . The corresponding tangent vectors at $f(z_0)$ are

$$(f \circ \gamma)'(0) = f'(z_0)\gamma'(0), \quad (f \circ \Gamma)'(0) = f'(z_0)\Gamma'(0).$$

Hence the angle between the images is

$$\begin{aligned} (24.6) \quad & \arg[(f \circ \gamma)'(0)] - \arg[(f \circ \Gamma)'(0)] = \arg(f'(z_0)\gamma'(0)) - \arg(f'(z_0)\Gamma'(0)) \\ & = \arg f'(z_0) + \arg \gamma'(0) - (\arg f'(z_0) + \arg \Gamma'(0)) \\ & = \arg \gamma'(0) - \arg \Gamma'(0) = \theta. \end{aligned}$$

Thus the image of two curves intersecting at an angle θ is a pair of curves intersecting at the same angle θ . In other words, analytic maps with $f'(z_0) \neq 0$ preserve angles (and orientation) at z_0 . Equivalently, a sufficiently small domain is transformed into a similar one — rescaled and rotated, but not distorted.

Exercise 24.123. More precisely, consider a sequence of similar triangles whose sides go to zero and with a common point z_0 , with $f'(z_0) \neq 0$. Consider their images through f . For each triangle, rescale both the triangle by the size L of one of its sides, and rescale their images through

f by $L|f'(z_0)|$. Show that in the limit of vanishing sides, the limit of the rescaled triangles and the limit of the rescaled images of these triangles are congruent. Hence, applying a similar procedure to polygons, we will obtain congruent polygons. The same extends to shapes that are limits of polygons.

24.4. Rescaling of arc length. The arc length along a curve $\gamma(t)$ is given by

$$(24.7) \quad L(\gamma) = \int_{\gamma} d|z| := \int_a^b |\gamma'(t)| dt.$$

If f is analytic, then

$$(24.8) \quad L(f(\gamma)) = \int_a^b |(f \circ \gamma)'(t)| dt = \int_a^b |f'(\gamma(t))| |\gamma'(t)| dt = \int_{\gamma} |f'(z)| d|z|.$$

Thus the arc length is locally stretched by the factor $|f'(z)|$, in agreement with Exercise 24.123.

24.5. Transformation of areas. The area of a domain A is its Lebesgue measure,

$$(24.9) \quad \iint_A dx dy.$$

Let $f = u + iv : B \rightarrow A$ be biholomorphic. Under the change of variables $(x, y) \mapsto (u(x, y), v(x, y))$, the area of A can be computed as

$$(24.10) \quad \iint_B |J(x, y)| dx dy,$$

where J is the Jacobian determinant

$$J(x, y) = \det \begin{pmatrix} u_x & u_y \\ v_x & v_y \end{pmatrix}.$$

Using the Cauchy–Riemann equations (see (9.3)) one finds that $J(x, y) = |f'(z)|^2$.

Geometrically, this means that areas are locally stretched by the factor $|f'(z)|^2$, just as lengths are stretched by $|f'(z)|$ (cf. Exercise 24.123).

Note 24.124. It is interesting to observe that if a smooth transformation (u, v) of the plane preserves angles, then $u + iv$ must be analytic (i.e., conformal). Likewise, if it rescales every infinitesimal figure by the same factor, then the map is either analytic, or it coincides with

the complex conjugate of an analytic function.⁷ This is not difficult to prove; see [3, §II.3], p. 74.

Thus analytic functions can be characterized geometrically as those maps which are *locally Euclidean*: they transform small shapes into similar shapes, preserving angles (orientation-preserving in the analytic case, orientation-reversing in the conjugate case).

Note 24.125. Observe that we did not require f to be globally one-to-one. The simple fact that f is analytic with nonzero derivative makes it conformal.

Remark 24.126 (On bijectivity). In $\mathbb{R}$, it suffices that $f' \neq 0$ everywhere for f to be conformal. This is **not** the case in $\mathbb{C}$. For instance, $\exp'(z) = \exp(z) \neq 0$ for all $z \in \mathbb{C}$, yet the exponential function is not injective: $\exp$ wraps the plane around the punctured plane infinitely many times.

24.6. Puiseux series.

Proposition 24.127 (Ramified expansions: Puiseux series). Assume that $f : \mathcal{D}_1 \rightarrow \mathcal{D}_2$ is analytic, and that for some $z_0 \in \mathcal{D}_1$ we have

$$f^{(j)}(z_0) = 0 \quad \text{for } j = 1, \dots, m-1, \quad f^{(m)}(z_0) = a \neq 0.$$

(As usual, we arrange $z_0 = 0$ and $f(z_0) = 0$.) Let $\{\omega_k\}_{1 \leq k \leq m}$ be the m th roots of unity. Then there exists a biholomorphic function φ in a neighborhood of 0 with $\varphi(0) = 0$ such that, for a chosen branch of $t^{1/m}$, all the solutions of the equation $f(z) = t$ near 0 are given by

$$z_k = \varphi(\omega_k t^{1/m}), \quad 1 \leq k \leq m.$$

Proof. In a neighborhood of 0 we may write

$$f(z) = z^m(a + zE(z)) = z^m g(z),$$

where g is analytic and $g(0) = a \neq 0$. Thus $g(z) \neq 0$ in some disk $\mathbb{D}_{\varepsilon_1}(0)$. There exists an analytic branch $h = g^{1/m}$ in $\mathbb{D}_{\varepsilon_1}(0)$. Choosing also a branch of $t^{1/m}$, our equation becomes

$$(zh(z))^m = t,$$

equivalent to m equations

$$zh(z) = \omega_k t^{1/m}, \quad k = 1, \dots, m.$$

The function $zh(z)$ is analytic at 0, and

$$(zh)'(0) = h(0) \neq 0.$$

⁷In this case the transformation is *orientation-reversing*. Geometrically, this means that locally the map looks like $z \mapsto \overline{f(z)}$ for some analytic f .




FIGURE 17. Ramified expansions (to be made rigorous in the sequel).

By the inverse function theorem, each equation $zh = \omega_k t^{1/m}$ has exactly one solution near 0, namely

$$z_k(t) = \varphi(\omega_k t^{1/m}),$$

where φ is a biholomorphism in a neighborhood of 0. ■

Note 24.128. Expanding φ as a Taylor series,

$$\varphi(s) = \sum_{j \geq 0} a_j s^j,$$

we obtain

$$z_k(t) = \varphi(\omega_k t^{1/m}) = \sum_{j \geq 0} a_j (\omega_k t^{1/m})^j = \sum_{j \geq 0} a_j \omega_k^j t^{j/m}.$$

This is a *ramified expansion*, i.e. a convergent series in fractional powers of t . Such expansions are called *Puiseux series*.

This gives another proof of the open mapping theorem:

Theorem 24.129. *Let $\mathcal{D}$ be a domain and $f : \mathcal{D} \rightarrow \mathbb{C}$ a nonconstant analytic function. Then f is open; that is, the image of every open set in $\mathcal{D}$ is open.*

Proof. This follows immediately from the fact that φ is a biholomorphism. ■

25. EVALUATION OF DEFINITE INTEGRALS

Contour integrals can be evaluated using the residue theorem. Many improper integrals with endpoints at infinity can also be computed this way: they can be regarded as contour integrals on the Riemann sphere, where ∞ is often a singular point of the integrand. Similarly, many integrals with singular endpoints can be evaluated in closed form. We begin with a simple but important case.

Proposition 25.130. *Let R be a rational function, continuous on $\mathbb{R}$, such that there exists $C > 0$ with*

$$|R(z)| \leq C|z|^{-2} \quad \text{for all } z \in \mathbb{C}.$$

(In particular, this holds if the degree of the numerator is at least two less than that of the denominator.) Then

$$(25.2) \quad \int_{-\infty}^{\infty} R(x) dx = 2\pi i \sum_{z_i \in \text{UHP}} \text{Res}(R; z = z_i) = 2\pi i \sum_{z_i \in \text{LHP}} \text{Res}(R; z = z_i),$$

where the sums run over the poles of R in the upper (resp. lower) half-plane.

In particular, if all poles of R lie strictly in the upper (or strictly in the lower) half-plane, then the integral vanishes.

Proof. Consider the rectangular contour shown in Fig. 42, consisting of the interval $[-A, A]$ on the real line together with the top horizontal segment C_1 and two vertical segments. By the residue theorem,

$$\oint_{[-A, A] \cup C_1} R(z) dz = 2\pi i \sum_{z_i \in \text{UHP}} \text{Res}(R; z = z_i).$$

Hence

$$(25.3) \quad \begin{aligned} \int_{-\infty}^{\infty} R(x) dx &= \lim_{A \rightarrow \infty} \int_{-A}^A R(x) dx = \lim_{A \rightarrow \infty} \left[\oint_{[-A, A] \cup C_1} R(z) dz - \int_{C_1} R(z) dz \right] \\ &= 2\pi i \sum_{z_i \in \text{UHP}} \text{Res}(R; z = z_i) - \lim_{A \rightarrow \infty} \int_{C_1} R(z) dz. \end{aligned}$$

But by the assumption $|R(z)| = O(|z|^{-2})$ as $|z| \rightarrow \infty$,

$$\left| \int_{C_1} R(z) dz \right| \leq \left(\sup_{z \in C_1} |R(z)| \right) \cdot \text{length}(C_1) = O(A^{-2}) \cdot O(A) = O(A^{-1}) \rightarrow 0,$$

so the integral over C_1 vanishes in the limit and we obtain the claim.

■




FIGURE 18. Rectangular contour used to evaluate $\int_{-\infty}^{\infty} R(x) dx$.

Note. It is useful to interpret the procedure above as “pushing” (or deforming) the original contour of integration $\mathbb{R}$ upwards toward $+i\infty$, collecting residues along the way. Since a rational function has only finitely many poles, once the contour has been pushed above the highest pole (say to $\text{Im } z = h$), it can be moved further to $+i\infty$ without encountering new singularities. On this final horizontal segment, the integral tends to zero as $h \rightarrow +\infty$ because of the decay assumption on R . By contour independence in singularity-free regions, the contribution of this part is therefore exactly zero.

Example 25.131. Evaluate

$$I = \int_{-\infty}^{\infty} \frac{dx}{1+x^4}.$$

Solution. The poles of $R(x) = \frac{1}{1+x^4}$ are the fourth roots of -1 , namely

$$e^{i\pi/4}, e^{3i\pi/4}, e^{5i\pi/4}, e^{7i\pi/4}.$$

In the upper half-plane we have $z_1 = e^{i\pi/4}$ and $z_2 = e^{3i\pi/4}$. The residue at each is

$$\text{Res}\left(\frac{1}{1+z^4}; z = z_i\right) = \frac{1}{(1+z^4)'|_{z=z_i}} = \frac{1}{4z_i^3}.$$

Therefore,

$$I = 2\pi i \left(\frac{1}{4z_1^3} + \frac{1}{4z_2^3} \right).$$

A short computation gives

$$\frac{1}{z_1^3} + \frac{1}{z_2^3} = \frac{1}{e^{3i\pi/4}} + \frac{1}{e^{9i\pi/4}} = e^{-3i\pi/4} + e^{-i\pi/4} = \sqrt{2}i.$$

Hence

$$I = 2\pi i \cdot \frac{1}{4} \cdot (\sqrt{2}i) = \frac{\pi}{\sqrt{2}}.$$

25.1. Fourier-type integrals. In calculating Fourier transforms, we encounter integrals of the form

$$(25.4) \quad \int_{-\infty}^{\infty} e^{iax} Q(x) dx,$$

where Q is a rational function, analytic on $\mathbb{R}$, and decaying at least like $1/x$ at infinity so that the (possibly improper) integral is well defined. We take $a > 0$ for definiteness, as in the inverse Fourier transform. As in the previous section, one way to evaluate the integral is to close the contour in the upper half-plane and push it towards $+i\infty$, where the exponential decays. To justify that this additional arc contributes nothing in the limit, Jordan's lemma is the standard tool.

Lemma 25.132 (Jordan). Assume $a > 0$, that Q is analytic in

$$\mathcal{D} = \{z : \operatorname{Im} z \geq 0, |z| > c\},$$

and let γ be the semicircle of radius $\rho > c$ in the UHP centered at the origin. Assume further that for large $|z|$, we have $|Q(z)| \leq M$. Then

$$(25.5) \quad \left| \int_{\gamma} e^{iaz} Q(z) dz \right| \leq \frac{M\pi}{a}.$$

In particular, if $Q(z) \rightarrow 0$ as $z \rightarrow \infty$, then

$$(25.6) \quad \int_{\gamma} e^{iaz} Q(z) dz \longrightarrow 0 \quad \text{as } \rho \rightarrow \infty.$$

Proof. Let ρ_0 be such that $|Q(z)| \leq M$ for $|z| > \rho_0$. For $\rho > \rho_0$ and γ as above,

$$(25.7) \quad \begin{aligned} \left| \int_{\gamma} e^{iaz} Q(z) dz \right| &= \left| \int_0^{\pi} e^{ia\rho e^{i\phi}} Q(\rho e^{i\phi}) \rho i e^{i\phi} d\phi \right| \\ &\leq M \int_0^{\pi} \rho e^{-\rho a \sin \phi} d\phi = 2M \int_0^{\pi/2} \rho e^{-\rho a \sin \phi} d\phi, \end{aligned}$$

where the last equality uses the symmetry $\sin \phi = \sin(\pi - \phi)$.

Next, note that $\theta^{-1} \sin \theta$ is decreasing on $[0, \pi/2]$, so

$$\sin \theta \geq \frac{2}{\pi} \theta, \quad 0 \leq \theta \leq \frac{\pi}{2}.$$

Hence

$$(25.8) \quad \left| \int_{\gamma} e^{iaz} Q(z) dz \right| \leq 2M \int_0^{\pi/2} \rho e^{-\frac{2\rho a}{\pi} \phi} d\phi \leq \frac{M\pi}{a},$$

which proves (25.5). The convergence to zero in the case $Q(z) \rightarrow 0$ as $|z| \rightarrow \infty$ is then immediate. ■

Proposition 25.133. Assume $a > 0$ and Q is a rational function, continuous on $\mathbb{R}$, and vanishing as $|z| \rightarrow \infty$. Then

$$(25.9) \quad \int_{-\infty}^{\infty} Q(x)e^{iax} dx = 2\pi i \sum_{z_i \in \mathbb{H}^u} \text{Res}(Q(z)e^{iaz}; z = z_i),$$

where the sum runs over the poles of Q in the open upper half-plane $\mathbb{H}^u$.

Proof. This follows immediately by combining Jordan's lemma (Lemma 25.132) with the residue computation in Proposition 25.130. Details are left as an exercise. ■

25.2. Further examples.

Example 25.134. Let $\tau > 0$ and evaluate

$$(25.10) \quad I(\tau) = \int_0^{\infty} \frac{\cos(\tau x)}{x^2 + 1} dx.$$

Solution. The integrand is even, so

$$(25.11) \quad 2I(\tau) = \int_{-\infty}^{\infty} \frac{\cos(\tau x)}{x^2 + 1} dx = \text{Re} \int_{-\infty}^{\infty} \frac{e^{i\tau x}}{x^2 + 1} dx.$$

This is of the form in Proposition 25.133, with poles in $\mathbb{H}^u$ at $z_1 = e^{i\pi/4}$ and $z_2 = e^{3i\pi/4}$. A short residue calculation gives

$$I(\tau) = \frac{\pi}{2} e^{-\tau}.$$

Note that $I(\tau)$ is the cosine Fourier transform of a function analytic in a neighborhood of $\mathbb{R}$, and it decays exponentially as $\tau \rightarrow \infty$.

Exercise 25.135 ([14], p. 116). Assume $\text{Re } z > 0$. Show that

$$(25.12) \quad I(z) = \int_0^{\infty} \frac{e^{-t} - e^{-tz}}{t} dt = \log z.$$

Hint. The integrand is continuous at $t = 0$, so the integral is well defined. Moreover $I(z)$ depends analytically on z , and by dominated convergence,

$$(25.13) \quad I'(z) = \int_0^{\infty} e^{-tz} dt = \frac{1}{z}.$$

Thus $I(z) = \log z + C$ for some constant C . Show that $C = 0$.

Example 25.136 (An important definite integral). Show that

$$(25.14) \quad \int_0^\infty \frac{\sin t}{t} dt = \frac{\pi}{2}.$$

Solution. A naive attempt to write

$$(25.15) \quad \int_{-\infty}^\infty \frac{\sin t}{t} dt \stackrel{?}{=} \operatorname{Im} \int_{-\infty}^\infty \frac{e^{it}}{t} dt$$

fails, since the right-hand side is ill defined. Instead, proceed as follows:

- (1) Using the “box” contour argument (see figure below), with $A = (2N + 1)\pi/2$, $N \in \mathbb{Z}$, show that

$$\int_{-\infty}^\infty \frac{\sin t}{t} dt = \int_{-\infty+i}^{\infty+i} \frac{\sin t}{t} dt = \int_{-\infty}^\infty \frac{\sin(t+i)}{t+i} dt.$$


- (2) Now compute

$$\int_{-\infty}^\infty \frac{\sin(t+i)}{t+i} dt = \int_{-\infty}^\infty \frac{e^{i(t+i)} - e^{-i(t+i)}}{2i(t+i)} dt.$$

The first term vanishes by Proposition 25.133. The second term equals

$$\overline{\int_{-\infty}^\infty \frac{-e^{i(t-i)}}{-2i(t-i)} dt},$$

to which Proposition 25.133 applies again. Carrying out the residue computation yields the desired result.



Remark 25.137. The same method shows more generally that

$$\int_0^\infty \frac{\sin(ax)}{x} dx = \begin{cases} \frac{\pi}{2}, & a > 0, \\ -\frac{\pi}{2}, & a < 0. \end{cases}$$

This fact will reappear naturally in Fourier analysis, where it corresponds to the Fourier transform of the sign function.

26. INTEGRALS OF BRANCHED FUNCTIONS

We now show that, for $\alpha \in (0, 1)$,

$$(26.2) \quad \int_0^\infty \frac{t^{-\alpha}}{t+1} dt = \frac{\pi}{\sin(\pi\alpha)}.$$

Note first that the improper integral is well defined.

We place a cut along $\mathbb{R}^+$, and adopt the convention that $t^{-\alpha} = |t|^{-\alpha}$ on the upper side of the cut. If we turn clockwise around the origin, then on the lower side of the cut

$$t^{-\alpha} = |t|^{-\alpha} e^{-2\pi i \alpha}.$$

Hence the integral in (26.2) equals

$$(26.3) \quad \frac{1}{1 - e^{-2\pi i \alpha}} \int_\gamma \frac{t^{-\alpha}}{t+1} dt = \frac{1}{1 - e^{-2\pi i \alpha}} \lim_{R \rightarrow \infty} \int_{\partial D_R} \frac{t^{-\alpha}}{t+1} dt,$$

where γ is the boundary of $\mathbb{C} \setminus [0, \infty)$ traversed from 0 to ∞ on the upper side of the cut and from ∞ back to 0 on the lower side, and ∂D_R is the boundary of the disk D_R of radius R centered at 0 with a cut along $[0, R]$. This is a simply connected domain containing the pole of the integrand at -1 , traversed counterclockwise. See Fig. 19.

The latter integral can be calculated by residues: the residue of the integrand at $t = -1$ is

$$\text{Res}_{t=-1} \frac{t^{-\alpha}}{t+1} = e^{-\pi i \alpha},$$

which yields the result.

More generally, we have the following result.

Proposition 26.138. Assume $\text{Re } a \in (0, 1)$ and let Q be a rational function, continuous on $\mathbb{R}^+$, such that $x^a Q(x) \rightarrow 0$ both as $x \rightarrow 0$ and as $x \rightarrow \infty$. Then

$$(26.4) \quad \int_0^\infty x^{a-1} Q(x) dx = -\frac{\pi e^{-\pi i a}}{\sin(\pi a)} \sum_{z_i} \text{Res}(z^{a-1} Q(z); z_i),$$

where the sum runs over all poles z_i of Q .

Exercise 26.139. Prove Proposition 26.138.

26.0.1. Conformally Equivalent Domains.

Definition 26.140. Two domains $\mathcal{D}_1$ and $\mathcal{D}_2$ are **conformally equivalent** if there exists a biholomorphic bijection $f: \mathcal{D}_1 \rightarrow \mathcal{D}_2$. In this case, we say that f **maps conformally** $\mathcal{D}_1$ onto $\mathcal{D}_2$.





FIGURE 19. Contour for the evaluation of (26.2).

The boundary behavior of analytic maps is a crucial tool in determining their conformal mapping properties.

Recall that a curve is traversed in a positive (counterclockwise) direction if its parameterization is such that the interior lies to the left of the curve as the parameter increases.



Proposition 26.141. Assume that $f: \mathcal{D} \rightarrow \mathbb{C}$ is analytic and that γ is a simple, piecewise differentiable closed curve contained in $\mathcal{D}$ along with its interior.

If f is one-to-one from γ onto $f(\gamma)$, then f maps $\text{Int}(\gamma)$ one-to-one and conformally onto $f(\text{Int}(\gamma))$, and it preserves the orientation of the curve.

Proof. Since f is continuous and one-to-one on the simple closed curve γ , its image $f(\gamma)$ is also a simple closed curve. Let $w_0 \in \text{Int}(f(\gamma))$, and set $z_0 \in \text{Int}(\gamma)$ such that $f(z_0) = w_0$. By Theorem 19.93 (the argument principle), the number of roots of $f(z) - w_0$ in $\text{Int}(\gamma)$ equals

$$(26.5) \quad I := \frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z) - w_0} dz,$$

which is some positive integer n .

By the change of variables $w = f(z)$, and since f is injective on γ , this becomes

$$I = \frac{1}{2\pi i} \int_{f(\gamma)} \frac{dw}{w - w_0}.$$

This integral equals ± 1 , depending on the orientation of $f(\gamma)$. The minus sign is impossible since $I = n > 0$. Hence $n = 1$, and the orientation of $f(\gamma)$ is positive.

Therefore, f maps $\text{Int}(\gamma)$ injectively and conformally onto $\text{Int}(f(\gamma))$.

■

26.1. Automorphisms. Automorphisms are biholomorphic maps from a domain $\mathcal{D}$ onto itself. They play an important role in complex analysis, since they describe the symmetries of the domain and often form groups with rich geometric and analytic structure. The group of automorphisms of $\mathcal{D}$ is denoted by $\text{Aut}(\mathcal{D})$.

They are also central in understanding the celebrated uniformization theorem, which we anticipate here and will return to in more detail later.

Theorem 26.142 (Uniformization Theorem: simply connected case). Every simply connected Riemann surface is conformally equivalent to exactly one of the following three domains:

- (1) the Riemann sphere $\hat{\mathbb{C}}$,
- (2) the complex plane $\mathbb{C}$,
- (3) the unit disk $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$.

Theorem 26.143 (Uniformization Theorem: general form). Every Riemann surface X admits a universal covering surface $\tilde{X}$ which is simply connected.⁸ By the simply connected case, $\tilde{X}$ is conformally

⁸A *universal covering surface* of a Riemann surface X is a simply connected Riemann surface $\tilde{X}$ together with a surjective holomorphic map $\pi : \tilde{X} \rightarrow X$ such that for every $p \in X$ there exists a neighborhood U of p with $\pi^{-1}(U)$ a disjoint union of open sets, each mapped biholomorphically onto U by π . It is unique up to conformal equivalence, and any other covering surface of X is a quotient of $\tilde{X}$.

equivalent to exactly one of the three canonical surfaces above. Moreover, X is conformally equivalent to a quotient

$$X \cong \tilde{X}/\Gamma,$$

where Γ is a properly discontinuous group of automorphisms of $\tilde{X}$.




FIGURE 20. The covering space of an annulus (adapted from Wikimedia Commons).

Hence, before turning to the general theory, let us recall some basic examples of automorphism groups: those of the complex plane, the Riemann sphere, and the unit disk.

26.2. Automorphisms of the plane.

Theorem 26.144. *The group of biholomorphic self-maps of $\mathbb{C}$ is exactly the affine group $z \mapsto az + b : a \in \mathbb{C} \setminus 0, b \in \mathbb{C}$.*

Proof. The maps $z \mapsto az + b$ with $a \neq 0$ are entire and bijective with holomorphic inverse, so they lie in $\text{Aut}(\mathbb{C})$. For the converse, let $f \in \text{Aut}(\mathbb{C})$. After a translation we may assume $f(0) = 0$.

Because f is a homeomorphism, its inverse is continuous at 0. Hence there exists $c > 0$ such that

$$|w| < c \implies |f^{-1}(w)| < 1.$$

Equivalently,

$$(26.6) \quad |z| \geq 1 \implies |f(z)| \geq c.$$

Consider $h(z) = f(1/z)$ on $0 < |z| < \infty$. If 0 were an essential singularity of h , then by the Casorati–Weierstrass theorem the image of every punctured neighborhood of 0 would be dense in $\mathbb{C}$, contradicting injectivity. Thus ∞ is either a removable singularity or a pole of f .

If ∞ were a removable singularity of f , then f would be bounded and hence, by Liouville's theorem, be constant—contra-dicting injectivity. Thus ∞ is a pole of f , so f is a polynomial (cf. Exercise 6.43). By the Fundamental Theorem of Algebra, an injective polynomial must have degree 1, and therefore $f(z) = az + b$ with $a \neq 0$. ■

26.3. The Riemann sphere. The Riemann sphere is a model of the extended complex plane (also called the closed complex plane): the complex plane plus one point at infinity. This extended plane represents the extended complex numbers, that is, the complex numbers together with the value ∞ . Analytic functions on $\hat{\mathbb{C}}$ are defined using bi-holomorphic charts.




FIGURE 21. The Riemann sphere (adapted from Wikimedia Commons).

26.4. Linear fractional transformations (LFTs): a first view.

A linear fractional transformation, or Möbius transformation, is a map of the form

$$(26.7) \quad L(z) = \frac{az + b}{cz + d}, \quad ad - bc \neq 0,$$

which is meromorphic on $\hat{\mathbb{C}}$ with a single pole at $z = -d/c$.

Exercise 26.145. Associate to an LFT its coefficient matrix

$$(26.8) \quad \hat{M}(L) := \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

If L_1 and L_2 are LFTs, show that

$$(26.9) \quad \hat{M}(L_1 \circ L_2) = \hat{M}(L_1)\hat{M}(L_2),$$

Hence LFTs with composition are (projectively) isomorphic to $\mathrm{GL}(2, \mathbb{C})$ with multiplication.⁹

26.5. Automorphisms of the Riemann sphere.

Theorem 26.146. A map $F : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ is analytic if and only if F is a rational function.

Proof. If F is analytic on $\hat{\mathbb{C}}$, then it has no essential singularities anywhere on the sphere, though it may have poles (since ∞ is a regular point of $\hat{\mathbb{C}}$). In particular, the restriction of F to $\mathbb{C} = \hat{\mathbb{C}} \setminus \{\infty\}$ is meromorphic.

Because $\hat{\mathbb{C}}$ is compact, the set of poles must be finite—otherwise the poles would have an accumulation point (necessarily at ∞), contradicting analyticity there (why?). Let P be a polynomial whose zeros are exactly the poles of F (counted with multiplicity). Then PF is entire on $\mathbb{C}$.

Consider the function h defined on a punctured neighborhood of 0 in $\mathbb{C}$ by

$$h(z) = P(1/z) F(1/z).$$

Since F is analytic at ∞ , h has at most a pole at 0. Equivalently, the Laurent expansion of h about 0 has only finitely many negative powers of z ; otherwise PF would have an essential singularity at ∞ . Thus

$$PF(z) = \sum_{j=-m}^n c_j z^j, \quad m, n \geq 0.$$

Analyticity of PF at $z = 0$ forces $m = 0$, so PF is a polynomial Q . Hence $F = Q/P$ is a rational function.

Conversely, any rational function $F = Q/P$ is meromorphic on $\mathbb{C}$ and has at most a pole at ∞ , so it defines an analytic map on $\hat{\mathbb{C}}$. ■

Theorem 26.147. The automorphism group of $\hat{\mathbb{C}}$ is precisely the Möbius group:

$$\mathrm{Aut}(\hat{\mathbb{C}}) = \left\{ z \mapsto \frac{az + b}{cz + d} : ad - bc \neq 0 \right\}.$$

Proof. Every Möbius map is a rational function with at most one simple pole, and its inverse is again a Möbius map, so it is clearly an automorphism of $\hat{\mathbb{C}}$.

⁹Multiplying all entries of a 2×2 matrix by the same nonzero scalar does not change the associated linear fractional transformation. Thus what really matters is the matrix up to overall scaling. This is what is meant by “projectively isomorphic” here.

Conversely, let f be an automorphism of $\hat{\mathbb{C}}$. By Theorem 26.146, $f = P/Q$ with P, Q polynomials having no common factors. Since f is bijective, it can have at most one pole, and that pole must be simple; hence $Q(z) = cz + d$ is linear.

If $\deg P \geq 2$, then, by the Fundamental Theorem of Algebra (Theorem 7.45), the equation $f(z) = 0$ has $\deg P$ solutions (counting multiplicities). Hence $\deg P = 1$, so $P(z) = az + b$ is linear.

Finally, $ad - bc \neq 0$, otherwise f would be constant. Thus f is a Möbius map. ■

27. LINEAR FRACTIONAL TRANSFORMATIONS (MÖBIUS TRANSFORMATIONS)

Linear fractional transformations were defined in (26.7).

If $c = 0$ we have a linear function. If $c \neq 0$ we write

$$(27.1) \quad \frac{az + b}{cz + d} = \frac{a}{c} - \frac{ad - bc}{c^2(z + d/c)}; \quad z \neq -d/c$$

It is clear from (27.1) that $S(z_1) = S(z_2)$ iff $z_1 = z_2$ and in particular $S'(z) \neq 0$: If a LFT L is defined in the domain $\mathcal{D}$ (that is, $-d/c \notin \mathcal{D}$), then $\mathcal{D}$ and $L(\mathcal{D})$ are conformally equivalent. It also follows from the decomposition (27.1) that

Proposition 27.1. The Möbius group is generated by $z \mapsto z + a$, $z \mapsto az$ and $z \mapsto 1/z$. This implies that $\text{PSL}(2, \mathbb{C})$ is generated by the matrices of the form

$$(27.2) \quad \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

(we took a negative sign in the last matrix to have determinant one.)

Hence, the automorphisms of $\hat{\mathbb{C}}$ correspond to $\text{PSL}(2, \mathbb{C})$ while those of $\mathbb{C}$ correspond to the subgroup generated by the matrices $\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix}$.

Exercise 27.2. Show that the inversion $z \mapsto 1/z$ maps a line or a circle into a line or a circle. Hint: Show first that the equation

$$\alpha z \bar{z} + \beta z + \bar{\beta} \bar{z} + \gamma = 0$$

where $\alpha, \gamma \in \mathbb{R}$ and $\beta \in \mathbb{C}$ and $|\beta|^2 > \alpha\gamma$ is the most general equation of a line or a circle. Then apply the inversion to the equation.

As a result we have an important property of LFTs

Proposition 27.3. A LFT maps a line or a circle into a line or a circle.

Proof. By Exercise 27.2 inversion has the property above, and clearly, so do Euclidean transformations. $\square$

27.1. Finding specific LFTs. A line is determined by two points and a circle is determined by three. We now show that for any two circles or lines there exists a LFT mapping one into the other, and we describe how to find it explicitly.

Let z_1, z_2, z_3 be three distinct points in $\mathbb{C}$. Then the transformation

$$(27.3) \quad S(z) = \frac{z_1 - z_3}{z_1 - z_2} \frac{z - z_2}{z - z_3}$$

maps z_1, z_2, z_3 into $1, 0, \infty$ in this order; this is straightforward to check. If one of z_1, z_2, z_3 is ∞ , we pass to the limit. For example, the transformation taking (∞, z_1, z_2) to $(1, 0, \infty)$ is

$$(27.4) \quad L(z) = \frac{z - z_1}{z - z_2}.$$

To find a transformation that maps z_1, z_2, z_3 into $\tilde{z}_1, \tilde{z}_2, \tilde{z}_3$ in this order, we may write it as $\tilde{L} = \tilde{L}^{-1} \circ L$, where L maps (z_1, z_2, z_3) to $(1, 0, \infty)$ and $\tilde{L}$ maps $(\tilde{z}_1, \tilde{z}_2, \tilde{z}_3)$ to $(1, 0, \infty)$. By Exercise 27.4, this transformation is unique.

Exercise 27.4. Show that a LFT that takes $(1, 0, \infty)$ into itself is the identity.

27.1.1. Cross ratio. If z_i , $i = 1, \dots, 4$ are four distinct points and $w_i = L(z_i)$, then (check!)

$$\frac{w_1 - w_2}{w_1 - w_3} \cdot \frac{w_3 - w_4}{w_2 - w_4} = \frac{z_1 - z_2}{z_1 - z_3} \cdot \frac{z_3 - z_4}{z_2 - z_4}.$$

The quantity

$$[z_1, z_2; z_3, z_4] = \frac{z_1 - z_2}{z_1 - z_3} \cdot \frac{z_3 - z_4}{z_2 - z_4}$$

is called the *cross ratio* of the four points. It is invariant under Möbius transformations. This provides a handy way to determine the image of a fourth point once a transformation has been fixed using three points.

27.2. Mappings of regions. We know that LFTs are *conformal and one-to-one* and transform circles and lines onto circles and lines. Proposition 26.141 shows that LFTs map disks and half-planes to disks and half-planes. To see the latter, note that on the Riemann sphere LFTs map circles to circles, hence disks to disks as well. A rigorous proof requires verifying that the interior/exterior of a circle are mapped to

interiors/exterior respectively — we leave this as a straightforward exercise.

Exercise 27.5. (i) Show that the *Cayley transform*

$$C(z) = \frac{z - i}{z + i}$$

maps the upper half-plane (UHP) conformally onto the unit disk $\mathbb{D}$. Its inverse

$$C^{-1}(w) = i \frac{1 + w}{1 - w}$$

maps $\mathbb{D}$ conformally back onto the UHP.

- (ii) Find an LFT that maps the disk $(x - 1)^2 + (y - 2)^2 = 4$ onto the unit disk and sends the center $(1, 2)$ to $i/2$.
 (iii)* Find the most general LFT that maps the unit disk onto itself.

Hint: The general solution has the form

$$z \mapsto e^{i\theta} \frac{z - a}{1 - \bar{a}z}, \quad |a| < 1, \theta \in \mathbb{R}.$$

27.3. The Schwarz lemma. As usual, we let $\mathbb{D}$ denote the open unit disk.

Theorem 27.6 (Schwarz lemma). Let $f : \mathbb{D} \rightarrow \mathbb{D}$ be analytic with $f(0) = 0$. Then:

- (i) For all $z \in \mathbb{D}$,

$$(27.5) \quad |f(z)| \leq |z|.$$

- (ii) If equality in (27.5) holds for some nonzero $z_0 \in \mathbb{D}$, then there exists $\phi \in \mathbb{R}$ such that $f(z) = e^{i\phi}z$.
 (iii) We have $|f'(0)| \leq 1$, with equality if and only if $f(z) = e^{i\phi}z$ for some $\phi \in \mathbb{R}$.

Proof. (i) Since $f(0) = 0$, the quotient $f(z)/z$ extends analytically to all of $\mathbb{D}$ (by defining its value at 0 to be $f'(0)$). Because $f(\mathbb{D}) \subset \mathbb{D}$, the maximum modulus principle gives

$$\left| \frac{f(z)}{z} \right| \leq \limsup_{r \uparrow 1} \max_{|z|=r} \left| \frac{f(z)}{z} \right| \leq 1,$$

which implies (27.5).¹⁰

(ii) Suppose equality holds at some nonzero z_0 . Then $|f(z_0)/z_0| = 1$, so the analytic function $f(z)/z$ attains its maximum modulus inside $\mathbb{D}$.

¹⁰You can show that $\limsup$ can be replaced by $\lim$ based on the monotonicity implied by the maximum principle.

By the maximum modulus principle, $f(z)/z$ must be constant, say C . Thus $f(z) = Cz$, and since $|C| = 1$, we may write $C = e^{i\phi}$ with $\phi \in \mathbb{R}$.

(iii) The bound $|f'(0)| \leq 1$ follows immediately by taking the limit $z \rightarrow 0$ in (27.5). If equality holds, write $f(z) = e^{i\phi}z + a_m z^{m+1} + \dots$ with $m \geq 1$. Then

$$\frac{f(z)}{z} = e^{i\phi}(1 + a_m z^m + \dots).$$

Choosing $z = \varepsilon e^{-i \arg(a_m)/m}$ with ε small, we obtain $|f(z)| > |z|$, contradicting (i). Hence $f(z) \equiv e^{i\phi}z$. $\square$

Corollary 27.7. If h is an automorphism of the unit disk and $h(0) = 0$, then

$$h(z) = e^{i\phi}z \quad \text{for some } \phi \in \mathbb{R}.$$

Proof. By Theorem 27.6, we have $|h(z)| \leq |z|$ for all $z \in \mathbb{D}$. Since h is an automorphism, its inverse h^{-1} is also an automorphism of $\mathbb{D}$ and satisfies $h^{-1}(0) = 0$. Applying Theorem 27.6 to h^{-1} gives $|h^{-1}(w)| \leq |w|$ for all $w \in \mathbb{D}$. In particular, with $w = h(z)$ we obtain

$$|z| = |h^{-1}(h(z))| \leq |h(z)|.$$

Combining this with the first inequality, we deduce $|h(z)| = |z|$ for all $z \in \mathbb{D}$. By Theorem 27.6(ii), it follows that $h(z) = e^{i\phi}z$ for some $\phi \in \mathbb{R}$. $\square$

27.4. Automorphisms of the unit disk. We have seen that the subgroup of LFTs

$$(27.6) \quad L(z) = e^{i\phi} \frac{z - \alpha}{1 - \bar{\alpha}z}, \quad \phi \in \mathbb{R}, \quad |\alpha| < 1,$$

maps the unit disk one-to-one onto itself.

The converse is also true:

Theorem 27.8. Every automorphism f of $\mathbb{D}$ is of the form (27.6), with $\alpha = f^{-1}(0)$.

Proof. Let $\alpha = f^{-1}(0)$. Consider the map $S(z) = \frac{z - \alpha}{1 - \bar{\alpha}z}$, which sends α to 0. Then $h := S \circ f^{-1}$ is an automorphism of $\mathbb{D}$ satisfying $h(0) = 0$. By Corollary 27.7, we must have $h(z) = e^{i\phi}z$ for some $\phi \in \mathbb{R}$. Thus f is of the form (27.6). $\square$

Exercise 27.9. Show that the automorphisms of the upper half-plane $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ are precisely the LFTs

$$z \mapsto \frac{az + b}{cz + d}, \quad a, b, c, d \in \mathbb{R}, \quad ad - bc > 0.$$

- (i) Verify directly that such maps preserve $\mathbb{H}$ and are bijective.
- (ii) Show that if $\varphi \in \text{Aut}(\mathbb{H})$ satisfies $\varphi(0) = 0$ and $\varphi(1) = 1$, then $\varphi(z) = z$.
- (iii) Conclude that this subgroup of LFTs is isomorphic to $\text{PSL}(2, \mathbb{R})$.

Remark 27.10. The automorphism group of $\mathbb{D}$ acts *transitively* on $\mathbb{D}$: for any $z_1, z_2 \in \mathbb{D}$ there exists an automorphism sending z_1 to z_2 . In fact, $\text{Aut}(\mathbb{D})$ acts even more strongly on the boundary: it is *triply transitive* on $\partial\mathbb{D}$, meaning that any three distinct boundary points can be mapped to any other three distinct boundary points. This illustrates the very rich symmetry of the unit disk.

Remark 27.11. We see that the automorphism groups of all three “canonical” simply connected Riemann surfaces (cf. Theorem 26.142) are realized as subgroups of $\text{PSL}(2, \mathbb{C})$:

- For the Riemann sphere $\hat{\mathbb{C}}$, $\text{Aut}(\hat{\mathbb{C}}) = \text{PSL}(2, \mathbb{C})$.
- For the complex plane $\mathbb{C}$, the automorphisms are the affine maps $z \mapsto az + b$ with $a \in \mathbb{C} \setminus \{0\}$, $b \in \mathbb{C}$, corresponding to the subgroup of upper-triangular matrices in $\text{PSL}(2, \mathbb{C})$.
- For the unit disk $\mathbb{D}$, the automorphisms are the so called *Blaschke factors*

$$z \mapsto e^{i\phi} \frac{z - \alpha}{1 - \bar{\alpha}z}, \quad |\alpha| < 1, \phi \in \mathbb{R},$$

forming a subgroup of $\text{PSL}(2, \mathbb{C})$.

Thus, in each canonical case the symmetry group of the surface embeds naturally into $\text{PSL}(2, \mathbb{C})$.

28. SOME SPECIAL BIHOLOMORPHIC TRANSFORMATIONS

Many elementary conformal maps are compositions of Möbius maps, ramifications ($z \mapsto z^\alpha$), exponential and trigonometric maps, and other simple transformations such as the Joukowski map. We illustrate below a number of useful transformations; see [1], [12] for further examples. A good number of interesting domains can be mapped to the unit disk using combinations of these transformations. Note that by Proposition 26.141 it suffices to examine carefully how the boundaries are mapped in order to understand the action of a map on a domain.

1. **Möbius transformations.** We have already discussed these in detail. A particularly useful special case is the Cayley transform

$$C(z) = \frac{z - i}{z + i},$$

which maps the upper half-plane conformally onto $\mathbb{D}$.

2. Power maps. The map $z \mapsto z^{1/\alpha}$ transforms sectors into half-planes. See Fig. 22.




FIGURE 22. The map $z \mapsto z^{1/\alpha}$ maps the open sector of opening $\pi\alpha$ conformally onto the open upper half-plane. The boundary point 0 maps to 0, and the upper boundary of the sector is rotated to $-\mathbb{R}$.

3. Maps of the cut plane.




FIGURE 23. $\mathbb{C}$ with one, two, four, and eight symmetric cuts. More generally, the directions of the cuts can be taken along the roots of unity $\omega_1, \dots, \omega_k$, $k \in \mathbb{N}$.

Exercise 28.12. (a) Show that $\phi_1(z) = \frac{4z}{(1+z)^2}$ maps conformally the unit disk onto $\mathbb{C} \setminus [1, \infty)$. Verify, using Proposition 26.141, that the two sides of $(1, \infty)$ are treated as distinct boundary arcs, as discussed in §26.

(b) Show that for any $n \in \mathbb{N}$ the map

$$\phi_n(z) = [\phi_1(z^n)]^{1/n}$$

maps $\mathbb{D}$ conformally onto $\mathbb{C}$ with symmetric cuts along ω_k , $k = 1, \dots, n$.

4. The Joukowski transformation. This map sends the region in the upper half-plane lying outside the unit circle conformally onto the upper half-plane (see Fig. 24). By rescalings and translations, one can adjust the radius or center along $\mathbb{R}^+$. It is defined by

$$(28.7) \quad J(z) = z + \frac{1}{z}.$$

Exercise 28.13. Check, by examining boundary behavior and applying Proposition 26.141, that J indeed maps the indicated region conformally onto the UHP. What is the image under J of the reflection across $\mathbb{R}$ of the yellow region?

5. Exponential map. The exponential function maps conformally the strip

$$S = \{z \in \mathbb{C} : 0 < \operatorname{Im} z < \pi\}$$

onto $\mathbb{C} \setminus (-\infty, 0]$.

Exercise 28.14. Use some of the maps discussed above to construct a conformal map from $\mathbb{D}$ onto the strip S .

5. The cosine maps the region $\{z \mid \operatorname{Im} z > 0, 0 < x < \pi\}$ conformally to the UHP.

Exercise 28.15. Note that the cosine and the exponential are linked via a Joukowski map. Can you explain the mapping of $\cos$ based on **5.** and J ?

Note 28.16. Keep in mind that the conformal maps above **are not tied to the domains presented**, except loosely through the fact that these domains are often chosen to be some maximal region of conformality. Often subdomains of a maximal domain are interesting in applications.




FIGURE 24. Velocity field lines in Joukovski's domain, the domain in the UHP above the half-circle.

Application: Ideal flow above a semicircular bump (normalized case $U = 1$). We consider irrotational, incompressible, inviscid flow in the upper half-plane above the semicircle $\{|z| = 1, \operatorname{Im} z \geq 0\}$. The boundary condition is free-slip (no penetration). As $x \rightarrow \pm\infty$, the flow approaches the uniform stream $(v, u) = (1, 0)$; see Fig. 24.

Complex potential. The Joukowski map $w = J(z) = z + \frac{1}{z}$ flattens the boundary, sending the upper half-plane above the semicircle to the upper half w -plane, with the boundary mapping to the real line. In the w -plane, the uniform stream with unit speed has potential

$$F(w) = w.$$

Therefore the complex potential in the z -plane is

$$W(z) = F(J(z)) = z + \frac{1}{z}.$$

Velocity field. The complex velocity is

$$\frac{dW}{dz} = 1 - \frac{1}{z^2}.$$

Thus, writing $z = x + iy$,

$$V_x(x, y) = v(x, y) = 1 - \frac{x^2 - y^2}{(x^2 + y^2)^2}, \quad V_y(x, y) = u(x, y) = \frac{2xy}{(x^2 + y^2)^2}, \quad |z| > 1, \operatorname{Im} z > 0.$$

Potential and stream function. The potential and stream function are

$$\phi(x, y) = \operatorname{Re} W(z) = x + \frac{x}{x^2 + y^2}, \quad \psi(x, y) = \operatorname{Im} W(z) = y - \frac{y}{x^2 + y^2}.$$

Boundary conditions. On the boundary $|z| = 1$, one checks $\psi = 0$, so the semicircle and real axis are streamlines (no penetration). As $x \rightarrow \pm\infty$, $(v, u) \rightarrow (1, 0)$, the uniform flow. Stagnation points occur at $z = \pm 1$, where $W'(z) = 0$.

Other common conformal maps are depicted in the following figures. The Cayley transform

$$\frac{z - i}{z + i}$$

maps conformally the upper half-plane (UHP) onto $\mathbb{D}$. It plays an important role in the theory of unbounded operators. Any other function f that maps conformally the UHP to $\mathbb{D}$ must be the Cayley transform composed on the left with an automorphism of the disk.

Exercise 28.17. Write the general form of such a map.




FIGURE 25. A conformal mapping example.

- Exercise 28.18.**
- (1) Check the transformations illustrated in the figures above.
 - (2) Draw a similar picture for the mapping $\sin z$ from the upper half-strip bordered by the half-lines $x = \pm\pi/2$, $y > 0$.




FIGURE 26. Mapping between a circle and a half-plane.




FIGURE 27. Another conformal mapping illustration.

- (3) Find a conformal map of the quarter disk $|z| < 1$, $\arg(z) \in (0, \pi/2)$ onto the upper half-plane.
- (4) Find a conformal map of the half-disk $|z| < 1$, $\arg(z) \in (0, \pi)$ onto the half-strip $x < 0$, $y \in (0, \pi)$.
- (5) Find a conformal map of the right half-plane (RHP) with a cut along $[0, 1]$ in the RHP.

It is useful to remark that we can find linear fractional transformations that map conformally a region between two circles into a half-plane (or disk) using very simple transformations. As an illustration, let us map the “moon crescent” M in Fig. 28 onto a half-plane.

The equations of the bounding circles are

$$x^2 + y^2 = 16, \quad (x - 3)^2 + y^2 = 9.$$




FIGURE 28. Mapping a crescent-shaped domain to a half-plane.

Solving for their intersection points $a_{\pm}$ we obtain

$$a_{\pm} = \frac{8}{3} \pm \frac{4i\sqrt{5}}{3}.$$

The angle between the two circles is the same as the angle between the tangent vectors at the intersection, which can be computed at a_+ as

$$\arg\left(\frac{a_+ - 3}{a_+}\right) = \arctan\left(\frac{\sqrt{5}}{2}\right) =: \phi_0.$$

The key idea is that if a Möbius map sends one of the intersection points to infinity, the circular arcs are mapped to straight rays, which simplifies the geometry of the domain. Specifically, if we choose an LFT sending

$$a_- \mapsto 0, \quad 0 \mapsto 1, \quad a_+ \mapsto \infty,$$

then both arcs become rays (since they meet at ∞). One of them is mapped to the positive real axis $\mathbb{R}^+$, while the other is mapped to a ray making angle ϕ_0 with $\mathbb{R}^+$. Such a transformation is for instance

$$(28.8) \quad S(z) = \frac{a_+}{a_-} \frac{z - a_-}{z - a_+}.$$

Finally, to straighten the resulting sector of opening ϕ_0 into the upper half-plane, we apply the power map

$$z \mapsto z^{\pi/\phi_0}.$$

- Exercise 28.19.** (1) Verify the coordinates $a_{\pm}$ of the intersection points.
 (2) Check that the angle between the circles at a_+ equals $\phi_0 = \arctan(\sqrt{5}/2)$.
 (3) Confirm that under the Möbius transformation (28.8) the small arc becomes $\mathbb{R}^+$ and the larger one becomes a ray at angle ϕ_0 .
 (4) Show that the ramified map $z \mapsto z^{\pi/\phi_0}$ sends the resulting sector conformally onto the upper half-plane.




FIGURE 29. Color density plot of the heat equation solution for a quarter disk

Example 2. Solve $\Delta u = 0$ in the quarter disk

$$\Omega = \{ |z| < 1, \arg(z) \in (0, \frac{\pi}{2}) \}$$

with boundary conditions: $u = 1$ on the circular arc, and $u = 0$ on the two radii.

Solution. The strategy is to find a conformal map of Ω onto a horizontal strip

$$S = \{ w = x + iy : 0 < y < 1 \}$$

such that the circular arc maps to the line $y = 1$, while the radii map to the line $y = 0$. In the strip S , the harmonic solution of the boundary problem is immediate:

$$u(w) = \operatorname{Im} w.$$

We then pull back u through the conformal transformations.

Step 1. The map $z \mapsto z^2$ sends the quarter disk onto the upper half-disk $\{|z| < 1, \operatorname{Im} z > 0\}$. On the boundary we still have $u = 1$ on the arc, $u = 0$ on the interval $[-1, 1]$.

Step 2. Map the upper half-disk to the first quadrant by a Möbius transformation sending $z = 1$ to ∞ :

$$z \mapsto i \frac{1+z}{1-z}.$$

This sends the interval $[-1, 1]$ to $\mathbb{R}^+$ and the semicircle to $i\mathbb{R}^+$, so the quadrant is obtained with boundary data $u = 0$ on $\mathbb{R}^+$ and $u = 1$ on $i\mathbb{R}^+$.

Step 3. The squaring map $z \mapsto z^2$ opens the quadrant into the upper half-plane $\{\operatorname{Im} z > 0\}$, with boundary values $u = 0$ on $\mathbb{R}^+$ and $u = 1$ on $\mathbb{R}^-$.

Step 4. Finally, the logarithm maps the upper half-plane to a horizontal strip. Specifically,

$$w = \frac{2}{\pi} \ln \left(\frac{1+z^2}{1-z^2} \right)$$

maps Ω conformally onto the strip $\{0 < \operatorname{Im} w < 1\}$, with the desired correspondence of boundary parts.

Thus the solution is

$$u(z) = \operatorname{Im} \left(\frac{2}{\pi} \ln \frac{1+z^2}{1-z^2} \right),$$

or, explicitly,

$$u(x, y) = 2 - \frac{2}{\pi} \arctan \left(\frac{4xy}{-1 + (x^2 + y^2)^2} \right).$$

Exercise* The temperature distribution also satisfies Laplace's equation. (1) Map M onto a strip as in Example 2. What is the distribution of temperature in the domain M if the temperature on the larger arc is 1 and 0 on the smaller one? What shape do the lines of constant temperature have?

(2) What is the distribution of temperature in the domain and with the boundary conditions described in example 2? Draw an approximate picture of the constant temperature lines.

29. THE RIEMANN MAPPING THEOREM

Using elementary transformations we can conformally map a limited family of domains onto $\mathbb{D}$; with the maps we used in the previous section, the boundary is always very simple. The Schwarz–Christoffel formulas in §30, generally nonelementary, provide conformal maps between any polygons and the upper half plane (which, as we saw, can be mapped onto the unit disk). For polygons with more than four sides, some numerical calculations are needed. Conformally mapping more general domains, even in a computer-assisted fashion, is a difficult problem.

In principle, however, **any simply connected domain other than $\mathbb{C}$** can be mapped onto $\mathbb{D}$:

Theorem 29.20 (Riemann Mapping Theorem). Given any simply connected domain $\mathcal{D}$ other than $\mathbb{C}$, there exists a biholomorphic map f between $\mathcal{D}$ and $\mathbb{D}$.

Normalizing f by $f(z_0) = 0$ and $f'(z_0) \in \mathbb{R}^+$ for some $z_0 \in \mathcal{D}$, such a map is unique.

Riemann stated this theorem in 1851 for domains with piecewise regular boundary, and gave a proof based on solving a Dirichlet problem for the Laplacian. The method, *Dirichlet's principle*, contained a gap: the existence of a minimizer for the associated variational problem was not established. This gap was pointed out by Weierstrass in 1870.

In full generality, the theorem finally received a rigorous proof in 1900 by Osgood. Further important proofs were given by Koebe (1907) and Carathéodory (1912).

The next two pages, reproduced from Greene and Kim, *Complex Analysis*, describe Riemann's original proof in detail.

holomorphic function F on the set with $F' = f$. We shall say then that U is *holomorphically simply connected*.

It is easy to show that topological simple connectivity as defined implies the holomorphic antiderivative property just described. The theorem itself in the following form will show among other things the converse, namely, that the holomorphic simple connectivity implies topological simple connectivity.

Theorem 1.1 (Riemann) *Suppose that U is a connected open subset of $\mathbb{C}$ with $U \neq \mathbb{C}$. If U is holomorphically simply connected, then U is biholomorphic to the unit disc, i.e., there is a one-to-one holomorphic function from U onto the unit disc $D = \{z \in \mathbb{C} : |z| < 1\}$.*

In the proof of this result, it will be useful to be able to assume that U is bounded. For this, we recall the familiar fact that such a U as in the theorem is always biholomorphic to a bounded open set. The proof of this in summary form goes like this. Since $U \neq \mathbb{C}$, we can replace U by a translate to suppose that $0 \notin U$. The function $1/z$ is then holomorphic on U and hence has an antiderivative $L(z)$, say. Changing L by an additive constant will arrange that $\exp(L(z)) = z$ (this is the usual process for finding complex logarithms). Then $\exp(L(z)/2)$ is one-to-one on U . Choose an open disc in the image of $\exp(L(z)/2)$. The negative of this disc is disjoint from the image of $\exp(L(z)/2)$. So the image of $\exp(L(z)/2)$, which is biholomorphic to U , is itself biholomorphic to a bounded open subset of $\mathbb{C}$, via a linear fractional transformation.

Note that it is not clear by definition that the holomorphic simple connectivity is preserved by a biholomorphic mapping since the meaning of taking the derivative is different when the coordinates change; but by the complex chain rule this is a matter of a holomorphic factor which can be assimilated into the original function. Checking the details of this is left to the reader as an exercise.

So now we can assume without loss of generality that the open set U is bounded. And by translation we can now assume $0 \in U$. We shall look for a biholomorphic mapping from U to the unit disc D which takes 0 to 0. Of course, if there is a biholomorphic map from U to the unit disc at all, there is one that takes 0 to 0 since a linear fractional transformation taking the unit disc D to itself will take any given point to the origin, and in particular the image of 0 to begin with can be moved to the origin.

Now if $H:U \rightarrow D$ is biholomorphic and has $H(0) = 0$, then $H(z)/z$ has a removable singularity at 0. Hence, H can be written as $zh(z)$, where h is holomorphic on U and $h(0) \neq 0$. That $h(0) \neq 0$ follows because H is supposed to be one-to-one and hence must have derivative vanishing nowhere. Of course, $h(z)$ is also nonzero for every other $z \in U$ because 0 is the only point of U with $H(z) = 0$.

Now there is an antiderivative L of h'/h on U . The product $h(z) \exp(-L(z))$ is constant since it has derivative identically equal to 0 and U is connected. Changing L by an additive constant, we can assume $h(z) \exp(-L(z)) = 1$ for all $z \in U$. (This familiar argument will occur several times here).

The essential point of Riemann's method was to consider the harmonic function $\operatorname{Re} L(z)$. This is of course equal to $\ln |h(z)|$. Since the "boundary values" of $|H(z)|$ have to be 1, it must be that the boundary value of $|h|$ at a boundary point z_0 of U has to be

$1/|z_0|$. In particular, the harmonic function $\ln |h|$ has to have boundary value at z_0 equal to $-\ln |z_0|$.

At this point, Riemann appealed to what he referred to as the Dirichlet Principle. The Euler–Lagrange equation for the variational problem of minimizing the so-called Dirichlet (energy) integral for a real-valued function $f(x, y)$, namely minimizing this integral

$$\int_U \left[\left(\frac{\partial f}{\partial x} \right)^2 + \left(\frac{\partial f}{\partial y} \right)^2 \right] dx dy$$

under the condition that $f = g$ on the boundary ∂U of U , is easily computed to satisfy $\Delta f = 0$ (§18 of [10, 11]).

So Riemann proposed that the harmonic function with the boundary values $-\ln |z_0|$ at each boundary point z_0 could be found by minimization of the Dirichlet integral. And Riemann was well aware of how to construct h and hence H from knowing $\ln |h(z)|$. Riemann actually expressed this all in terms of $\ln |H|$ and the idea of Green's function, a function with boundary value 0 and a specified singularity at (in our case) the point 0, namely the function had to be of the form $\ln |z| + u(z)$ with u harmonic near the point 0. This is equivalent for open sets in $\mathbb{C}$ to our discussion, though the Green's function notion is useful when one tries to extend the Riemann Mapping Theorem to the uniformization problem where there is no *a priori* global z -coordinate.

The main difficulty is that there is no particular reason to suppose that there is in fact any minimum for the Dirichlet integral in this situation. There is also a less serious difficulty of explaining why the resulting function is one-to-one and onto—intuitively this is just a matter of winding numbers if one can approximate U from inside by domains with smooth or piecewise-smooth closed curve boundaries. One supposes that Riemann may have taken this part for obvious, though it is actually quite subtle if one does not appeal to any pre-existing topological intuitions. We shall give a precise argument later on. Riemann apparently considers only domains, the boundary of which is smooth in some sense. Osgood made the major forward step treating simply connected open sets in general, thus proving what we call today the Riemann Mapping Theorem. The Osgood proof is acknowledged directly by Carathéodory [2] where the ideas involved in the usual proof of today, via normal families, are presented. See the footnote (**) of page 108 of [2]. But, for some reason, Osgood's proof fell from favor or even recognition for the history of the Theorem in [9]; there is a reference to Osgood's paper but no comment on it, no acknowledgment that this is in fact the reasonably complete first proof of the general result.

2 The application of the Perron method

Even simply connected bounded open sets in $\mathbb{C}$ can have complicated boundaries. The boundary of the Koch snowflake, for example, has Hausdorff dimension greater than 1 [15]. And Osgood [6] already gave an example with boundary having positive (2-dimensional) measure. Thus, it is appropriate to introduce carefully what is to be meant by finding functions with specified boundary values. For this purpose, let U be a bounded open set in $\mathbb{C}$ and ∂U be its boundary, that is the complement of U within the closure $\text{cl}(U)$ of U in $\mathbb{C}$, or equivalently the intersection of $\text{cl}(U)$ with $\mathbb{C} - U$. Suppose that $b: \partial U \rightarrow \mathbb{R}$ is a continuous function. Then we say that a harmonic function $h: U \rightarrow \mathbb{R}$ is

29.1. Equicontinuity and Normal Families. We look at functions $f : M \rightarrow M'$ where M, M' are metric spaces. Recall that if the metrics are d and d' , a function is *uniformly continuous* if

$$(29.9) \quad \forall \varepsilon > 0 \exists \delta > 0 \left(\forall (z, z_0) \in M^2, d(z, z_0) < \delta \Rightarrow d'(f(z), f(z_0)) < \varepsilon \right).$$

We can assume that the metric d' is a **bounded function**, for we can always replace it by

$$d'' = \frac{d'}{1 + d'},$$

which is a metric topologically equivalent to d' , with $d'' \leq 1$. (This standard trick allows us to avoid unbounded distances.)

Definition 29.21 (Equicontinuity). *A family $\mathcal{F}$ of continuous functions $f : M \rightarrow M'$ is equicontinuous if for every $x \in M$ and every $\varepsilon > 0$ there exists $\delta(x, \varepsilon) > 0$ such that*

$$(29.10) \quad d(x, y) < \delta(x, \varepsilon) \implies d'(f(x), f(y)) < \varepsilon \quad \forall f \in \mathcal{F}.$$

Definition 29.22 (Normal Families). Let M, M' be complete metric spaces. A collection of continuous functions $\mathcal{F}$ from M to M' is a **normal family** if it is pre-compact in the topology of uniform convergence on compact sets.¹¹

Exhaustion by Compact Sets. If the metric space M is σ -compact, then by definition it can be covered by a countable nested family of compact sets. For instance, in $\mathbb{C}$ or $\mathbb{R}^n$ we may take

$$(29.11) \quad M = \bigcup_{n \in \mathbb{N}} K_n, \quad K_n = \{x \in M : |x| \leq n\}.$$

Metrizability of Uniform Convergence on Compact Sets. On each K_n in (29.11) we define the distance between two functions f, g analogously to the L^∞ norm:

$$(29.12) \quad \delta_n(f, g) = \sup_{x \in K_n} d''(f(x), g(x)).$$

We then define a distance on the whole of M by

$$(29.13) \quad \rho(f, g) = \sum_{n=1}^{\infty} 2^{-n} \delta_n(f, g).$$

¹¹A subset of a metric space is called *pre-compact* (or *relatively compact*) if its closure is compact. Equivalently, every sequence in the set has a subsequence converging to a point in the ambient space.

Since $d''' \leq 1$, the sum converges. This construction metrizes the topology of uniform convergence on compact sets.

Exercise 29.23. Let M be a σ -compact metric space. Check that ρ in (29.13) is a metric on $C(M)$. Check that convergence with respect to ρ is equivalent to uniform convergence on compact sets. Finally, check that $C(M)$ is complete with respect to ρ if M' is a complete metric space.

Proposition 29.24. A family $\mathcal{F}$ is normal if and only if its closure $\overline{\mathcal{F}}$ with respect to ρ is compact in $C(M)$.

Example 29.25 (Motivation). Let $f_n : \mathbb{C} \rightarrow \mathbb{C}$ be the translates $f_n(z) = f(z+n)$ for some fixed entire function f . Even if f is bounded on $\mathbb{C}$, the family $\{f_n\}$ is not normal on $\mathbb{C}$, because the translates “escape to infinity” and there is no subsequence that converges uniformly on compact sets. This illustrates that normality is a much stronger property than pointwise boundedness or equicontinuity of each individual function.

29.1.1. *The Ascoli–Arzelà Theorem.*

Theorem 29.26 (Ascoli–Arzelà). A family $\mathcal{F}$ of continuous functions in the region $\Omega \subset \mathbb{C}$ with values in a metric space M' is **normal** in Ω if and only if the following conditions are satisfied:

- (i) $\mathcal{F}$ is equicontinuous at every $x \in \Omega$.
- (ii) $\mathcal{F}$ is equibounded: for every $z \in \Omega$ there exists a compact set $K_1 = K_1(z) \subset M'$ such that $f(z) \in K_1$ for all $f \in \mathcal{F}$.

This is a standard theorem; we leave the proof for the Appendix, see §29.26.

Proposition 29.27. Let $M' \subset \mathbb{C}$ and $\mathcal{F}$ be a normal family from Ω to M' . Let $K \subset \Omega$ be compact. Then $\mathcal{F}$ is uniformly bounded on K :

$$(29.14) \quad \sup_{z \in K, f \in \mathcal{F}} |f(z)| = m < \infty.$$

Proof. Since $\mathcal{F}$ is a normal family, by Theorem 29.26 and equicontinuity, it follows that for any point a we can find $\delta(a) > 0$ such that

$$(29.15) \quad |a - b| < \delta(a) \Rightarrow |f(a) - f(b)| < 1, \quad \forall f \in \mathcal{F}.$$

Cover K by finitely many disks $\mathbb{D}_{\delta(a_j)}(a_j)$, and let δ_0 be the minimum radius in this cover. For each j , set $m_j = \sup\{|f(a_j)| : f \in \mathcal{F}\}$ and let $m = 1 + \max_j m_j$. Then, for any $x \in K$, there exists a_j with $|x - a_j| < \delta_0$, and for all $f \in \mathcal{F}$ we have

$$(29.16) \quad |f(x)| \leq |f(x) - f(a_j)| + |f(a_j)| < 1 + m_j \leq m,$$

as desired. ■

Theorem 29.28 (Montel). Let $\mathcal{D} \subset \mathbb{C}$ be a domain and let $\mathcal{F}$ be a family of analytic functions on $\mathcal{D}$ such that for every compact set $K \subset \mathcal{D}$,

$$\sup\{|f(z)| : z \in K, f \in \mathcal{F}\} = m(K) < \infty.$$

Then $\mathcal{F}$ is a normal family on $\mathcal{D}$.

Proof. The family is clearly equibounded. We show that the derivatives are also equibounded, from which equicontinuity follows.

Let $K \subset \mathcal{D}$ be compact. Take $r < \text{dist}(K, \partial\mathcal{D})$, and cover K by finitely many disks $\mathbb{D}_r(a_j)$. Let $O = \bigcup_{j=1}^m \mathbb{D}_r(a_j)$ and set $r' = \text{dist}(\partial O, K)$. Then $\overline{O} \subset \mathcal{D}$.

By Cauchy's integral formula, for $z \in K$ we have
(29.17)

$$|f'(z)| = \left| \frac{1}{2\pi i} \oint_{\partial O} \frac{f(s)}{(s-z)^2} ds \right| \leq \frac{1}{2\pi} \cdot \frac{2\pi \cdot m(K)}{(r')^2} \cdot \text{length}(\partial O) \leq \frac{2m(K)}{r'},$$

so $|f'(z)|$ is uniformly bounded on K . Equicontinuity follows, and the result is proved. ■

29.2. The Riemann Mapping Theorem.

Definition 29.29. An injective analytic function is called *univalent* (or *schlicht*).

Theorem 29.30 (Riemann Mapping Theorem). Given any nonempty simply connected domain $\Omega \subset \mathbb{C}$ other than $\mathbb{C}$ itself, a point $z_0 \in \Omega$, and the normalization conditions $\varphi(z_0) = 0$, $\varphi'(z_0) \in \mathbb{R}^+$, there exists a unique biholomorphism $\varphi(z)$ between Ω and $\mathbb{D}$.

Note 29.31. The exclusion of $\mathbb{C}$ is necessary, since an entire bounded function must be constant by Liouville's theorem.

Note 29.32. Equivalently, any two nonempty simply connected domains different from $\mathbb{C}$ are conformally equivalent (cf. Exercise 27.9).

29.3. The Riemann Mapping Theorem.

Definition 29.33. An injective analytic function is called *univalent* (or *schlicht*).

Theorem 29.34 (Riemann mapping theorem). Given any nonempty simply connected domain $\Omega \subset \mathbb{C}$ other than $\mathbb{C}$ itself, a point $z_0 \in \Omega$ and the normalization conditions $\varphi(z_0) = 0$, $\varphi'(z_0) \in \mathbb{R}^+$ there exists a unique biholomorphism $\varphi(z)$ between Ω and $\mathbb{D}$.

Note 29.35. The fact that $\mathbb{C}$ must be an exception follows from the fact that an entire bounded function is constant.

Note 29.36. Equivalently, any two nonempty simply connected domains different from $\mathbb{C}$ are conformally equivalent. (cf. Exercise 27.9).

Proof of the Riemann Mapping Theorem

Uniqueness. This part is easier. If φ_1 and φ_2 are two functions with the stated properties, then $S := \varphi_1(\varphi_2)^{-1}$ is an automorphism of $\mathbb{D}$ and $S(0) = 0$. By Schwarz's lemma $S(z) = ze^{i\theta}$ for some θ . $S'(0) > 0$ implies $\theta = 0$.

Existence.

Note 29.37. Clearly, in the proof we can replace the arbitrary domain $\mathcal{D}$ with any set conformally equivalent to it. Therefore, we first simplify the domain by elementary transformations.

Note 29.38. By assumption there is a point in $\mathbb{C} \setminus \Omega$, which, by translation if needed, we can assume to be 0.

Note 29.39. (1) We can always replace Ω by a conformally equivalent Ω' which is bounded. Indeed, since $0 \notin \Omega$, the map $f = z \mapsto 1/z$ is bounded on Ω . $f(\Omega)$ is conformally equivalent to Ω and it is bounded. Indeed:

Lemma 29.40. Assume Ω satisfies the conditions of Theorem 29.34. Then,

(i) There is a biholomorphic branch of the log between Ω and $L = \log(\Omega)$ (with $\exp$ as its inverse, of course).

(ii) $L \cap (L + 2\pi i) = \emptyset$

(iii) There is a branch of the square root such that $\sqrt{\Omega} \cap (-\sqrt{\Omega}) = \emptyset$.

(iv) L omits an open set.

Proof. (i) In one direction, fix a in $\mathcal{D}$ and let $\text{Log}(z) = \int_1^a \frac{ds}{s} + \int_a^z \frac{ds}{s}$. Log is analytic in Ω since any closed curve in $\mathcal{D}$ can be homotopically shrunk to a point without crossing zero. Now, if $\log z_1 = \log z_2$, by taking the exponential we see that $z_1 = z_2$.

(ii) If w and $w + 2\pi i$ were both in L , then $w = \log z_1$ and $w + 2\pi i = \log z_2$. By exponentiating, we get $z_1 = z_2$ violating bijectivity of the log.

(iii) follows from (i) and (ii).

(iv) If $w_0 \in L$ together with a disk $\mathbb{D}_\varepsilon(w_0)$, then, by (ii), $2\pi i + \mathbb{D}_\varepsilon(w_0) \cap L = \emptyset$. ■

If w_1 is the center of an omitted disk, then the map $w \mapsto \frac{1}{z-w_1}$ is bounded in L .

- (2) We can assume that $\Omega \subset \mathbb{D}$. Indeed, this is simply achieved by rescaling.

Lemma 29.41. Let $z_0 \in \Omega \subset \mathbb{D}$. Then there exists a biholomorphism g of Ω onto $g(\Omega) \subset \mathbb{D}$ with $g(z_0) = 0$ and $g'(0) > 0$.

Proof. Since $\Omega \subset \mathbb{D}$, choose an automorphism ϕ of the unit disk with $\phi(z_0) = 0$. Then $g := \phi|_{\Omega}$ is a biholomorphism from Ω onto $g(\Omega) \subset \mathbb{D}$ and satisfies $g(z_0) = 0$. Finally, multiplying ϕ by a unimodular constant $e^{i\theta}$ if necessary, we may arrange $g'(z_0) > 0$. ■

We can thus assume wlog that $\Omega \subset \mathbb{D}$ and $z_0 = 0$.

Preview of the rest of the proof.

Note 29.42. (1) Let ϕ be any biholomorphism between Ω and $S \subset \mathbb{D}$. We want to understand how $\phi'(z_0)$ relates to the “size” of S , in the sense of set inclusion. Suppose for the moment that the Riemann Mapping Theorem is already known. Let $\phi_1 : \Omega \rightarrow S_1$ and $\phi_2 : \Omega \rightarrow S_2$ with $S_1 \subsetneq S_2$. For simplicity, assume $z_0 = 0$. Consider the following chains of biholomorphisms:

$$(29.18) \quad \mathbb{D} \xrightarrow{\psi} \Omega \xrightarrow{\phi_2} S_2 \xrightarrow{\chi} \mathbb{D},$$

normalized so that $\chi \circ \phi_2 \circ \psi$ vanishes at 0 with positive derivative there. By Schwarz’s lemma it must be the identity. Similarly,

$$(29.19) \quad \mathbb{D} \xrightarrow{\psi} \Omega \xrightarrow{\phi_1} S_1 \xrightarrow{\chi} \tilde{S}_1 \subsetneq \mathbb{D}.$$

The strict inclusion follows from $S_1 \subsetneq S_2$. By the chain rule and Schwarz’s lemma, we conclude that $\phi_1'(0) < \phi_2'(0)$.

Hence, our strategy will be to maximize $\phi'(z_0)$.

- (2) Next, we consider the supremum M of the values $\phi'(z_0)$ over all such maps. By compactness arguments, there exists a subsequence $\phi_n \rightarrow \phi$ with $\phi'(z_0) = M$. We then show that $\phi(\Omega) = \mathbb{D}$: if not, one can construct a new map ϕ_1 with $\phi_1'(z_0) > M$ by composing with a suitable branch of the square root, see below, contradicting maximality.
- (3) Since in $\mathbb{D}$ we have $0 < |z| < 1 \implies |z| < \sqrt{|z|} < 1$, the map $z \mapsto \sqrt{z}$ is *expanding*. If $\Omega \neq \mathbb{D}$, one can select a point outside $\phi(\Omega)$, choose a branch of the square root, and enlarge $\phi(\Omega)$ by applying this branch, thereby increasing $\phi'(z_0)$.

Definition 29.43. Let

$$\mathcal{F} = \left\{ \phi \mid \phi : \Omega \longrightarrow \phi(\Omega) \subset \mathbb{D}, \phi(0) = 0, \phi'(0) > 0, \phi \text{ biholomorphic} \right\}.$$

Proposition 29.44. (i) $M := \sup\{f'(z_0) : f \in \mathcal{F}\}$ is attained by some $F \in \mathcal{F}$.

(ii) If $F(\Omega) \neq \mathbb{D}$, then there exists $F_1 \in \mathcal{F}$ with $F'_1(z_0) > M$, a contradiction which completes the proof of the theorem.

Proof. (i) For any $f \in \mathcal{F}$ and $z \in \Omega$, we have $|f(z)| < 1$. By Montel's Theorem 29.27, $\mathcal{F}$ is a normal family. Thus, if $f_n \in \mathcal{F}$ with $f'_n(z_0) = m_n \rightarrow M$, then a subsequence converges uniformly on compacta to some F . By Weierstrass's theorem 8.49, F is analytic and $F'(z_0) = M$. Hurwitz's theorem 19.100 ensures F is injective: indeed, F cannot be constant since $F'(z_0) = M \neq 0$, and if $F(z_1) = F(z_2)$ for some $z_1 \neq z_2$, then $F(z) - F(z_1)$ would vanish in $\Omega^* = \Omega \setminus \{z_1\}$, contradicting Hurwitz.

(ii) Suppose $\mathbb{D} \setminus F(\Omega) \neq \emptyset$, and let $a \in \mathbb{D} \setminus F(\Omega)$. First apply the automorphism

$$(29.20) \quad f_1(z) = \frac{F(z) - a}{1 - \bar{a}F(z)},$$

which maps Ω biholomorphically into $\mathbb{D} \setminus \{0\}$. Hence there exists a branch of the square root on $f_1(\Omega)$, and we set

$$(29.21) \quad f_2(z) = \sqrt{f_1(z)}, \quad z \in \Omega.$$

By Lemma 29.40, f_2 is biholomorphic and $f_2(\Omega) \subset \mathbb{D}$. We have $f_2(z_0) = b = \sqrt{-a} \neq 0$. Now apply another automorphism of the disk to move b to 0 and normalize the derivative:

$$(29.22) \quad F_1(z) = \frac{f_2(z) - b}{1 - \bar{b}f_2(z)} \cdot \frac{|f'_2(z_0)|}{f'_2(z_0)}, \quad z \in \Omega.$$

A direct calculation gives

$$(29.23) \quad F'_1(z_0) = \frac{1 + |a|}{2\sqrt{|a|}} F'(z_0) > F'(z_0) = M,$$

a contradiction. ■

Note 29.45. This proof is nonconstructive: it does not provide a concrete sequence realizing the extremal map F , nor a direct procedure for obtaining F . The square-root trick ensures that any failure of surjectivity would yield a larger derivative at z_0 , but in practice the convergent sequence with $f'_n(z_0) \rightarrow M$ cannot be prescribed explicitly.

29.4. Behavior at the boundary of biholomorphisms: a general but weaker result. We now establish a weaker, but easier and useful fact [3]: if $\mathcal{D}$ is a simply connected domain and φ maps it conformally onto $\mathbb{D}$, then $\varphi(z)$ approaches $\partial\mathbb{D}$ whenever z approaches $\partial\mathcal{D}$,

in the sense defined below. This does not necessarily imply that $\varphi(z)$ converges to a boundary point.

Definition 29.1. Let $\mathcal{D}$ be a domain.

- A sequence $\{z_n\} \subset \mathcal{D}$ satisfies $z_n \rightarrow \partial\mathcal{D}$ as $n \rightarrow \infty$ if for every compact set $K \subset \mathcal{D}$ there exists n_0 such that $z_n \notin K$ for all $n > n_0$.

- Similarly, for an arc $\gamma : [0, 1] \rightarrow \mathcal{D}$ we say that $\gamma(t) \rightarrow \partial\mathcal{D}$ as $t \rightarrow 1$ if for every compact set $K \subset \mathcal{D}$ there exists $t_0 \in (0, 1)$ such that $\gamma(t) \notin K$ for all $t > t_0$.

30. BOUNDARY BEHAVIOR.

It is known, but would take us some time to prove, that *if $\mathcal{D}$ and $\mathcal{D}'$ are Jordan regions (domains bounded by Jordan curves), then the conformal map between $\mathcal{D}$ and $\mathcal{D}'$ extends to a homeomorphism between $\overline{\mathcal{D}}$ and $\overline{\mathcal{D}'}$.*

30.1. Behavior at the boundary of biholomorphisms: a general but weaker result. We derive an easy but useful result [3]: if $\mathcal{D}$ is a simply connected domain and φ maps it conformally onto $\mathbb{D}$, then $\varphi(z)$ approaches $\partial\mathbb{D}$ as z approaches $\partial\mathcal{D}$, in a sense defined below (which does not necessarily imply that $\varphi(z)$ converges).

Let $\mathcal{D}$ be a domain. Informally, a sequence or an arc approaches $\partial\mathcal{D}$ if eventually recedes away from any point in $\mathcal{D}$. The precise definition is:

Definition 30.1. A sequence $z_n \rightarrow \partial\mathcal{D}$ as $n \rightarrow \infty$ if for any compact set $K \subset \mathcal{D}$ there exists n_0 such that for all $n > n_0$ we have $z_n \notin K$. Similarly, for an arc $\gamma : [0, 1] \rightarrow \mathcal{D}$, $\gamma(t) \rightarrow \partial\mathcal{D}$ if $\forall K \exists t_0 \in (0, 1)$ s.t. $\gamma(t) \notin K$ if $t > t_0$.

Theorem 30.2. *If $\varphi : \mathcal{D} \rightarrow \mathcal{D}'$ is a domain biholomorphism and $z_n \rightarrow \partial\mathcal{D}$, as $n \rightarrow \infty$ then $\varphi(z_n) \rightarrow \partial\mathcal{D}'$ as $n \rightarrow \infty$.*

Proof. We prove the statement for sequences; the one for arcs is almost identical. Since φ is biholomorphic, any compact covering of $\mathcal{D}$ generates a compact covering of $\mathcal{D}'$ and vice-versa. Let $z_n \rightarrow \partial\mathcal{D}$ and let $K' \subset \mathcal{D}'$ be any compact set and $K = \phi^{-1}(K')$. By definition,

$$z_n \rightarrow \partial\mathcal{D} \Rightarrow \exists n_0 \text{ s.t. } \forall n > n_0 \ z_n \notin K$$

Since φ is one to one, for any $n > n_0$, $\varphi(z_n) \notin K'$ either.

Corollary 30.3. *If $\varphi : \mathcal{D} \rightarrow \mathbb{D}$ is a biholomorphism, then $|\varphi(z)| \rightarrow 1$ as $z \rightarrow \partial\mathcal{D}$.*




FIGURE 30. Boundary behavior across line segments.

30.2. A reflection principle for harmonic functions. Let $\mathcal{D}^u$ be a domain in the UHP such that $\partial\mathcal{D}^u \supset I := [a, b] \in \mathbb{R}$. We denote by $\mathcal{D}_I$ the reflection of $\mathcal{D}^u$ across I .

Theorem 30.4. *Assume v is harmonic in $\mathcal{D}^u$ and continuous in $\mathcal{D}^u \cup I$ and $v = 0$ on I . Then v extends to a harmonic function on $\mathcal{D} = \mathcal{D}^u \cup I \cup \mathcal{D}_I$.*

Note 30.5. Note that we cannot simply use the Schwarz reflection principle to prove this theorem, which is stronger. Nothing is known a priori about the continuity of u the harmonic conjugate of v . In fact, the Schwarz reflection principle follows from Theorem 30.4, see Exercise 30.6.

Proof. [10] As in the Schwarz reflection principle, the extension of v is defined through $V(z) = v(z)$ for $z \in \mathcal{D}^u$

$$(30.1) \quad V(\bar{z}) := -v(z) \quad \forall \bar{z} \in \mathcal{D}_I$$

and $V(z) = 0$ on I . The property of a function being harmonic is a local one, so it is enough to check that V is harmonic in any disk in $\mathcal{D} = \mathcal{D}^u \cup I \cup \mathcal{D}_I$ for some family of disks covering $\mathcal{D}$. For disks in $\mathcal{D}^u$ this is so by assumption while for disks $\subset \mathcal{D}_I$ it follows directly from (30.1). Consider now a disk $\mathbb{D}_\varepsilon$ containing part of I as its diameter. Laplace's equation $\Delta v = 0$ with continuous boundary condition $V(z), z \in \partial\mathbb{D}_\varepsilon$ has a unique solution v_1 . We have $V(\bar{z}) = -V(z)$ on $\partial\mathbb{D}_\varepsilon$, and thus $v_2(z) = -v_1(\bar{z})$ is another solution of Laplace's equation with the same boundary condition, hence $v_1 = v_2$. It follows in particular that $v_1 = 0$ on I . But now V and v_1 satisfy Laplace's equation with the same boundary condition on $\overline{\mathbb{D}_\varepsilon \cap \mathcal{D}^u}$ and on $\overline{\mathbb{D}_\varepsilon \cap \mathcal{D}_I}$, hence they are equal in $[\mathbb{D}_\varepsilon \cap \mathcal{D}^u] \cup [\mathbb{D}_\varepsilon \cap \mathcal{D}_I] \cup I$ (the latter because they both vanish on I). ■

Exercise 30.6. Show that if $f = u + iv$ is analytic in a domain $\mathcal{D}$ as in the theorem and v is zero on I , then u is continuous at I . In particular, for the Schwarz reflection principle for analytic functions f in a domain $\mathcal{D}$ in the UHP whose closure contains a real segment I , we only need to show that $\operatorname{Im} f$ is continuous and vanishes on I .

Exercise 30.7. Assume that $\mathcal{D}$ is a simply connected domain in the UHP such that $\overline{\mathcal{D}} \supset I \subset \mathbb{R}$. Let ϕ be the conformal map from $\mathbb{D}$ to $\mathcal{D}$. Consider a half disk strictly contained in $\mathcal{D}$ with diameter on I as in the figure. The arc is the image of an analytic curve in $\mathbb{D}$ bounding a domain $\mathcal{D}'$ in $\mathbb{D}$. Show that $\operatorname{Im} \phi(z) \rightarrow 0$ as $z \rightarrow \partial\mathbb{D} \cap \overline{\mathcal{D}'}$ and apply Theorem 30.4 and Exercise 30.6 to show that ϕ has analytic continuation to the reflection of $\mathcal{D}$ through I . In particular ϕ^{-1} extends continuously to I , and ϕ maps part of $\partial\mathbb{D}$ to I .

Note 30.8. The following ingenious construction is due to Ahlfors [3, p. 27]. If f is analytic, then so is its *reflection*

$$f^*(z) := \overline{f(\bar{z})}.$$

We observe that

$$(30.2) \quad 2u(x, y) = f(x + iy) + \overline{f(x - iy)} = f(z) + f^*(z),$$

for all $x + iy = z$ in the domain where the right-hand side makes sense.

Assuming in addition that the expression $u(z/2, z/2i)$ is defined, we obtain

$$(30.3) \quad 2u\left(\frac{z}{2}, \frac{z}{2i}\right) = f(z) + f^*(0) \\ = f(z) + C \quad \Rightarrow \quad f(z) = 2u\left(\frac{z}{2}, \frac{z}{2i}\right) - C,$$

where $C = f^*(0)$ is a constant.

Consequently,

$$v(z) = \operatorname{Im} \left(2u\left(\frac{z}{2}, \frac{z}{2i}\right) - C \right).$$

Thus, at least in simple cases (rational functions, for instance), one can recover f and v directly from u without integration. Can you extend this construction to general analytic functions?

30.3. Analytic arcs.

Definition 30.9. Let $I = [0, 1]$. A *proper analytic arc* is the image of I under a function γ which extends analytically and injectively to a neighborhood of I in $\mathbb{C}$.

Equivalently, a proper analytic arc is one whose parametrization

$$\gamma : I \longrightarrow \mathbb{C}$$

extends to a biholomorphism between a neighborhood of I and its image.

30.4. An extension of the Schwarz reflection principle. *Informally*, the principle states that if f is analytic in a domain $\mathcal{D}$ which contains an analytic arc γ , if f is continuous up to γ , and if the image of γ is itself an analytic arc, then f extends analytically beyond γ .




FIGURE 31. Analytic continuation across an analytic arc.

Definition 30.10 (One-sided boundary). Let γ be a proper analytic arc bounding a domain $\mathcal{D}$. We say that $\gamma : I \rightarrow \mathbb{C}$, $I = [a, b]$, is a *one-sided boundary* of $\mathcal{D}$ if, for every $z_0 \in \gamma(I)$, there exists a disk $\mathbb{D}_\varepsilon(z_0)$ such that

$$\gamma^{-1}(\mathbb{D}_\varepsilon(z_0) \cap \mathcal{D})$$

lies entirely in either the upper half-plane or the lower half-plane (independently of z_0).¹²

In this case, we also say that $\mathcal{D}$ lies *on one side* of γ . As usual, by analytic continuation of a function f across a curve, we mean the existence of an analytic function $\hat{f}$ in a neighborhood of the curve which coincides with f wherever both are defined. Note that such analytic continuation is necessarily unique.

Definitions The theorem below is the “conformally mapped” Schwarz reflection principle. jjjjj

Exercise 30.11. Let $\mathcal{D}$ be a simply connected nonempty domain with nonempty complement and let ϕ be the conformal map of $\mathbb{D}$ to $\mathcal{D}$ assumed continuous up to the boundary. Show that ϕ has analytic continuation through an arc of $\partial\mathbb{D}$ iff the image of the arc is an analytic arc in $\partial\mathcal{D}$.

Theorem 30.12 (Analytic reflection across arcs). Let $\mathcal{D} \subset \mathbb{C}$ be a simply connected domain, and assume that $\mathcal{D}$ lies on one side of a proper analytic arc γ . Let f be analytic in $\mathcal{D}$ and continuous on $\mathcal{D} \cup \gamma$.

¹²Recall that γ extends to a biholomorphism in a neighborhood of I .

Assume that $f(\gamma) \subset \tilde{\gamma}$, where $\tilde{\gamma}$ is a proper analytic arc, and that $f(\mathcal{D})$ lies on one side of $\tilde{\gamma}$. Then f extends analytically across γ .

Proof. By assumption, γ and $\tilde{\gamma}$ are images of closed intervals I and $\tilde{I}$ under biholomorphisms Γ and $\tilde{\Gamma}$, respectively. The map f is analytic in $\mathcal{D}$ and continuous up to γ , while Γ and $\tilde{\Gamma}$ are biholomorphic in neighborhoods of I and $\tilde{I}$.

Since $f(\Gamma(I)) = \tilde{\Gamma}(\tilde{I})$, we may assume without loss of generality that I is approached from above. Define

$$\psi := \tilde{\Gamma}^{-1} \circ f \circ \Gamma.$$

Then ψ is analytic in a domain in the upper half-plane, continuous on I , and real-valued on I ; hence, the classical Schwarz reflection principle applies to ψ . The desired analytic continuation of f is then given by

$$\hat{f} := \tilde{\Gamma} \circ \psi^* \circ \Gamma^{-1},$$

where ψ^* denotes the analytic continuation of ψ reflected across I . ■

Theorem 30.13. Let v be harmonic in a domain $\mathcal{D}$, and let $\gamma \subset \partial\mathcal{D}$ be a proper analytic arc such that $\mathcal{D}$ lies on one side of γ . Assume that v is continuous up to γ and vanishes on γ . Then v extends to a harmonic function in an open set containing $\mathcal{D} \cup \gamma$.

Proof. Exercise: use Theorem 30.4 and the strategy in Theorem 30.12. ■

Theorem 30.14. Let $\mathcal{D} \subsetneq \mathbb{C}$ be a simply connected domain and φ a conformal map between $\mathcal{D}$ and $\mathbb{D}$. If $\partial\mathcal{D}$ contains a proper analytic arc γ and lies on one side of γ , then

- (i) φ extends analytically across γ .
- (ii) if $z_n \rightarrow z_0 \in \gamma$, then $\varphi(z_n) \rightarrow \varphi(z_0) \in \gamma$.
- (iii) Furthermore, this extension is one-to-one on γ , thus in a neighborhood of $\mathcal{D} \cup (\mathcal{D} \cap \gamma)$.

Note 30.15. By applying a linear fractional transformation, we see that an analogous statement holds if $\mathbb{D}$ is replaced by a half-plane.

In particular, if the boundary of $\mathcal{D}$ is piecewise analytic, then φ extends analytically to a domain $\mathcal{D}' \supset \mathcal{D}$. (It is also biholomorphic in some domain $\mathcal{D}'' \supset \mathcal{D}$.)

Proof. (i) Follows from Theorem 30.12.

(ii) Follows immediately from (i).

(iii) Let $v = \log |\varphi|$. By (ii), we have $|\varphi| \rightarrow 1$ as $\partial\mathcal{D}$ is approached, hence v is well defined close to $\partial\mathcal{D}$ and

$$\lim_{z \rightarrow \partial\mathcal{D}} v(z) = 0.$$

By Theorem 30.4, v extends to a harmonic function on an open set $\mathcal{D}'$ containing $\gamma \cup \mathcal{D}$. Since v is harmonic, there exists an analytic function g such that $v = \operatorname{Re} g$. In $\mathcal{D}$ this g coincides (up to an additive constant) with $\log \varphi$. Choosing the constant to be zero, we have $g = \log \varphi$. Thus $\log \varphi$ has analytic continuation across γ , and in particular it is continuous up to γ . Hence $\varphi = \exp(\log \varphi)$ also extends analytically across γ and maps γ into an arc of a circle.

If φ were not injective on an open subsegment of γ , that portion of γ would be mapped onto a closed curve, violating the continuity of φ (think of the behavior near the endpoints of that segment of γ). ■

30.4.1. Discussion: Tilings with parallelograms. An application of the Schwarz reflection principle to conformal mapping. Consider a conformal map $\varphi : P \rightarrow \text{UHP}$, where P is one of the rectangles in Figure 32. Properly normalized, $\varphi = \operatorname{sn}$, the elliptic sine function (one of the Jacobi elliptic functions; we will return to it later). Theorem 30.12 shows that φ admits analytic continuation through any side of the rectangle to the adjacent one.

By successive reflections, any point in the plane can be reached from one rectangle, and although the same point may be reached by different reflection sequences, the number of reflections is always either even or odd, independently of the chosen path. Hence we obtain a single-valued analytic continuation of φ to the whole of $\mathbb{C}$ except at the corners. This continuation is *meromorphic*, since the point at infinity must be mapped to a point on the boundary of the rectangle.

Because two successive reflections across $\mathbb{R}$ return a point to itself, the extended function is doubly periodic. At the corners, where the sides meet at right angles, the mapping locally straightens the corner, behaving as

$$\varphi(z) = a + (z - z_0)^2(1 + o(1)).$$

Since the function is single-valued and bounded near each corner, it is analytic there. We will prove later that, more precisely, for $k \in [0, 1]$, $\operatorname{sn}(z, k)$ gives a conformal map of the closed rectangle $[-K, K] \times [0, K']$ onto the upper half-plane, with

$$0, \pm K, \pm K + iK', iK' \mapsto 0, \pm 1, \pm k^{-2}, \infty,$$

respectively; iK' is a simple pole.




FIGURE 32. Tiling by reflections of a rectangle.

Source:

<https://upload.wikimedia.org/wikipedia/commons/3/33/Stacked.bond.png>

30.5. Behavior at the boundary, a stronger result.

Definition 30.16. A point $\zeta \in \partial\mathbb{D}$ is called accessible if there is a sequence $\{z_n\}_{n \in \mathbb{N}} \subset \mathcal{D}$ s.t. $\lim_{n \rightarrow \infty} z_n = \zeta$ and a continuous function $\gamma : [0, 1] \rightarrow \mathcal{D}$ which passes through all z_n : $t_0 = 0 < t_1 < \dots < t_n < 1$, $\lim_{n \rightarrow \infty} t_n = 1$, $\gamma(t_n) = z_n$ and $\gamma(1) = \zeta$.

Theorem 30.17 (Boundary behavior). *Let $\mathcal{D}$ be a bounded domain.*

(i) *If z_0 is an accessible point of $\partial\mathcal{D}$ then the biholomorphism φ with the unit disk has a limit (call it $\varphi(z_0)$) as $z \rightarrow z_0$, $z \in \mathcal{D}$ and $\varphi(z_0) \in \partial\mathbb{D}$.*

(ii) *If z_1 and z_2 are accessible points of $\partial\mathcal{D}$ then $\varphi(z_1) \neq \varphi(z_2)$.*

We recall that a Jordan curve in $\mathbb{C}$ is a *continuous map* γ defined (say) on $[0, 1]$ with values in $\mathbb{C}$ which is injective, that is $\gamma(t_1) = \gamma(t_2)$ only if $t_1 = t_2$ or $t_1 = 0$ and $t_2 = 1$ where in the latter case it is a closed Jordan curve. We also recall that a closed Jordan curve divides $\mathbb{C}$ into exactly two regions, one bounded and one unbounded. The bounded region is called the interior of the curve. A Jordan domain is the interior of a Jordan curve. It follows from Theorem 30.17 above that if $\mathcal{D}$ is the interior of a simple, closed Jordan curve, then the map φ extends to a continuous injective function in $\overline{\mathcal{D}}$.

For space limitations we do not prove this interesting result; a proof, essentially based on the first proofs by Lindelöf and Koebe, is found in [10]. We shall not use conformal maps of this generality. In §31 we will find that for polygons, φ can be expressed by quadratures.




FIGURE 33. The origin is inaccessible

30.5.1. *A negative result.* The following is a standard example of an inaccessible point [12] shows that, in full generality, continuity of the conformal mapping cannot be expected. Such is the case of the *inaccessible point* below. Take $\mathcal{D}$ to be an open horizontal rectangle with a vertex at zero from which vertical line segments of length, say, $1/2$ have been removed, see Fig. 33 (check that 0 is inaccessible!). The image of 0 on $\partial\mathbb{D}$ (which wlog can be taken as $z = 1$) must be a point of discontinuity of the conformal map φ into the unit disk, for $\varphi^{-1}(1) = 0$ and in any neighborhood of 1 there are infinitely many points where φ^{-1} is $1/2$.

Illustration of nonanalytic behavior at all boundary points of the conformal image of the unit disk.

Exercise 30.18. Show that:

- The functional equation for the function G ,

$$(30.4) \quad G(z^2) = \lambda^{-1} \frac{G(z)^2}{1 - G(z)}, \quad z \in \mathbb{D}, \quad G'(0) = 1,$$

for $\lambda = 0.5i$, defines a unique function G analytic in the unit disk.

- The points in $G(\partial\mathbb{D})$ are exactly the points $w \in \mathbb{C}$ for which the solution of the one-step recurrence

$$x_{n+1} = \lambda x_n(1 - x_n), \quad x_0 = w,$$

converges to 0.

- $G : \mathbb{D} \rightarrow G(\mathbb{D})$ is a conformal map.

Figure 34 shows the domain $G(\mathbb{D})$. The boundary $J = \partial G(\mathbb{D})$ is called the *Julia set* for the iteration of the quadratic map $x \mapsto \lambda x(1-x)$.

Theorem 30.19. *The unit circle is a natural boundary for G .*

Sketch of proof. By continuity,

$$G(1) = \frac{\lambda}{1 + \lambda}.$$

Differentiating the functional equation (30.4), one finds that if $G'(1) = 0$, then G' must vanish at all binary rational angles. If instead $G'(1) \neq 0$, we can divide by it and obtain

$$G(1) = 1 \pm (1 + 2\lambda)^{-1/2},$$

which contradicts the previous value. Hence $G'(1) = 0$ and, by iterating the argument, G' either vanishes or fails to exist at all binary rational points on the unit circle.

See [7] for details. ■

Exercise 30.20. *Fill in the missing steps in the argument above, and verify that the unit circle is indeed a natural boundary of G .*




FIGURE 34. The conformal image of the unit disk under the map G .

31. CONFORMAL MAPPINGS OF POLYGONS AND THE SCHWARZ-CHRISTOFFEL FORMULAS

For polygonal regions, the conformal map to the unit circle (or to $\mathbb{H}^u$ obviously) can be done by quadratures. The transformation is still usually nonelementary, but the integral representation gives us enough control to describe the transformation quite well.




FIGURE 35. Schwarz-Christoffel transformations: a closed polygon.

31.1. Heuristics. If ψ is biholomorphic at z_0 , the angle between the tangent of a curve γ through z_0 and the tangent to its image through ψ is $\arg \psi'(z_0)$; we write this in differential form, $dw = \psi'(z)dz$. We want to map $\mathbb{H}^u$ to the interior of a polygon. We then choose the positive orientation when traversing $\partial\mathbb{H}^u$ (which leaves the domain to its left): this means traversing the boundary from $\mathbb{R}^-$ to $\mathbb{R}^+$. For later convenience we denote by $\pi\alpha_i$ the interior angles of the polygon.

We place without loss of generality a vertex at zero, and rotate the polygon so that one side is in $\mathbb{R}^+$. The red arrow indicates the positive orientation of the polygon. Suppose that we want to map 0 to 0 and the segment in blue on $\partial\mathbb{H}^u$ to the blue segment on the polygon, see Fig. 36. We see that, to the left of $z = 0$, dw is rotated by $-\pi(1 - \alpha)$ with respect to dz , while to the right of $z = 0$ (red arrow) dz and dw are parallel. A transformation that behaves like this on the boundary is $\psi'(z) = z^{-(1-\alpha)}$. We see that indeed the argument of ψ' ($\text{Im} \ln \psi'$ which exists locally for $z \neq 0$ since $\psi' \neq 0$) does not change except at the singularity, $z = 0$:

$$(31.1) \quad (\ln \psi')' = \frac{\alpha - 1}{z} \in \mathbb{R} \quad (\text{since } z \in \mathbb{R}) \Rightarrow d \arg \psi' = 0 \text{ for } z \neq 0$$

Also, note that by Schwarz reflection, ψ is analytic at points z which are *not* mapped to vertices of the polygon.

Proposition 31.1. *Any transformation ψ of a one-sided neighborhood $\mathcal{N}$ in $\mathbb{H}^u$ of a segment $I = [-a, b]$; $a, b > 0$ which maps 0 to the vertex $w = 0$ of the polygon and is continuous up to the boundary has the property $\psi(z) = z^{\alpha-1}h(z)$ where h is holomorphic in $\mathcal{N} \cup I \cup \bar{\mathcal{N}}$. In particular, $\psi'(z) = z^{\alpha-1} \sum_{k=0}^{\infty} h_k z^k$ where $h(0) \neq 0$ and the sum is convergent.*




FIGURE 36. Schwarz-Christoffel transformations: two adjacent sides of the polygon.

Definition 31.2. We will call functions which are (locally) of the form $(z - z_0)^\alpha H(z)$ with H holomorphic, **ramified-analytic**.

Proof. Let $H = \int z^{-(\alpha-1)} \psi'$. Note that $dH = z^{-(\alpha-1)} \psi' dz$ maps the blue vector on the left side of the polygon $\mathcal{N}$ into a vector parallel to the red one, is continuous down to I . The image of I through $H = \int z^{-(\alpha-1)} \psi'$ will, by construction, be an interval in $\mathbb{R}$. Schwarz's reflection principle and the continuity of H ensure that H has analytic continuation in $\mathcal{N} \cup I \cup \overline{\mathcal{N}}$. Thus H' is analytic too. Since the transformation h is one-to one ($H(z)$ is strictly monotonic in $z \in I$, implying $H' = h \neq 0$). ■

More generally, a transformation of the form

$$(31.2) \quad \psi' = \prod_{i=1}^{n-1} (z - a_i)^{\alpha_i - 1}$$

satisfies

$$(31.3) \quad (\ln \psi')' = \sum_{i=1}^{n-1} \frac{\alpha_i - 1}{z - a_i} \in \mathbb{R} \Rightarrow d \arg \psi' = 0 \text{ for } z \neq a_i$$

and $\arg \psi'$ changes by $\pi(1 - \alpha_i)$ upon traversing a_i . The points $A_i := \psi(a_i)$ on the polygon are the only ones where $\arg \psi'$ changes, and it changes by $+\pi(1 - \alpha_i)$ (check the signs!). If the polygon P is a closed

curve, then infinity must be mapped into one of the vertices of P , and, since ψ is conformal (to be proven later) it cannot be one of the $\{A_j, j = 1..n-1\}$ so it must be A_n . Recall that the sum of exterior angles of a closed polygon is 2π , and thus $B = \sum_1^{n-1} \beta_i = 1 + \alpha_n$, $\beta_i := a_i - 1$. Note also that $1 - a_i/z$ is analytic at infinity: as $z \rightarrow \infty$ we have

(31.4)

$$\psi' = z^B \prod_{i=1}^{n-1} (1 - a_i/z)^{\beta_i} = \zeta^{-B} g(\zeta); \quad g(\zeta) = \prod_{i=1}^{n-1} (1 - a_i \zeta)^{\beta_i}, \quad \zeta = 1/z$$

If $|\zeta| < \varepsilon < \max\{1/|a_i|, i = 1, \dots, n-1\}$, g is analytic and $\operatorname{Re} g(\zeta) > 0$, implying that $\ln g$ is well defined. If z traverses $\partial\mathbb{D}_R$ from $+R$ to $-R$ where $R = 1/\varepsilon$ in a positive direction the change in $\arg \psi'$ comes solely from z^B and it equals πB , the same change in argument that $(z - A)^{1-\alpha_n}$ would produce. This “closes the polygon” with the last vertex $A_n = \lim_{z \rightarrow \infty} f(z)$. See figure. In the case of a closed polygon we see that ψ is bounded on $\mathbb{R}$. In the case of an open polygonal line, the arguments are similar.

Theorem 31.3. (i) Let P be any polygon (closed or open) with n vertices $A_1, \dots, A_n$ and interior angles $\pi\alpha_1, \dots, \pi\alpha_n$. There is a choice of $a_1, \dots, a_n \in \mathbb{R}$ and $C, C' \in \mathbb{C}$ such that the function

$$(31.5) \quad \psi = z \mapsto C \int_0^z \prod_{k=1}^n (s - a_k)^{\alpha_k - 1} ds + C'$$

maps $\mathbb{H}^u$ into P (and $\psi(a_i) = A_i$).

(ii) The function mapping $\mathbb{D}$ conformally into P is given by

$$(31.6) \quad \psi = z \mapsto C \int_0^z \prod_{k=1}^n (b_k - s)^{\alpha_k - 1} ds + C'$$

for a choice of $b_1, \dots, b_n, C, C'$. (iii) Moreover, all transformations between $\mathbb{H}^u$ or $\mathbb{D}$ and polygons are of this form.

(iv) One of the points a_k can be chosen to be ∞ . In this case the transformation is

$$(31.7) \quad \psi = z \mapsto C \int_0^z \prod_{k=1}^{n-1} (s - a'_k)^{\alpha_k - 1} ds + C'$$

(simply the point a_n is omitted). (v) The map from $\mathbb{D}$ to $\text{ext}(P)$ taking 0 to ∞ is

$$(31.8) \quad \psi = z \mapsto C \int_0^z s^{-2} \prod_{k=1}^{n-1} (s - a'_k)^{\alpha_k - 1} ds$$

Note 31.4. (i) The formulas (31.5) and (31.6) are the same, up to constants: (31.5) is invariant under a Cayley transform since the sum of exterior angles of a closed polygon is 2π . Do the calculation.

(ii) One may ask what kind a conformal map is that in (v)? The answer is best seen if we think of the Riemann sphere as the target.

Remark 31.5. (i) It is important to note what freedom we have in such transformations. Suppose we want to map a triangle Δ . All triangles with same angles are similar, and a mapping between two similar triangles reduces to scaling, rotation and translation. Thus we need to understand one triangle with given angles α_i . We take say $a_1 = 0$ and $a_2 = 1$, use the α 's, and see what triangle Δ_1 is obtained. Then, we can choose C and C' so that we remap Δ_1 to Δ . Thus we are able to map any triangle to $\mathbb{H}^u$, prescribing the position of the images of the vertices at will. See §31.2.

(ii) For $n > 3$, we can still place three points at will but the position of the fourth one etc cannot be chosen arbitrarily. We see that we have the freedom of $n - 1$ real constants, the a'_i 's, and of two complex ones C and C' , a total of $n + 3$ real constants, whereas an arbitrary closed polygon has $2n$ real constants as degrees of freedom (the position of its vertices); $2n > n + 3$ if $n > 3$. The a_k for $k \geq 4$ are called accessory parameters, and they are determined by the polygon and the values of $a_1, \dots, a_3$; except for very symmetric cases (such as regular polygons), these parameters cannot be determined in closed form.

Given the disparity in the number of available parameters versus the degrees of freedom of the problem, it is not a priori clear that the Schwarz-Christoffel transformation should always work (but it does).

First proof of the Schwarz-Christoffel formula. We analyze closed polygons, open ones being similar. Once we prove the formula for the UHP, the formula for $\mathbb{D}$ follows through a simple calculation applying the Cayley transform.

We can first arrange that one vertex is at zero and a second one at 1. Indeed, we can transform P by translation and multiplications by a constant (changing C , C' in the Theorem) into one geometrically similar to it, $\tilde{P}$ that has these properties. By composition to the right

with $az+b$ we can arrange that 0 and 1 in $\mathbb{H}^u$ are mapped into 0 and 1 in $\tilde{P}$. Note that by Proposition 31.1 the function $F(z) = \psi'(z) \prod_{i=1}^{n-1} (z - a_i)^{1-\alpha_i}$ is analytic in $\overline{\mathbb{H}^u}$, does not change its phase except perhaps at a_i . But at every a_i , F is real and positive. Thus F is real on the real line and extends to an entire function. For the behavior at ∞ note that the transformation $\zeta = -1/z$, $\psi(z) = G(\zeta)$. $z \mapsto -1/z$ is an automorphism of $\mathbb{H}^u$ which maps ∞ to 0. Traversing $\mathbb{R}^+$ in z from 0 to ∞ , returning to $-\infty$ on a “large” circle and moving in $(-\infty, 0)$ to the right corresponds to ζ traversing $(-\infty, \infty)$ from negative to positive values. To straighten the angle at a_n ($= 0$ in ζ) we need, as in Proposition 31.1 to multiply $G'(\zeta)$ by $\zeta^{1-\alpha_n}$: $\zeta^{1-\alpha_n} \frac{dG}{d\zeta} = H(1/z)$ where H is analytic in particular bounded. Since $\frac{d}{d\zeta} G = \zeta^{-2} \psi'(-1/\zeta)$ it follows that $z^{1+\alpha_n} \psi'$ is analytic at infinity.

Now, $1 - \alpha_1 + \dots + 1 - \alpha_n = 2$ implying $1 - \alpha_1 + \dots + 1 - \alpha_{n-1} = 1 + \alpha_n$ and thus

$$(31.9) \quad (z - a_1)^{1-\alpha_1} \dots (z - a_{n-1})^{1-\alpha_{n-1}} \psi' \sim \text{const.} z^{1+\alpha_n} \psi' \text{ as } z \rightarrow \infty$$

is entire and bounded, thus constant and hence

$$(31.10) \quad (z - a_1)^{1-\alpha_1} \dots (z - a_{n-1})^{1-\alpha_{n-1}} \psi' = C$$

Corollary 31.6 (Analytic structure of ψ). *The function ψ is ramified-analytic (cf. Definition 31.2). In a neighborhood of a_j we have*

$$(31.11) \quad \psi = C_{1j} x^{\alpha_j} H_j + C_{2j}$$

where H_j is holomorphic.

Proof. This follows by straightforward integration of (31.10). ■

■

“Geometric” proof of the Schwarz-Christoffel formula. This proof is largely based on [12]. A slightly different argument is used in [3]. We prove the statement for a closed polygon, the one for open ones being very similar. We use the notations in Fig. 37 and will assume first that none of the points a_i is infinity (infinity must however go somewhere on ∂P).

Let ψ be the conformal map between the UHP and P . The figure shows various successive reflections of a polygon P across its sides. Since ψ transforms an interval I of $\mathbb{R}^+$ into the line segment ℓ bounding P and P_1 , it has analytic continuation across I , and in fact the image of the UHP is P and that of the LHP is P_1 . This continuation can be reflected back, and we obtain P_2 . By simple geometry, the polygons P and P_2 are Euclidian transformations of each-other, of the form




FIGURE 37. Successive reflections.

$P_2 = aP + c$ for some constants a and c with $|a| = 1$, whereas P and P_1 are related by a flip. We note also that ψ is analytic at all points except for $a_1, \dots, a_n$, where it is singular. Indeed, at a_i ψ is not conformal: it maps two collinear successive intervals of $\mathbb{R}$ into two segments forming an angle $\pi\alpha_i$. However, consider the auxiliary function

$$h(z) = [\psi(z) - \psi(a_i)]^{1/\alpha_i}$$

Then, h maps the two successive intervals into a straight line (clearly an analytic arc), and by Schwarz reflection, it is analytic at a_i . Hence,

$$(31.12) \quad \psi(z) = \psi(a_i) + [h(z)]^{\alpha_i}$$

Since $h(a_i) = 0$, a_i is a branch point of ψ . Consider the function

$$g(z) = \psi''(z)/\psi'(z)$$

Using (31.12) we see that g has a simple pole of residue $1 - \alpha_i$ at a_i . Since a_i are the only singularities of g , g is single valued and extends to a meromorphic function in $\mathbb{C}$. Thus,

$$(31.13) \quad G(z) = g(z) + \sum_{j=0}^{n-1} \frac{1 - \alpha_j}{z - a_j}$$

is entire. Since $\psi(\infty) \in \partial P$ but is different from $a_1, \dots, a_n$, by the Schwarz reflection principle, ψ is analytic at infinity (what that means is best understood if we map the UHP conformally to $\mathbb{D}$: ∞ corresponds to some point on $\partial\mathbb{D}$, other than the images of the a_i). Hence, in a

neighborhood of $z = \infty$ we have $\psi(z) = \sum_{k=0}^{\infty} c_k z^{-k}$, where $c_0 = \psi(\infty)$ whence $g(z) \sim \text{const.}/z$ for large z and in particular it vanishes at ∞ . Then G is an entire function that vanishes at infinity implying $G = 0$ and we have

$$(31.14) \quad g(z) = - \sum_{j=0}^{n-1} \frac{1 - \alpha_j}{z - a_j}$$

meaning

$$(31.15) \quad \frac{\psi''(z)}{\psi'(z)} = \sum_{j=0}^{n-1} \frac{\alpha_j - 1}{z - a_j}$$

Integrating 31.15, we get

$$\psi(z) = c_1 + c \int_0^z \prod_{k=1}^n (t - a_k)^{\alpha_k - 1} dt$$

and Theorem 31.3 (i) and (iii) follow, and, as discussed, a composition with the Cayley transform proves (ii). To make one of the $a_k = \infty$ we use an LFT substitution: $z = a_k - 1/z_1$, and the rest of the proof of (iv) is a simple calculation.

Finally, for the mapping of the exterior of the polygon, which contains the point at infinity, making g unbounded there, the formula has to be reworked. Assume that $\infty = \psi(0)$. Thus ψ is unbounded, but must be conformal at zero, and the only possibility is a first order pole: $\psi(z) = \psi_1(z)/z$ where ψ_1 is regular at 0 and $\psi_1(0) \neq 0$. Now $\psi''(z)/\psi'(z) + 2/z$ is regular and a calculation shows that

$$\frac{\psi''(z)}{\psi'(z)} - \sum_{i=1}^n \left(\frac{1 - \alpha_i}{z - a_i} + \frac{1 - \alpha_i}{a_i} \right) + \frac{2}{z}$$

is entire, bounded in $\mathbb{C}$ and zero at zero, hence it vanishes identically, and (v) follows. ■

Note 31.7. The second order linear ODE obtained after multiplying (31.15) by f' is a *Fuchsian equation*: it has only *regular singularities* in $\hat{\mathbb{C}}$ [4].

31.2. Conformal map of the UHP onto the interior of a triangle of angles $\pi\alpha, \pi\beta, \pi\gamma$. We choose for convenience $a_1 = 0, a_2 = 1, a_3 = \infty$ resp. Theorem 31.3 (iv) gives

$$(31.16) \quad \psi(z) = A \int_0^z z^{\alpha-1} (1-z)^{\beta-1} dz + B$$

We choose $A = 1$. From the choice above, we know that the length of the first side is

$$\int_0^1 |\psi'(s)| ds = \int_0^1 s^{\alpha-1} (1-s)^{\beta-1} ds = \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)}$$

The lengths of the other sides can be most easily determined from the relation

$$\frac{a}{\sin \pi \alpha} = \frac{b}{\sin \pi \beta} = \frac{c}{\sin \pi \gamma}$$

We note that $\psi(z) = B_z(\alpha, \beta)$, the incomplete Beta function, which, for $\alpha, \beta \notin -\mathbb{N}$ satisfies

$$B_z(\alpha, \beta) = \Gamma(\alpha) z^\alpha {}_2F_1(\alpha, 1-\beta; \alpha+1; z)$$

where ${}_2F_1$ is the hypergeometric function.

31.3. Schwarz triangle functions and hypergeometric functions.

We can attempt the reflection-continuation procedure of §34.1. We now imagine the reflections having a common vertex. To insure a single valued function upon successive reflections about the sides, we must return to the starting triangle with no overlap or gap. If the rotations preserve one vertex, for this to happen we need $\alpha_i = 1/n_i$, $n_i \in \mathbb{N}$. The constraints are thus:

$$(31.17) \quad \alpha + \beta + \gamma = 1; \quad 1/\alpha, 1/\beta, 1/\gamma \text{ positive integers}$$

Check that the only solutions are: $(\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$ (equilateral triangle) $(\frac{1}{2}, \frac{1}{3}, \frac{1}{6})$ (half of an equilateral triangle) and $(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ (isosceles right triangle). Then the reflected images cover the whole plane and the mapping functions are restrictions of meromorphic functions. These are special cases of the *Schwarz triangle functions*. The group of reflections is a special case of an infinite *Coxeter group*.

Each triangle function corresponds to an elliptic function. We will return to this topic later.

Differential equation. We can derive an equation with polynomial coefficients for ψ as follows. In $\psi' = z^{\alpha-1}(z-1)^{\beta-1}$ we write $\psi = z^\alpha \psi_1$ and get, after dividing by $z^{\alpha-1}$,

$$(31.18) \quad z\psi_1' + \alpha\psi_1 - (1-z)^{\beta-1} = 0 \Rightarrow \frac{z}{(1-z)^{\beta-1}}\psi_1' + \frac{\alpha}{(1-z)^{\beta-1}}\psi_1 - 1 = 0$$

which we differentiate one more time to eliminate the constant 1 and we get

$$(31.19) \quad z(1-z)\psi_1'' + [\alpha+1-(2-\beta-\alpha)z]\psi_1' + \alpha(\beta-1)\psi_1 = 0$$

The differential equation for the hypergeometric function ${}_2F_1(a, b; c; z)$ is the *Riemann equation*

$$(31.20) \quad z(1-z)h'' + [c - (a+b+1)z]h' - abh = 0$$

From the way we went from (31.18) to (31.19) (or by direct verification) we see that one solution is $z^{-\alpha}$. Comparing with (31.20) the second one is

$$(31.21) \quad {}_2F_1(\alpha, 1-\beta; \alpha+1; z) \Rightarrow \psi = z^\alpha \cdot {}_2F_1(\alpha, 1-\beta, \alpha+1, z)$$

That is, in this case, the Schwarz-Christoffel transformation is a ratio of two independent solutions of the special hypergeometric equation (31.19).

32. OPTIONAL MATERIAL: CURVILINEAR TRIANGLES

In general, the map from $\mathbb{H}^u$ into a *curvilinear triangle*, one whose sides are arccircles is given by the ratio of two independent solutions of (31.20), where the angles α, β, γ are related to a, b, c by (cf. [12])

$$(32.22) \quad a = \frac{1}{2}(1 + \beta - \alpha - \gamma), \quad b = \frac{1}{2}(1 - \alpha - \beta - \gamma), \quad c = 1 - \alpha$$

Note 32.8. To see qualitatively why that is, before we work out the mathematical details, since the sought-for function f transforms segments into arccircles, Möbius transforms of f map real segments into real segments. Möbius transforms are conformal wherever defined, so they preserve angles, and thus f should be ramified analytic at three points, say $\{0, 1, \infty\}$.

Secondly, note that any function which is real-analytic (with real values) on some interval I and such that $f' \neq 0$ on I conformally maps a neighborhood of I into a neighborhood of $f(I)$ (where, of course, $f(I)$ is an interval in $\mathbb{R}$).

Finally, this approach would apply to any curvilinear polygon, and the final ODE that we construct will still be second order linear, but the solutions are generally quite complicated.

To find an f as in Note 32.8, a natural candidate would be a ratio of two solutions of a real-valued linear second order ODE with analytic coefficients, whose solutions have only ramified singularities, at in $\{0, 1, \infty\}$. Such an ODE will have two linearly independent solutions, f_1 and f_2 ¹³. Define $S = \{a_1 f_1 + a_2 f_2 \mid A_{1,2} \in \mathbb{C}\}$ (this of course is the space of all solutions). Then,

- (1) $f_{1,2}$ are analytic in $\mathbb{C}^* \setminus \{0, 1, \infty\}$ while at the points $0, 1, \infty$ they are *ramified-analytic*;

¹³Meaning that $A, B \in \mathbb{C}$ and $Af_1 + Bf_2 = 0 \Rightarrow A = B = 0$.

- (2) on each interval $(-\infty, 0)$, $(0, 1)$ and $(1, \infty)$ there are two positive linearly independent functions in S ;
- (3) S is invariant under Schwarz reflections¹⁴. This is because any analytic continuation of a solution is a solution, by permanence of relations.
- (4) $(f_1/f_2)' \neq 0 \forall x \in \mathbb{R}$.¹⁵

Then some ratio F_1/F_2 of two linearly independent $F_{1,2} \in S$ maps the upper half plane into a curvilinear triangle, that is, one whose sides are line segments or arccircles (a line segment is a limiting case of an arccircle, so we will call arccircle too). Indeed, take F_1 and F_2 in S linearly independent, choose one of the intervals in (2) above, and let $\tilde{F}_1$ and $\tilde{F}_2$ the functions which are real valued on the interval I . We choose $\tilde{F}_1$ and $\tilde{F}_2$ such that $\tilde{F}_1/\tilde{F}_2$ is bounded at zero. By assumption, $F_{1,2} = A_{1,2}\tilde{F}_1 + B_{1,2}\tilde{F}_2$ and

$$(32.23) \quad \frac{F_1}{F_2} = \frac{1 + a\tilde{F}_1/\tilde{F}_2}{b + c\tilde{F}_1/\tilde{F}_2}$$

Now, $\tilde{F}_1/\tilde{F}_2$ is real-valued and one-to-one, and thus the right side of (32.23) is a Möbius transformation of a line segment: an arccircle. The image of $\mathbb{R}$ through F_1/F_2 consists of three arccircles, a general curvilinear triangle, provided that the singularities are such that F_1/F_2 are continuous on $\mathbb{R}$.

We now show that (31.20) is such an equation, for a, b, c satisfying (32.22). The following integral representation due to Euler can be checked to solve (31.20):

$$(32.24) \quad {}_2F_1(a, b; c; z) = K \int_0^1 H(z, x) dx,$$

where $H(z, x) = x^{b-1}(1-x)^{c-b-1}(1-zx)^{-a}$ assuming $c > b > 0$

The standard choice of K , immaterial here, is $K^{-1} = B(b, c-b)$ with B the Beta function¹⁶. For the integral to exist, we need

$$(32.25) \quad b-1 > -1, c-b-1 > -1, -a > -1$$

¹⁴Equivalently, the space generated by $f_{1,2}$, $S := \{C_1 f_1 + C_2 f_2 : C_1, C_2 \in \mathbb{C}\}$ is closed under continuations at the branch points (e.g., $f_1(ze^{2\pi i}) = C_1 f_1(z) + C_2(z)$).

¹⁵This in fact is implied by linear independence. Check this by first showing that two functions are linearly independent iff their Wronskian $W(f_1, f_2) := f_1' f_2 - f_2' f_1 \neq 0$.

¹⁶The formula is valid under the more general condition $\operatorname{Re}(c) > \operatorname{Re}(b) > 0$, but here we only need real a, b, c .




FIGURE 38. Hypergeometric contours

which are satisfied if $\alpha + \beta + \gamma < 1$, $\alpha, \beta, \gamma \in \mathbb{R}^+$, which is the case for a hyperbolic triangle (with concave sides) as can be verified from (32.22). Under this same condition, we have

$$(32.26) \quad (b-1) + c - b - 1 - a < -1 \Rightarrow H(x, z) \sim s^{-p}, \quad p > 1 \text{ as } x \rightarrow \infty$$

Singularities of ${}_2F_1$ For general a, b, c the behavior of h in (31.20) at the singular points of the ODE, $0, 1, \infty$ ¹⁷ follows general results about regular singular points of ODEs (Frobenius theory, cf. [4]); we will not assume this here however. We will find the behavior of the solutions in three different way, to illustrate various approaches.

Directly from the integral representation, the behavior can be calculated in the following way.

Clearly, (32.24) and Corollary ?? imply that ${}_2F_1$ is analytic except possibly on $\mathbb{R}^+$. If $z \rightarrow a \in \mathbb{R}^+ \setminus \{1\}$ then we can use the analyticity of the integrand at a to first homotopically deform the contour as shown in Fig. 38. The new integral (of course, equal to the original one) is manifestly analytic in z near a . Thus the only possible singularity is at 1 (since the integrand is manifestly analytic for small z , cf. Corollary ??).

¹⁷Writing the equation in the form $h'' + Q(x)h + R(x) = 0$, $\{0, 1\}$ are singular points of P, Q . Ditto after the change of variable $z = 1/t$, $y(z) = t^{-c}Y(t)$ at $t = 0$.

At $z = 0$ we get

$$(32.27) \quad \frac{{}_2F_1(a, b; c; z)}{K} = \int_0^1 x^{b-1}(1-x)^{c-b-1} dx > 0$$

To find the type of behavior at $z = 1$ it is convenient to take $z = 1 - \varepsilon$ and change the variable to $x = 1 - s$ in the integral:

$$(32.28) \quad K \int_0^1 h(z, (1-s)) ds = K \int_0^1 (1-s)^{b-1} s^{c-b-1} (\varepsilon + s - \varepsilon s)^{-a} ds$$

By (32.26) We can then push the contour up toward $+i\infty$,

$$(32.29) \quad \int_0^1 h(z, (1-s)) ds = \left(\int_0^{+i\infty} - \int_1^{+i\infty} \right) (1-s)^{b-1} s^{c-b-1} (\varepsilon + s - \varepsilon s)^{-a} ds$$

The second integral is analytic in ε by the same Corollary ??.

In the first one, we change variable to $s = \varepsilon u$, to get

$$(32.30) \quad -\varepsilon^{c-b-1-a+1} \int_0^{+i\infty} (1-\varepsilon u)^{b-1} u^{c-b-1} (1+u-\varepsilon u)^{-a} ds \\ = -\varepsilon^{c-b-a} \int_0^{-\infty} (1-\varepsilon u)^{b-1} u^{c-b-1} (1+u-\varepsilon u)^{-a} ds$$

where the contour change is justified by (32.26).

Exercise 32.9. *What is the phase of the last integral in (32.30)?*

The integral in (32.30) is analytic in ε and thus

(32.31) ${}_2F_1(a, b; c; z) = A_1 + A_2(1-z)^\gamma$ where $A_{1,2}$ are analytic at 1 and $A_2(1) = \int_1^{+\infty} u^{c-b-1}(1+u)^{-a} ds > 0$. A second solution of (31.20) is obtained, cf. [12], by noticing that the substitution $h(z) = g(1-z)$ leads to the equation (31.20) with c replaced by $C = a + b + 1 - c$. The integral representation (32.24) still holds with c replaced by $C = 1 - \gamma$, and converges under the same conditions on α, β, γ as (32.24). The behavior of the integrand at infinity is still of the form in (32.26). Thus the same analysis applies, to show that there is a second solution which is analytic at $z = 1$ and has a singularity at zero, with $c - b - a$ replaced by $1 - c = \alpha$. This also suggests making the substitution $h_1 = z^{1-c}h = z^\alpha h$ directly into (31.20); we get an equation of the form (31.20), with A, B, C replaced by

$$(32.32) \quad A = a + 1 - c, B = b + 1 - c, C = c - 2$$

which has an integral representation of the form (32.24) valid under the same conditions on α, β, γ . Thus, near $z = 0$ we have two linearly independent solutions, A_1 analytic and A_2 of the form $z^\alpha A(z)$ with A

analytic and $A(0) \neq 0$. These are clearly linearly independent since the second solution is not analytic at zero. In the same way, there are two solutions, one analytic and positive at $z = 1$ and another one of the form $z^\gamma A(z)$, A analytic and positive for $z < 1$.

From the ODE. Here we are assuming knowledge of basic properties of linear ODEs: a linear combination (with constant coefficients) of solutions is a solution (this can be checked directly) and the fact that the space of solutions of a second order linear equation is two-dimensional (i.e., there are exactly two free constants, or, in other words, there are two linearly independent solutions which form a basis in the space of all solutions).

With F_{01} given by (32.24), we notice as before that F is analytic near zero and $F(0) > 0$. We look for a second solution in the form $F_{02} = F_{01}g$. The equation for g is

$$(32.33) \quad \begin{aligned} \frac{g''}{g'} &= -2\frac{F'}{F} + q(x); \quad \left(q(x) := \frac{(a+b+c)x-c}{x(1-x)} \right) = -2\frac{F'}{F} - \frac{c}{x} + A(x) \\ &\Rightarrow g' = -\frac{c}{x}A_1(x) \Rightarrow F_{01}(x) = x^{-c+1}A_2(x) = x^\alpha A_2(x); \end{aligned}$$

with A, A_1, A_2 analytic (check the conclusions above). Since F is real-valued for real $z \in (0, 1)$, g is also real-valued, and it is an independent solution (it has a manifestly different behavior at zero). At $x = 1$ we make the substitution $z = 1 - y$ and we get

$$(32.34) \quad y(1-y)h'' + [C - (a+b+1)y]h' - abh = 0, \quad C = a+b+c-1 = 1-\gamma$$

In the same way as above we get two independent real valued solutions for y real, $F_{10}(y)$ analytic and a second one F_{11} of the form $y^\gamma A_1$ with A_1 analytic. They are in general different from the solutions $F_{00}(y)$ and $F_{01}(y)$.

Finally the substitution $h(z) = z^a H(1/z)$ $z = 1/Z$ in (31.20) results in an equation of the same type as (31.20). The ratio of two solutions behaves like $z^{-\beta}$ for large z . Now, any transformation of the type F_1/F_2 where F_1 and F_2 are real maps the interval $(0, 1)$ into an interval. A different choice f_1/f_2 is a Möbius transformation of F_1/F_2 , and thus it maps the interval $(0, 1)$ into an arccircle. Keeping one combination F_1/F_2 chosen so that $F_1/F_2 \rightarrow 0$ as $z \rightarrow \infty$ will then map $\mathbb{R}$ into a curvilinear triangle, of angles α, β, γ .

Note 32.10. In the simplest nondegenerate case, Frobenius theory allows to determine the behavior of solutions of linear meromorphic ODEs, Lf , with regular singularities, say at z_0 by a very simple method:

take $L(z - z_0)^r$ and keep only the lowest power of $(z - z_0)$. This gives a quadratic equation for r . If the solutions $r_{1,2}$ do not differ by an integer, then $Lf = 0$ has two linearly independent solutions in the form $(z - z_0)_{1,2}^r A_{1,2}(z)$, A analytic at z_0 .

Note 32.11. The equation satisfied by f_1/f_2 , a ratio of solutions of the hypergeometric equation is

$$(32.35) \quad \{w, z\} = \frac{1 - \alpha^2}{2z^2} + \frac{1 - \gamma^2}{2(z - 1)^2} + \frac{\alpha^2 + \gamma^2 - \beta^2 - 1}{2z(z - 1)} = 0$$


where $\{w, z\}$ is the important *Schwarzian derivative* [12]:

$$(32.36) \quad \{w, z\} = \left(\frac{w''}{w'} \right)' - \frac{1}{2} \left(\frac{w''}{w'} \right)^2 \quad \left(w' = \frac{dw}{dz} \right)$$

The Schwarzian derivative is *invariant under any Möbius transformations of w* as expected from our discussion and can be checked by straightforward calculation.

33. TWO OTHER IMPORTANT EXAMPLES OF SCHWARZ-CHRISTOFFEL TRANSFORMATIONS

33.1. Another look at the sine function. *Problem.* Map the strip indicated into $\mathbb{H}^u$ preserving the points marked with circles and the positive orientation.



Solution The α 's at $-\pi$ and π are both $1/2$. We note that Theorem 31.3 still applies, by passing the angles to this limit. We apply formula (31.5) with $a_1 = -1$, $a_2 = 1$ and the integrand is then $(s^2 - 1)^{-1/2}$. Eq.

(31.5) therefore gives, for two arbitrary constants,

$$(33.37) \quad \psi = C \arcsin z + C'$$

and therefore our map $f = \psi^{-1}$ has the general form

$$(33.38) \quad \psi^{-1}(w) = \sin(cw + c')$$

We have now to choose c and c' to match the prescribed points. We must have $\sin(-\pi c + c') = -1$ and $\sin(c\pi + c') = -1$; the choice $c' = 0$ and $c = 1/2$ matches these conditions. We get

$$(33.39) \quad f(w) = \sin(w/2)$$

34. MAPPING OF A RECTANGLE: ELLIPTIC FUNCTIONS

Placing one of the a_i at infinity, and choosing the remaining three to be $0, 1, \rho$ where $\rho > 0$, we get the Schwarz-Christoffel formula for a rectangle in the form

$$(34.1) \quad F(z) = \int_0^z \frac{ds}{\sqrt{s(s-1)(s-\rho)}}$$

The double symmetry of the rectangle suggests a symmetric choice of a_i :

$$(34.2) \quad \psi(z, k) = \int_0^z \frac{ds}{\sqrt{(1-s^2)(1-k^2s^2)}}$$

Here $k \in (0, 1)$. We have $\psi(z, k) = F(\arcsin z, k)$ where F is the incomplete elliptic integral of the first kind.

The branch of the square root is taken to be positive for $z \in (0, 1)$ and otherwise it is obtained by analytic continuation in the UHP, passing to the limit $\text{Im } z \rightarrow 0^+$ for $z \in \mathbb{R}$. Thus, as z grazes $\mathbb{R}^+$, the square root is first in $\mathbb{R}^+$, then changes to $i\mathbb{R}^+$ at 1 and then to $\mathbb{R}^-$ after $1/k$. Also, since the square root is even for small z , by analyticity this property is preserved along $\mathbb{R}$, and, for $k \in \mathbb{R}$, $\psi(z, k) = -\psi(-z, k)$. $\psi(\mathbb{H}^u)$ it is a rectangle with vertices

$$(34.3) \quad (-K, 0), (K, 0), (K, iK') \text{ and } (-K, iK)$$

where

$$(34.4) \quad K = \int_0^1 \frac{ds}{\sqrt{(1-s^2)(1-k^2s^2)}}$$

$$(34.5) \quad iK' = \int_1^{1/k} \frac{ds}{\sqrt{(1-s^2)(1-k^2s^2)}}$$




FIGURE 39. The fundamental rectangle.

Traversing $\mathbb{R}$ gives

$$(34.6) \quad \int_{-\infty}^{\infty} \frac{ds}{\sqrt{(1-s^2)(1-k^2s^2)}} = 0$$

(by symmetry, or since the contour can be pushed up to $i\infty$). Observe that for $s > 1/k$ we note that the square root is real and negative and we have

$$(34.7) \quad \sqrt{(1-s^2)(1-k^2s^2)} = -ks^2 \sqrt{(1-s^{-2})(1-(ks)^{-2})}$$

(where the square roots on the right side are positive) and that the function $\sqrt{1-\zeta}$ $\zeta = 1/s^2$ is *analytic* for $|\zeta| < 1$. Thus $\psi(z, k)$ is analytic at infinity and for $z > 1/k$, writing $\int_0^z = \int_0^\infty - \int_z^\infty$ we get the expansion at infinity

$$(34.8) \quad \begin{aligned} \psi(k, z) &= iK' - \int_z^\infty -\frac{1}{ks^2} - \left(\frac{k^{-1} + k^{-3}}{2s^4} + \cdots \right) ds \\ &= iK' + \frac{1}{kz} + \frac{k^{-1} + k^{-3}}{6z^3} \cdots \end{aligned}$$

The value iK' is gotten by the symmetry of the function: when $z \rightarrow \infty$, exactly half of the rectangle has been traversed.

Exercise 34.1. ** Find changes of variables that connect (34.6) to (34.1)

34.1. Continuation to the whole of $\mathbb{C}$. Double periodicity. As mentioned, after an even number of Schwarz reflections of ψ the function returns to its original value. Recall that an even number of successive Schwarz reflections results in ψ returning to its original value. This is due to the symmetry of the rectangle and would not hold for a general polygon; see the discussion in §30.4.1 The *inverse function* $\text{sn} = \psi^{-1}$ is, by this a doubly periodic function:

$$(34.9) \quad \text{sn}(z) = \text{sn}(z + 2K) = \text{sn}(z + 2iK')$$

R_{-11}	R_{01} iK' ●	R_{11}
R_{-10}	R_{00}	R_{10}
R_{-1-1}	$-K$ 0 K R_{0-1}	R_{1-1}

35. RIEMANN SURFACES: A FIRST VIEW

Here I follow loosely the approach in [5].

By definition, an n -dimensional manifold is a Hausdorff topological space X such that every point $x \in X$ belongs to an open set in X which is homeomorphic to an open subset of $\mathbb{R}^n$.

In particular, when $n = 2$, one can endow a two-dimensional manifold with additional structure that reflects the complex plane. This leads to the notion of a *complex chart* and, more generally, of a *Riemann surface*.

Definition 35.2. Let X be a two-dimensional manifold. A *complex chart* on X is a homeomorphism

$$\varphi : U \rightarrow V$$

of an open subset $U \subset X$ onto an open subset $V \subset \mathbb{C}$. Two complex charts $\varphi_i : U_i \rightarrow V_i$ ($i = 1, 2$) are said to be *holomorphically compatible* if the transition map

$$\varphi_2 \circ \varphi_1^{-1} : \varphi_1(U_1 \cap U_2) \longrightarrow \varphi_2(U_1 \cap U_2)$$

is biholomorphic.

A *complex atlas* on X is a family $\mathcal{F} = \{\varphi_i : U_i \rightarrow V_i\}_{i \in I}$ such that $\{U_i\}_{i \in I}$ covers X and the charts are pairwise holomorphically compatible.

Two complex atlases $\mathcal{F}$ and $\mathcal{F}'$ are said to be *analytically equivalent* if every chart in $\mathcal{F}$ is holomorphically compatible with every chart in $\mathcal{F}'$.

Remark 35.3.

- The restriction of a complex chart φ to a smaller open set $U' \subset U$ is holomorphically compatible with φ .
- Since biholomorphisms preserve analyticity, analytic equivalence of atlases is indeed an equivalence relation.

Definition 35.4. A *complex structure* Σ on a two-dimensional manifold X is an equivalence class of analytically equivalent atlases.

This complex structure Σ is determined by any representative atlas $\mathcal{F}$, and there exists a unique *maximal atlas* $\mathcal{F}^*$ consisting of all complex charts on X that are holomorphically compatible with every chart in $\mathcal{F}$.

Definition 35.5. A Riemann surface Ω is a pair (X, Σ) where X is a connected two-dimensional manifold and Σ is a complex structure on X .

By convention, the complex structure Σ is often omitted from the notation when it is clear from the context. By definition, a Riemann surface is locally equivalent to an open set of $\mathbb{C}$. Since each point of Ω comes equipped with a whole family of indistinguishable charts, the only notions and structures that pass from $\mathbb{C}$ to Ω are those which are *independent of the choice of chart*, i.e. those invariant under biholomorphic maps.

Definition 35.6. A Riemann surface Ω is a connected complex manifold of complex dimension one. This means that Ω is a connected Hausdorff space that is endowed with an atlas of charts to $\mathbb{D}$: for every point $x \in \Omega$ there is a neighborhood of x that is homeomorphic to $\mathbb{D}$, and the transition maps between two overlapping charts are required to be analytic.

Definition 35.7. A Riemann surface Ω is an oriented manifold of (real) dimension two – a two-sided surface – together with a conformal structure. Again, manifold means that locally at any point x of Ω , the space is homeomorphic to a subset of the real plane. The supplement “Riemann” signifies that Ω is endowed with an additional structure which allows angle measurement on the manifold, namely an equivalence class of so-called Riemannian metrics. Two such metrics are considered equivalent if the angles they measure are the same. Choosing an equivalence class of metrics on Ω is the additional datum of the conformal structure.

Remark 35.8. On a real 2-dimensional surface, a Riemannian metric g assigns to each point $x \in \Omega$ an inner product g_x on the tangent space $T_x\Omega$. This inner product defines angles between tangent vectors: for $v, w \in T_x\Omega$,

$$\cos(\theta) = \frac{g_x(v, w)}{\sqrt{g_x(v, v)} \sqrt{g_x(w, w)}}.$$

Two metrics g_1, g_2 are said to be conformally equivalent if $g_2 = \lambda(x)g_1$ for some smooth positive function λ . Conformally equivalent metrics yield the same angles by the formula above, though lengths may differ. A conformal structure on Ω is precisely such an equivalence class of metrics.

Clearly, $\mathbb{C}$ itself is a Riemann surface. So are domains $\mathcal{D} \subset \mathbb{C}$.




FIGURE 40. The Riemann sphere (taken from Google images)

An important surface in complex analysis is the Riemann sphere $S^2 = \hat{\mathbb{C}}$. The complex plane contains the equator of a sphere $\hat{\mathbb{C}}$, represented by $\partial\mathbb{D} \subset \mathbb{C}$. For any point z in the plane, one draws a line segment from the north pole of the sphere through the sphere, cutting the sphere at some point $p(z)$. Clearly, to any point z in the plane, there corresponds a unique $p(z)$ on $\hat{\mathbb{C}}$. $0 \in \mathbb{C}$ corresponds to the south pole in $\hat{\mathbb{C}}$. There is no point in $\mathbb{C}$ corresponding to the north pole on $\hat{\mathbb{C}}$; this is “the point at infinity”. Points in the southern hemisphere map to points in $\mathbb{D}$ while those in the northern hemisphere are in $\overline{\mathbb{D}}^c$.

We will later study the properties of bianalytic transformations, and will see that, in the limit where the size of a triangle goes to zero, a bianalytic (conformal) map takes such a triangle into a similar triangle: the angles are preserved. The triangle is simply rotated and rescaled, in this limit.

Bianalytic transformations are the natural isomorphisms of domains, and more generally of Riemann surfaces in complex analysis. The very important and deep *uniformization theorem* states that, up to such isomorphisms, there are only 3 distinct *simply connected* Riemann surfaces: $\hat{\mathbb{C}}, \mathbb{C}$ and $\mathbb{D}$. Riemann surfaces that are not “very simple” are conformally equivalent to $\mathbb{D}$.

Typically, we cannot embed usefully Riemann surfaces in $\mathbb{R}^3$ and they are not easy to depict. In common cases, a useful way to handle Riemann surfaces is to think of them as spaces of curves, closed or not, modulo homotopies. The endpoints of these curves are the points on

the Riemann surface. Thus, the log that we have defined by integrating ds/s over various paths avoiding zero is naturally generating such a set of equivalence classes.

An important object is **the fundamental group** $\pi(\Omega, p)$ generated by the homotopy classes of *closed curves* joining p to p under composition. Here, the product is concatenation of curves, the inverse of γ , γ^{-1} is $t \mapsto \gamma(1 - t)$ and $1 = \gamma\gamma^{-1}$.




FIGURE 41. The Riemann surface of the log (from https://en.wikipedia.org/wiki/Complex_logarithm)

Exercise 35.9. *Check that the fundamental group of $\mathbb{C} \setminus \{0\}$ is the free group generated by one element, an abelian group.*

Let us uniformize Ω_0 , the Riemann surface of the log. That means finding a bianalytic bijection from Ω_0 to one of $\hat{\mathbb{C}}, \mathbb{C}, \mathbb{D}$.

The log is well defined by $\log z = \int_1^z ds/s$ on curves from 1 to z modulo homotopies in $\hat{\mathbb{C}} \setminus \{0\}$, and is manifestly analytic in z . The fact that it is injective follows from the calculation $\log z_1 = \log z_2 \Rightarrow \exp \log z_1 = \exp \log z_2 \Rightarrow z_1 = z_2$. It is also onto since any $z \neq 0$ can be written as $\log x + i\phi + 2N\pi i$ for some $x \in \mathbb{R}^+$, $\phi \in [0, 2\pi)$ and $N \in \mathbb{Z}$.

The Riemann surface of the log is uniformized by the log onto $\mathbb{C}$.

In the opposite direction, let us look again at the action of the exponential on $\mathbb{C}$. As we have seen, $\exp$ is one-to-one bi-analytic from any vertical strip

$$S_n\{z = u + iv : u \in \mathbb{R}, v \in [2n\pi i, 2(n+1)\pi i)\} \rightarrow \mathbb{C} \setminus \{0\}$$

(think again of the polar representation of complex numbers). If we take $\exp$ to be defined on $\mathbb{C}$, it is not invertible because it is many-to-one. It is not univalent. But in each strip it is one-to-one and we can take as its inverse $\log |z| + i\phi + 2n\pi i$. By taking $\cup_n S_n$ we cover all the possible values of $\log$. We see that, if we define $\exp$ from $\mathbb{C}$ to the Riemann surface of the log, then it is one-to-one, onto and bi-analytic.

Covering maps. A covering map (also called a covering or projection) is a surjective open map $f : X \rightarrow Y$ that is locally a homeomorphism. In a covering map, the preimages $f^{-1}(y)$ are discrete sets in X whose cardinality (finite or infinite) do not depend on y .

Universal covers The universal cover of a connected topological space X is a simply connected space Y with a map $f : Y \rightarrow X$ that is a covering map. If X is simply connected, i.e., has a trivial fundamental group, then it is its own universal cover. For instance, the sphere $S^2 = \hat{\mathbb{C}}$ is its own universal cover. The universal cover is always unique and, under very mild assumptions, always exists. In fact, the universal cover of a topological space X exists iff the space X is connected, locally pathwise-connected, and semilocally simply connected.

We see that the exponential is the covering map of $\mathbb{C} \setminus \{0\}$ and $\mathbb{C}$ is its covering space. Note that, of course, $\mathbb{C} \setminus \{0\}$ is not simply connected while the covering space, $\mathbb{C}$, is. If we look at the image of $f(z) = e^z = ue^{iv}$ where u is fixed and $v \in (a, b) \subset \mathbb{R}$ where $b \leq \infty$ as a parametrized curve, this curves circles, infinitely many times as $b \rightarrow \infty$. around 0. The set of all these curves are the natural curves on which the log is analytic. Here $f^{-1}(y) = \{a + 2n\pi i : e^a = y\}$.

As another example, $f = z \mapsto z^2$ as a map from $\mathbb{C} \setminus \{0\}$ to $\mathbb{C} \setminus \{0\}$ is a covering map in which f^{-1} always consists of two points. The exponential is a universal cover of $\mathbb{C} \setminus \{0\}$ where the inverse image of a point is an infinite set.

As another example, consider the equivalence relation $x + iy \sim x + m + i(y + n)$ iff $(m, n) \in \mathbb{Z}^2$ and the quotient map $\pi : \mathbb{C} \rightarrow \mathbb{C}/\sim$. Then π is the universal cover of the torus $\mathbb{T}$.

The Riemann surface of the square root As the log, the square root, definable in terms of the log is analytic on the Riemann surface of the log. We note however, from its log-based definition, that closed curves starting at some $z_0 \neq 0$, say $z_0 = 1$ that encircle 0 twice yield

the same value of the square root, here 1. What is different from the case of the log is that the fundamental group has a relation: $a^2 = 1$. The Riemann surface of the square root is different from that of the log: the second floor of the “infinite parking lot” is glued to the first. It is not possible to embed this surface in $\mathbb{R}^3$, but we can draw a more or less suggestive picture, below.




FIGURE 42. An approximate depiction of the Riemann surface of the square root (from <https://commons.wikimedia.org/wiki/>)

36. THE MODULAR GROUP

In preparation for elliptic function theory, we now look into the properties of an important subgroup of $PSL(2, \mathbb{R})$ namely the **modular group** $\Gamma = PSL(2, \mathbb{Z})$, course isomorphic to the subgroup of LFT

$$z \mapsto \frac{az + b}{cz + d},$$

$a, b, c, d \in \mathbb{Z}$, $|ad - bc| = 1$, with M and $-M$ identified.

Note 36.10. *The fact that $ad - bc = \pm 1$ implies that the fractions $a/b, a/c, c/d, b/d$ are all irreducible. Check this.*

Definition 36.11 (Lattices). *A two-dimensional lattice generated by the complex numbers, or periods, ω_1 and ω_2 with $\omega_1/\omega_2 \notin \mathbb{R}$ is the set*

$$(36.2) \quad \Lambda_{\omega_1, \omega_2} := \Lambda = \{z \in \mathbb{C} : z = m_1\omega_1 + m_2\omega_2; m_1, m_2 \in \mathbb{Z}\}$$

Any such Λ is a **module** over $\mathbb{Z}$. Geometrically, a lattice provides a tessellation of the plane by identical parallelograms:

Definition 36.12. *A tessellation, or tiling of a region is a family of disjoint open sets whose union of closures cover the region.*

36.1. Bases of lattices. Lattices are clearly discrete sets. Hence there is a ω_1 of minimal absolute value. Choose any $\omega \in \Lambda$ which is not of the form $n\omega_1$ and among these ω choose one of minimal absolute value; call it ω_2 . We must have $\omega_2/\omega_1 \notin \mathbb{R}$, or else if n is s.t. $n < \omega_2/\omega_1 < n+1$, then $|n\omega_1 - \omega_2| < |\omega_1|$, a contradiction.

Proposition 36.13. *$PSL(2, \mathbb{Z})$ is a symmetry of any lattice Λ . In other words $PSL(2, \mathbb{Z})\Lambda = \Lambda$.*

Proof. (Simplified, from [3].) If

$$(36.3) \quad M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in PSL(2, \mathbb{Z})$$

with $a, b, c, d \in \mathbb{Z}, ad - bc = 1$, and $\Lambda = \Lambda_{\omega_1, \omega_2}$ with $\omega_2/\omega_1 \notin \mathbb{R}$, we want to show that $M\Lambda = \Lambda$. Let

$$(36.4) \quad \begin{pmatrix} \omega'_1 \\ \omega'_2 \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix}$$

Note that (ω'_1, ω'_2) and (ω_1, ω_2) play a symmetric role, for

$$(36.5) \quad \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix} = M^{-1} \begin{pmatrix} \omega'_1 \\ \omega'_2 \end{pmatrix}$$

Eq. (36.5) shows in particular that $\omega'_2/\omega'_1 \notin \mathbb{R}$. Check that, for instance by assuming the contrary and arriving at a contradiction. Now, the pair (36.4) and (36.5) shows that $\Lambda_{\omega_1, \omega_2} \subset \Lambda_{\omega'_1, \omega'_2}$ and $\Lambda_{\omega'_1, \omega'_2} \subset \Lambda_{\omega_1, \omega_2}$. ■

Corollary 36.14.

- Any two bases of a lattice Λ are related by a modular transformation (36.4).
- $PSL(2, \mathbb{Z})$ are automorphisms of lattices, and hence of the 2d torus $\mathbb{T}$ (in fact, these are all).




FIGURE 43. The fundamental region $\mathcal{R}$ (the grey region together with the right part of the circle, that is, starting at i) and the action of $PSL(2, \mathbb{Z})$ on it, figure from Wiki: [15].

36.2. The fundamental region of $PSL(2, \mathbb{Z})$. There is a special basis of importance for $PSL(2, \mathbb{Z})$ and for the theory of elliptic functions:

Theorem 36.15. *There exists a basis (ω_1, ω_2) such that the ratio $\tau = \omega_2/\omega_1$ satisfies the following conditions:*

- (1) $\text{Im } \tau > 0$
- (2) $-\frac{1}{2} < \text{Re } \tau \leq \frac{1}{2}$
- (3) $|\tau| \geq 1$
- (4) $\text{Re } \tau \geq 0$ if $|\tau| = 1$.

The ratio τ is uniquely determined by these conditions, and there is a choice of two, four, or six corresponding bases. The region $\mathcal{R}$ is shown in Fig. 36.2.

Proof. (Slightly simplified version of the proof in [3].) Choose ω_1 and ω_2 as in the beginning of §36.1. Then, $|\omega_1| \leq |\omega_2|$, $|\omega_2| \leq |\omega_1 + \omega_2|$ and $|\omega_2| \leq |\omega_1 - \omega_2|$. These translate in $|\tau| \geq 1$ and $|\text{Re } \tau| \geq \frac{1}{2}$. If $\text{Im } \tau < 0$, replacing (ω_1, ω_2) by $(-\omega_1, \omega_2)$ makes $\text{Im } \tau > 0$ while preserving $|\text{Re } \tau| \geq \frac{1}{2}$. Finally, if it happens that $\text{Re } \tau = -\frac{1}{2}$ we pass to $\omega_1, \omega_1 + \omega_2$. If $|\tau| = 1$ and $\text{Re } \tau < 0$ we replace τ by $-\tau^{-1}$, i.e. (ω_1, ω_2) by $(-\omega_2, \omega_1)$ finishing the required normalization. If τ' corresponding to another basis is also in $\mathcal{R}$, then a calculation based on Corollary 36.14 shows that

$$(36.6) \quad \tau' = \frac{a\tau + b}{c\tau + d} \quad \text{and} \quad \text{Im } \tau' = \frac{\text{sign}(ad - bc)\text{Im } \tau}{|c\tau + d|^2} = \frac{\text{Im } \tau}{|c\tau + d|^2}$$

since $\tau' \in \mathcal{R}$; this also implies $ad - bc = 1$. We want to show $\tau' = \tau$ and in this τ and τ' play symmetric roles and we are free to assume

$\text{Im } \tau' \geq \text{Im } \tau$. By (36.6) this gives

$$(36.7) \quad |c\tau + d| \leq 1$$

Writing $\tau = \alpha + i\beta$ we have $\alpha \in (-1/2, 1/2]$, $\beta^2 + \alpha^2 \geq 1$. Eq. (36.7) gives $(\alpha c + d)^2 + \beta^2 c^2 \leq 1$. We must have $|c| \leq 1$ since $|c| \geq 2$ gives $|\beta| \leq \frac{1}{2}$ forcing $|\tau| < 1$.

If $c = 0$, $ad - bc = 1$ implies $ad = 1$, hence $a = d = 1$ or $a = d = -1$. From the first equality in (36.6) we then have $\tau' = \tau \pm b$, hence $\text{Im } \tau = \text{Im } \tau'$, $|b| = |\text{Re } \tau - \text{Re } \tau'| < 1$, and from Assumption 2, $b = 0$ implying $\tau = \tau'$.

If $|c| = 1$, then $|\tau + d| \leq 1$ or $|\tau - d| \leq 1$ implies $d < 2$, hence $d = 0$ or $d = \pm 1$. If $d = 0$, since $|\tau| \geq 1$ we must have $|\tau| = 1$ and now $bc = -1$ gives $b/c = -1$, hence $\tau' = \pm a - 1/\tau = \pm a - \bar{\tau}$ implying $1 > \text{Re}(\tau + \tau') = \pm a$. Hence $a = 0$ and $\tau' = -1/\tau$ which is only possible if $\tau = i$ and $\tau' = i$.

If $d = \pm 1$ then $|\tau \pm 1| \leq 1$. The closest point to -1 in $\bar{\mathcal{R}}$ is $(-\frac{1}{2}, \frac{\sqrt{3}}{2}) \notin \mathcal{R}$, hence $d = -1$ and $|\tau - 1| \leq 1$. Proceeding as before, the closest point to $\bar{\mathcal{R}}$ is $e^{i\pi/3} = (\frac{1}{2}, \frac{\sqrt{3}}{2}) \in \mathcal{R}$, a point of minimal imaginary part, and where $|\tau - 1| = 1$. But by (36.6) $\text{Im } \tau' = \text{Im } \tau$, hence $\tau' = \tau$.

To classify all possible choices of bases we have to find the fixed points in $\mathcal{R}$ of $\text{PSL}(2, \mathbb{Z})$. The matrices -1 are identified and result in $\tau \rightarrow \tau$, but $(-\omega_1, -\omega_2)$ is a different basis. The analysis above shows that the only possible fixed points are $\tau = i$ and $\tau = e^{\pi i/3}$. These are fixed points of $\tau \mapsto 1/\tau$ and $\tau \mapsto (\tau - 1)/\tau$. ■

Exercise 36.16. Find all $z \in \text{UHP}$ for which there is some $M \in \text{PSL}(2, \mathbb{Z})$ s.t. $Mz = z$

We denote the interior of a set A by A° .

Corollary 36.17. If $M \in \text{PSL}(2, \mathbb{Z})$, then $(M\mathcal{R})^\circ \cap \mathcal{R}^\circ \neq \emptyset$ iff $M = 1$, or equivalently, if $A = M\mathcal{R}$ and $A' = M'\mathcal{R}$ then $A^\circ \cap (A')^\circ \neq \emptyset$ iff $M = M'$.

Proof. If $\tau \in (\mathcal{R})^\circ \cap (M\mathcal{R})^\circ$ and $M \neq 1$, then $\tau = M\tau'$, hence τ and τ' belong to the same lattice. By Theorem 36.15, we must have $\tau = M\tau$. Hence, if $M \neq 1$ then $\tau \in (\mathcal{R})^\circ$, contradiction. ■

Definition 36.18. A group G acts on a topological space X in a properly discontinuous way if for all $x \in X$ and $g \in G$ there is a neighborhood $\mathcal{N}$ such that $g(\mathcal{N}) \cap \mathcal{N} \neq \emptyset \Leftrightarrow g = 1$.

Exercise 36.19. Does $\text{PSL}(2, \mathbb{Z})$ act in a properly discontinuous way on the upper half plane?

36.3. The generators of the modular group.

Proposition 36.20. *The modular group is generated by $T = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $S = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, or in terms of LFTs, $\sigma = z \mapsto -1/z$ and $\tau = z \mapsto z + 1$.*

*Proof.*¹⁸ Let $F(z) = \frac{az+b}{cz+d}$. We distinguish three cases.

- (1) $a = 0$. Then $bc = -1$ which can only happen if $b = -c = \pm 1$. In this case, $F(z) = \pm 1/(\pm z + d) = \sigma\tau^{\pm d}(z)$.
- (2) $|a| = 1$. By projective equivalence, we may assume $a = 1$, and then $d - bc = 1$. We bring this transformation back to case 1 by τ, σ :

$$\tau^c \sigma \left(\frac{z+b}{cz+d} \right) = \tau^c \frac{-cz-d}{z+b} = \frac{-cz-d}{z+b} + c = -\frac{1}{z+b}$$

indeed of type (1).

- (3) $|a| > 1$ (it can't be < 1 !). If we show that we can strictly lower $|a|$. Since $a \in \mathbb{Z}$, it means we can eventually bring ourselves to case (2). By Note 36.10 (a and c are coprime) we have $|a| \neq |c|$. By applying σ if necessary we can arrange $|a| > |c|$. We can assume $c \neq 0$ or else σ brings us to (1). Then


$$(36.8) \quad \tau^{\pm 1} F = \frac{(a \pm c)z + (b - d)}{cz + d}$$

■

Exercise 36.21. * 1. Show that the modular group induces a tessellation of the UHP, namely $\{(M\mathcal{R})^\circ : M \in PSL(2, \mathbb{Z})\}$. One way is to apply Corollary 36.17 and note that for any $z \in \mathcal{R}$ and $x \in \mathbb{R}^+$, xz is in the closure of the tessellation by taking $r = p/q \in \mathbb{Q}^+$ and taking LFTs of the form $(npz + b)/(cz + nq)$, $n \in \mathbb{N}, n \rightarrow \infty$. This shows that the strip $\{|\operatorname{Re} z| \leq 1/2, \operatorname{Im} z \geq 0\}$ is in the closure of the tessellation.

2. Find the tessellations of the unit disk induced by conjugation of $PSL(2, \mathbb{Z})$ with the Cayley transform. Find the Poincaré metric on $\mathbb{D}$ using the Cayley transform.

¹⁸Adapted and simplified from a post by rmdmc89

FIGURE 44. Marked action of S, T

36.4. **The hyperbolic plane.** On the UHP define the metric tensor

$$(36.9) \quad ds^2 = \frac{dx^2 + dy^2}{y^2} = \frac{dz d\bar{z}}{y^2}$$

The metric is given by

$$(36.10) \quad \rho(z_1, z_2) = 2 \tanh^{-1} \left| \frac{z_1 - z_2}{z_1 - \bar{z}_2} \right| = \log \frac{|z_1 - \bar{z}_2| + |z_1 - z_2|}{|z_1 - \bar{z}_2| - |z_1 - z_2|}$$

Exercise 36.22. Check that the metric is invariant under LFT's, hence under $SL(2, \mathbb{R})$. Check that the geodesics for this metric are circular arcs $\perp \mathbb{R}$ and vertical lines ending on $\mathbb{R}$.

The hyperbolic plane has constant negative curvature.

36.5. **Schwarz reflection of domains about a circle.** We have seen in Theorem 22.116 that we can analytically continue functions from the upper half plane into a symmetric domain in the lower half plane provided they are continuous up to the real line and real-valued there. Consider a domain $\mathcal{D}$ bounded by a curve and an arccircle C , and let F be analytic in $\mathcal{D}$ and continuous up to C . At any point $z \in C$ we can decompose $f(z) = f_1 + f_2$ where the direction of f_1 is tangent to the circle and f_2 is perpendicular to it. Assume that $f_2 = 0$. Let M

be a Möbius transformation that takes C to a real interval J and $\mathcal{D}$ to a domain $\mathcal{D}'$ in the UHP. Then $f \circ M^{-1}$ is analytic in $\mathcal{D}'$, continuous down to J and real valued on J . Then there is a function analytic F in the domain $\mathcal{D}' \cup \mathcal{D}'' \cup J$ where $\mathcal{D}''$ is the reflection of $\mathcal{D}$ across J s.t. $F|_{\mathcal{D}} = f \circ M^{-1}$ and $F \circ M^{-1}$ provides analytic continuation of f to $\mathcal{D} \cup R\mathcal{D} \cup C$ where $R\mathcal{D}$ is the reflection of $\mathcal{D}$ across the arccircle C .

Exercise 36.23. Find the formula of the reflection of a point across an circle, defined as in the construction above.

37. ENTIRE AND MEROMORPHIC FUNCTIONS

Analytic and meromorphic functions share with polynomials and rational functions a number of very useful properties, such as decomposition by partial fractions and root-factorization. These notions have to be carefully analyzed though, since questions of convergence arise.

37.1. A historical context. Finding the *exact* value of the sum

$$(37.1) \quad S := \sum_{n=1}^{\infty} \frac{1}{n^2}$$

known as the “Basel problem” had been open for almost a century, in spite of efforts by great mathematicians of the time (the Bernoulli brothers, Leibniz, Goldbach, Stirling, Moivre and others) before Euler solved it in 1735 when he was 28; because the problem stumped so many brilliant minds, this attracted a lot of attention. Euler’s solution, well before there was any systematic theory of complex functions proceeds (roughly) as follows. Take the function

$$(37.2) \quad f(x) = \frac{\sin x}{x}$$

This has a Taylor series which converges for all $x \in \mathbb{C}$. If f were a polynomial with roots at $z = a_i$, then we would be able to write

$$(37.3) \quad f(x) = A \prod (x - a_i)$$

Assuming the same were true for an “infinite order polynomial” and noting that the roots of $\sin(x)/x$ are at $n\pi$, $n \in \mathbb{Z} \setminus \{0\}$ we get

$$(37.4) \quad \frac{\sin x}{x} = \prod_{n \geq 1} \left(1 - \frac{x}{n\pi}\right) \left(1 + \frac{x}{n\pi}\right) = \prod_{n \geq 1} \left(1 - \frac{x^2}{(n\pi)^2}\right)$$

(here the free constant must be 1, comparing the values at 0) Expanding out and collecting the coefficient of x^2 we get $\frac{\sin x}{x} = 1 - cx^2 + O(x^4)$

where

$$(37.5) \quad c = \frac{1}{\pi^2} + \frac{1}{4\pi^2} + \cdots = \frac{1}{\pi^2} S$$

On the other hand from the Maclaurin expansion of $f(x)$ we have $c = \frac{1}{3!}$. Thus

$$(37.6) \quad S = \frac{\pi^2}{6}$$

In fact, Euler went further and calculated $\sum \frac{1}{n^{2k}}$ for $k \geq 1$, in for all even k . This was mostly fine by the standards of the day, though it led to some criticisms that prompted more rigorous proofs later by Euler. Weierstrass was apparently inspired by this solution when he developed the theory of decomposition of entire functions as products. For a rigorous proof of (37.5) see §38.2

37.2. Partial fraction decompositions. First let $R = P_0/Q = P_1 + P/Q$ be a rational function. where P_i and Q are polynomials and $\deg(P) < \deg(Q)$. We aim at a partial fraction decomposition of R ; if $\deg(Q) = 0$ there is nothing further to do. Otherwise let $z_1, \dots, z_n$, $n \geq 1$, be the zeros of Q , where we don't count the multiplicities, and let m_j be the multiplicities of these roots. Let's look at the singular part of the Laurent expansion of P/Q at z_j :

$$(37.7) \quad \frac{P}{Q} = \sum_{k=1}^{m_j} \frac{c_{jk}}{(z - z_j)^k} + \text{analytic at } z_j$$

We claim that

$$(37.8) \quad \frac{P}{Q} = \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{c_{jk}}{(z - z_j)^k}$$

Indeed,

$$(37.9) \quad E(z) := \frac{P}{Q} - \sum_{j=1}^n \sum_{k=1}^{m_j} \frac{c_{jk}}{(z - z_j)^k}$$

is an entire function. By assumption, $P/Q \rightarrow 0$ as $z \rightarrow \infty$ and the rhs of (37.8) also, clearly, goes to zero as $z \rightarrow \infty$. Thus $E(z) \rightarrow 0$ as $z \rightarrow \infty$ and therefore $E \equiv 0$.

Second example. We examine in detail an example that has some of the features of the general case. The function

$$(37.10) \quad \frac{\pi^2}{\sin^2 \pi z}$$

has poles for $z_i = k$, $k \in \mathbb{Z}$, and the singular part of the Laurent series at $z = k$ is, as it can be quickly checked

$$(37.11) \quad \frac{1}{(z - k)^2}$$

We first note that

$$g(z) = \sum_{k \in \mathbb{Z}} \frac{1}{(z - k)^2}; \quad z \in \mathbb{C} \setminus \mathbb{Z}$$

converges uniformly and absolutely on compact sets in $\mathbb{C} \setminus \mathbb{Z}$ to an analytic function: take z_0 s.t. $\text{dist}(z, \mathbb{Z}) > \varepsilon$ and the disk $\mathbb{D}_{\varepsilon/2}(z_0)$. By translation symmetry we may assume that $\lfloor x_0 \rfloor = \alpha \in [0, 1)$. We can write

$$g(z) = \sum_{|k| \leq 1} \frac{1}{(z - k)^2} + \sum_{|k| \geq 2} \frac{1}{(z - k)^2}$$

The second sum is bounded in absolute value by $\sum_{|k| \geq 1} \frac{1}{k^2}$, while the finite sum is a manifestly analytic function. Next we note that

$$(37.12) \quad \frac{\pi^2}{\sin^2 \pi z} - g(z)$$

extended by continuity at the integers is entire. This can be immediately checked by noting that the only singularities of $h(z) = \frac{\pi^2}{\sin^2 \pi z}$ are $z = k \in \mathbb{Z}$, and calculating the Laurent series of h at $z = k$.

Proposition 37.1. *We have*

$$(37.13) \quad \frac{\pi^2}{\sin^2 \pi z} = \sum_{k \in \mathbb{Z}} \frac{1}{(z - k)^2}; \quad z \notin \mathbb{Z}$$

Proof. In the notations introduced above the proposition, we saw that, after extending it by continuity at the removable singularities, the function $h - g$ is entire. This function is clearly periodic of period 1, and it is clearly bounded in any compact subset of the strip $\text{Re } z \in [0, 1]$. By dominated convergence $g(z) \rightarrow 0$ as $\text{Im } z \rightarrow \infty$. Now, with $z = x + iy$ we have

$$|\sin(x + iy)^2| = \frac{1}{2} \left(\cosh(2y) - \cos(2x) \right) \geq \frac{1}{2} (\cosh(2y) - 1)$$

It follows that $h - g$ is entire and vanishes at infinity, thus it is identically zero.

■

Exercise 37.2. By integrating (37.13) show that

$$\cot(x) = \frac{1}{x} + \sum_{n=1}^{\infty} \frac{2x}{n^2\pi^2 - x^2}$$

By one more integration, show the product formula Euler used. (See also Lemma 38.5 below.)

37.3. The Mittag-Leffler theorem. As such, the procedure in Proposition 37.1 does not extend in general. For instance, for the function Γ'/Γ the poles are located at the negative integers and have residue 1, and the partial fraction decomposition would not converge without some modifications. Furthermore, even when it does converge, a function can be determined by a sum over its poles only up to an entire function. For a general theorem, we need to address these issues.

The theorem below shows that for any sequence of one-sided Laurent series centered at the points $\{b_n\}_{n \in \mathbb{N}} \in \mathbb{C}$ with no accumulation point, there is a meromorphic function with exactly that singular behavior and analytic elsewhere. Conversely, any meromorphic function can be decomposed as a sum of its negative powers-part of its Laurent series at the poles and an entire function. More precisely,

Theorem 37.3 (Mittag-Leffler). (i) Let $\{b_n\}_{n \in \mathbb{N}}$ be a sequence of complex numbers with no accumulation point (cf. Note 37.4) in $\mathbb{C}$ and let $\{P_n\}_{n \in \mathbb{N}}$ be a sequence of polynomials without constant term. Then there are meromorphic functions f in $\mathbb{C}$ whose only poles are at $z = b_n$ and having singular part $P_n((z - b_n)^{-1})$ at $z = b_n$.

(ii) Conversely let f be meromorphic in $\mathbb{C}$ whose poles are $\{b_n\}_{n \in \mathbb{N}}$ and singular part of the Laurent expansion $P_n((z - b_n)^{-1})$ at b_n , $n \in \mathbb{N}$. Then there exists a sequence of polynomials $\{p_n\}_{n \in \mathbb{N}}$ and an entire function g such that

$$(37.14) \quad f = \sum_{n \in \mathbb{N}} \left[P_n \left(\frac{1}{z - b_n} \right) - p_n(z) \right] + g(z) := S(z) + g(z)$$

where the series converges uniformly on compact sets in $\mathbb{C} \setminus \{b_n\}_{n \in \mathbb{N}}$.

Note 37.4. The condition that $\{b_n\}_{n \in \mathbb{N}}$ has no accumulation point in $\mathbb{C}$ automatically implies that $|b_n| \rightarrow \infty$ as $n \rightarrow \infty$.

Note 37.5 (Idea of the construction). The possible divergence of the infinite series $\sum_{n \in \mathbb{N}} P_n((z - b_n)^{-1})$ is due to the behavior for large n of the terms of the series. We fix z and take n large enough so that $|z/b_n| < 1$. For large n , $\frac{1}{z - b_n}$ roughly behaves like $-b_n^{-1}$. If we subtract out $-1/b_n$, the behavior is roughly $-zb_n^{-2}$. Continuing in this way, and

keeping in mind that $|b_n| \rightarrow \infty$ as $n \rightarrow \infty$, we obtain for each n a bound $|z|^{m_n-1}|b_n|^{-m_n}$ ultimately achieving convergence.

Proof. We start by proving (i). We can assume without loss of generality that $b_n \neq 0$ for if say $b_1 = 0$ then we can prove the theorem for $\tilde{f} = f - P_1(1/z)$. Note first that the series at infinity in $1/b_n$ of $P_n(z - b_n)^{-1}$ and its series in z at zero coincide.

Let p_n be a Maclaurin polynomial of P_n such that

$$(37.15) \quad |P_n((z - b_n)^{-1}) - p_n(z)| \leq 2^{-n}; \quad \forall z \text{ s.t. } |z| < |b_n|/4$$

Now we look at the series

$$(37.16) \quad f_1 = \sum_{n=1}^{\infty} [P_n((z - b_n)^{-1}) - p_n(z)]$$

and fix an R and analyze the series for $z \in \mathbb{D}_R$. We split the sum (37.15) into two parts:

$$(37.17) \quad f_1 = \sum_{n: |b_n| \leq 4R} [P_n((z - b_n)^{-1}) - p_n(z)] + \sum_{n: |b_n| > 4R} [P_n((z - b_n)^{-1}) - p_n(z)]$$

The first sum is finite, while for the second one we have $|z| < |b_n|/4$ and the estimate (37.15) applies. Thus (37.16) is convergent away from the poles uniformly on compact sets.

(ii) Clearly, $f - f_1$ is entire. ■

37.4. Further examples. We write the result in Exercise 37.2 in the form

$$(37.18) \quad \pi \cot \pi z = \frac{1}{z} + \sum_{n \in \mathbb{N} \setminus \{0\}} \frac{2z}{z^2 - n^2}$$

We can now use this identity to calculate easily some familiar sums. Note that the lhs of (37.18) has the Laurent expansion at $z = 0$

$$(37.19) \quad \pi \cot \pi z = \frac{1}{z} - \frac{\pi^2 z}{3} - \frac{\pi^4 z^3}{45} - \frac{2\pi^6 z^5}{945} - \dots$$

Since the series on the rhs of (37.18) converges uniformly near $z = 0$, by Weierstrass's theorem it converges together with all derivatives. On the other hand we have

$$(37.20) \quad \frac{2z}{z^2 - n^2} = -2 \left(\frac{z}{n^2} + \frac{z^3}{n^4} + \frac{z^5}{n^6} + \dots \right)$$

and we get immediately,

$$(37.21) \quad \sum_{n \geq 1} \frac{1}{n^2} = \frac{\pi^2}{6}, \quad \sum_{n \geq 1} \frac{1}{n^4} = \frac{\pi^4}{90}, \quad \sum_{n \geq 1} \frac{1}{n^6} = \frac{\pi^6}{945} \dots$$

Exercise 37.6. * The definition of the Bernoulli numbers B_k is

$$(37.22) \quad \frac{1}{e^z - 1} = \frac{1}{z} - \frac{1}{2} + \sum_{k=1}^{\infty} (-1)^{k-1} \frac{B_k}{(2k)!} z^{2k-1}$$

Show that

$$(37.23) \quad \sum_{n=1}^{\infty} \frac{1}{n^{2k}} = 2^{2k-1} \frac{B_k}{(2k)!} \pi^{2k}$$

38. INFINITE PRODUCTS

An infinite product is the limit

$$(38.1) \quad \prod_{n=1}^{\infty} p_n := \lim_{k \rightarrow \infty} \prod_{n=1}^k p_n = \lim_{k \rightarrow \infty} \Pi_k$$

We wish to express entire functions as infinite products. Then the following conventions are natural:

1. Only finitely many p_j are nonzero.

Hence there exists an n_0 such that $p_k \neq 0$ for any $k \geq n_0$ and we require:

- 2.

$$\prod_{k=n_0}^n p_k \rightarrow \ell \neq 0 \quad \text{as } n \rightarrow \infty$$

In issues of convergence we ignore the zero terms, so in the following we assume they simply do not exist. Write now $p_n = P_n/P_{n-1}$, with $P_0 = 1$ we see that the limit of Π_k is the same as the limit of the P_k , and since $P_k \rightarrow \ell \neq 0$ we have $P_{k+1}/P_k \rightarrow 1$ as $k \rightarrow \infty$. Thus $p_n \rightarrow 1$ is a necessary condition of convergence of the infinite product. We should then better write the products as

$$(38.2) \quad \prod_{n=1}^{\infty} (1 + a_n)$$

and then a necessary condition of convergence is $a_n \rightarrow 0$.

Theorem 38.1. The infinite product (38.2) converges iff

$$(38.3) \quad \sum_{n=1}^{\infty} \ln(1 + a_n)$$

converges. We omit, as before, the terms with $a_n = -1$. The log is defined with a cut along $(-\infty, 0]$, extended by continuity on the upper side of the cut.

Proof. If the sum (38.3) converges, then $\prod_n$ converges, since the exponential of a finite sum is a finite product.

In the opposite direction, a word of caution. We know that in the complex domain, $\ln ab$ is not always $\ln a + \ln b$. The limit of the sum will not, in general, be the log of the infinite product. So the reasoning is not that obvious. We write $(1 + a_n) = \rho_n e^{i\phi_n}$ and note that $\prod \rho_n$ is convergent, hence the real part of (3.14) converges. Since $(1 + a_{n+1})/(1 + a_n) \rightarrow 1$ we can inductively choose a multiple of $2\pi i$ in $\arg(1 + a_n)$ so that

$$(38.4) \quad \phi_{n+1} - \phi_n \rightarrow 0 \text{ as } n \rightarrow \infty$$

With this choice, we see that $\sum_k \phi_k$ must converge. Otherwise, there are two possibilities. If $S_n = \sum_{k=1}^n \phi_k$ is unbounded, then, by (38.4), $\text{dist}(\{S_n\}_{n \in \mathbb{N}}, 2\mathbb{Z} + 1) = \text{dist}(\{S_n\}_{n \in \mathbb{N}}, 2\mathbb{Z}) = 0$ entailing infinite oscillation and divergence of the product. If instead $\{S_n\}_{n \in \mathbb{N}}$ is confined to a compact set and has $a \neq b$ as accumulation points, then the conclusion is the same. Hence, the imaginary part of (3.14) converges as well. ■

Absolute convergence is easier to control in terms of series. An infinite product is absolutely convergent, by definition, iff

$$(38.5) \quad \sum_{n=1}^{\infty} |\ln(1 + a_n)|$$

is convergent.

Theorem 38.2. *The sum (38.5) is absolutely convergent iff $\sum a_k$ is absolutely convergent.*

Proof. Assume $\sum a_k$ converges absolutely. Then in particular $a_n \rightarrow 0$. Also, if $\sum_{n=1}^{\infty} \ln(1 + a_n)$ converges absolutely then $\ln(1 + a_n) \rightarrow 0$ and $a_n \rightarrow 0$. But (eliminating all the irrelevant zero terms which are zero) we have, as $n \rightarrow \infty$ $\lim_{n \rightarrow \infty} |a_n|^{-1} \ln(1 + |a_n|) = 1$, and the rest follows from the limit ratio theorem. ■

Note 38.3. *Conditional (not absolute) convergence of $\sum a_n$ and of $\prod(1 + a_n)$ are unrelated notions. (Consider, e.g., the product $\prod(1 - (-1)^n n^{-1/2})$. Is the associated series $\sum(-1)^n n^{-1/2}$ convergent? Is the product convergent?)*

38.1. Uniform convergence of products.

Exercise 38.4. *Assume that $p_n(z)$ are analytic in the domain $\mathcal{D}$. Show that $f(z) = \prod_{n \geq 1} p_n(z)$ converges absolutely and uniformly on every*

compact set in the domain $\mathcal{D}$ iff

$$(38.6) \quad f'(z) = \sum_{k=1}^{\infty} \prod_{n=1}^{\infty} \frac{p'_k}{p_k} p_n$$

converges absolutely and uniformly on every compact set in the domain $\mathcal{D}$ (if some p_k are zero, the terms are understood in the sense of removable singularities).

38.2. Example: the sin function. Let us prove directly Euler's product formula for the sin function. A good product formula candidate is indeed

$$(38.7) \quad C\pi \sin \pi z \stackrel{?}{=} z \prod_{n>0} \left(1 - \frac{z^2}{n^2}\right)$$

The constant C can only be $1/\pi^2$ if we look at the behavior near $z = 0$. Thus,

$$(38.8) \quad \frac{\sin \pi z}{\pi} \stackrel{?}{=} z \prod_{n>0} \left(1 - \frac{z^2}{n^2}\right)$$

This equality of course needs to be proved, but this is not difficult.

First we note that the product on the rhs of (38.8) is absolutely and uniformly convergent on any compact z set; this can be easily checked. It thus defines an entire function $g(z)$. Motivated by the way we obtained this possible identity, let us look at the expression $f'/f - g'/g$ where $\pi f(z) = \sin \pi z$. We get, using Exercise 38.4,

$$(38.9) \quad f'/f - g'/g = \pi \cot \pi z - \frac{1}{z} + \sum_{n \in \mathbb{N} \setminus \{0\}} \frac{2z}{z^2 - n^2} = 0$$

This means that

$$(38.10) \quad \frac{f'g - fg'}{fg} = 0$$

in $\mathbb{C} \setminus \mathbb{Z}$ or, equivalently,

$$(38.11) \quad \frac{f'g - fg'}{g^2} = 0 = \left(\frac{f}{g}\right)'$$

or $f/g = \text{const}$; we already calculated the constant based on the behavior at zero, it is one. Thus indeed,

$$(38.12) \quad \frac{\sin \pi z}{\pi z} = \prod_{n>0} \left(1 - \frac{z^2}{n^2}\right)$$

proving (37.5). More generally, we have the following:

Lemma 38.5. Assume that $\{f_n\}$ are analytic and nonzero in a domain $\mathcal{D}$ containing zero, that $f_n(0) = 1$ and that $\sum_{k=1}^n f'_k/f_k$ converges uniformly on compact sets in $\mathcal{D}$. Then $\{\prod_{k=1}^n f_k\}_{n \in \mathbb{N}}$ converges uniformly on compact sets in $\mathcal{D}$, to a function analytic in $\mathcal{D}$.

Proof. Under these assumptions, $h_n(z) = \int_0^z \sum_{k=1}^n f'_k(s)/f_k(s) ds$ also converges uniformly on compact sets in $\mathcal{D}$. Defining $g(z) = e^{-h_n(z)} \prod_{k=1}^n f_k$ and using Exercise 38.4 we have $g'_n(z) = 0$ and $g(0) = 1$. The rest is immediate. ■

38.3. Canonical products. The simplest possible case is that in which we have a function with no zeros.

Theorem 38.6. Assume f is entire and $f \neq 0$ in $\mathbb{C}$. Then f is of the form

$$(38.13) \quad f = e^g$$

where g is also entire.

Proof. Since f'/f is entire and $\mathbb{C}$ is simply connected, $h(z) = \int_0^z f'(s)/f(s) ds$ is well defined and also entire. Now we note that $(fe^{-h})' = 0$ in $\mathbb{C}$ and thus $f = \exp(h + C)$ proving the result. Another proof is by using the monodromy theorem and the fact that $\log f$ has no singularities in $\mathbb{C}$. ■

Assume now that f has finitely many zeros, a zero of order $m \geq 0$ at the origin, and the nonzero ones, possibly repeated are $a_1, \dots, a_n$.

Then

$$f = z^m \prod_{k=1}^n \left(1 - \frac{z}{a_k}\right) e^{g(z)}$$

where g is entire.

This is clear, since if we divide f by the prefactor of e^g we get an entire function with no zeros.

We cannot expect, in general, such a simple formula to hold if there are infinitely many zeros. Again we have to take care of convergence problems. This is done in a manner similar to that used in the Mittag-Leffler construction.

Theorem 38.7 (Weierstrass). (i) If $\{a_n\}_{n \in \mathbb{N}}$ is a sequence with no accumulation points, and $\{m_n\}_{n \in \mathbb{N}} \subset \mathbb{N}$ is a set of multiplicities then there exists an entire function with zeros at a_n of multiplicities m_n and no other zeros.

(ii) Assume f is an entire function with zeros at a_n of multiplicities m_n . Then there exist polynomials q_n , and an entire function $g(z)$ such that

$$(38.14) \quad f(z) = e^{g(z)} z^m \prod_{n \in \mathbb{N}} \left[\left(1 - \frac{z}{a_n} \right)^{m_n} e^{q_n} \right]$$

Proof. This is a consequence of Mittag-Leffler. For the proof, we may assume that $f(0) \neq 0$, for otherwise, if f has a zero of order m at 0 we prove the theorem first for $\tilde{f}(z) = f(z)z^{-m}$ understood as usual after removing the singularity at zero.

Take a meromorphic function h with simple poles at a_n of residues m_n as provided by Mittag-Leffler:

$$(38.15) \quad h(z) = \sum_{n \in \mathbb{N}} \left(\frac{m_n}{z - a_n} - p_n \right)$$

We claim that $e^{\int_0^z h(s)ds}$ understood by analytic continuation from a neighborhood of zero has the desired properties. Indeed, in a neighborhood of 0 we write

$$(38.16) \quad \frac{m_n}{z - a_n} = \frac{f'_n(z)}{f_n(z)} \quad \text{where} \quad f_n(z) = \left(1 - \frac{z}{a_n} \right)$$

we choose any branch of $\log(1 - z/a_n)$ (a multiple of $2\pi i$ will not matter); the natural choice is that given by the power series. We choose a disk of radius r $\mathbb{D}_r$ and choose N_r so that for $n \geq N_r$ $a_n \notin \mathbb{D}_r$. In $\mathbb{D}_r$ we have

$$(38.17) \quad \int_0^z h(s)ds = \sum_{n \leq N(r)} m_n \log(1 - z/a_n) - \int_0^z p_n(s)ds + \int_{n \geq N(r)} \left(\frac{m_n}{z - a_n} - p_n \right)$$

Once more, the integral is understood as being well defined near zero, and analytically continued to $\mathbb{D}_r$. In $\mathbb{D}_r$ we have

$$(38.18) \quad e^{\int_0^z h(s)ds} = \prod_{n \leq N(r)} (1 - z/a_n)^{m_n} e^{\int_{n \geq N(r)} \left(\frac{m_n}{s - a_n} - p_n(s) \right) ds}$$

where the exponential on the right side of (38.18) is analytic. For (ii) we note that if f is entire with the prescribed zeros and multiplicities and H is as in (i), then f/H is entire with no zeros and the result follows from Theorem 38.6. ■

Note 38.8. *Historically, Weierstrass' factorization theorem came first. Gösta Mittag-Leffler was inspired by Weierstrass and found his decomposition theorem.*

Corollary 38.9. *Any meromorphic function is a ratio of entire functions.*

Proof. Let F be meromorphic with poles at b_n of order m_n . Let G be any entire function with zeros at b_n of order m_n . Then FG has only removable singularities. ■

38.4. Counting zeros of analytic functions. Jensen's formula.

The rate of growth of an analytic function is closely related to the density of zeros. A quite effective counting theorem is due to Jensen.

Theorem 38.10 (Jensen). *Assume $f \not\equiv 0$ is analytic in the closed disk $\overline{\mathbb{D}_r}$ and $f(z) = cz^m g(z)$ with $m \geq 0$ and $g(0) = 1$. Let a_i be the nonzero roots of f in $\mathbb{D}_r$, repeated according to their multiplicity. Then*

$$(38.19) \quad \ln |c| = -m \ln r - \sum_{i=1}^n \ln \left(\frac{r}{|a_i|} \right) + \frac{1}{2\pi} \int_0^{2\pi} \ln |f(re^{i\theta})| d\theta$$

Proof. The proof will follow quite easily from the case when $f \not\equiv 0$ in $\mathbb{D}_r$. In this case we can define a consistent branch of $\ln f$ in $\mathbb{D}_r$ (see also the second proof of Theorem 38.6), and $\operatorname{Re} \ln f = \ln |f|$ is harmonic in $\mathbb{D}_r$. For $r' < r$, by the mean value theorem for harmonic functions, we have

$$(38.20) \quad \ln |f(0)| = \frac{1}{2\pi} \int_0^{2\pi} \ln |f(r'e^{i\theta})| d\theta$$

Since f is analytic in the closed disk and $\ln |x|$ is in $L^1(\mathbb{R})$, it is easy to see by dominated convergence (check) that (38.20) holds in the limit $r = r'$ too, even if there are zeros on the circle of radius r :

$$(38.21) \quad \ln |f(0)| = \frac{1}{2\pi} \int_0^{2\pi} \ln |f(re^{i\theta})| d\theta$$

Assume now f has zeros, with the convention in the statement of the theorem. We then build a function which has no zeros inside $\mathbb{D}_r$ and has the same absolute value for $|z| = r$. Such a function is

$$(38.22) \quad h(z) = \frac{r^m}{z^m} f(z) \prod_{i=1}^n \frac{r^2 - \overline{a_i}z}{r(a_i - z)}$$

(understood as usual with the removable singularities removed). Then

$$(38.23) \quad \ln |h(0)| = \frac{1}{2\pi} \int_0^{2\pi} \ln |h(re^{i\theta})| d\theta = \frac{1}{2\pi} \int_0^{2\pi} \ln |f(re^{i\theta})| d\theta$$

We have $h(0) = r^m c \prod_{i=1}^n \frac{r}{a_i}$ and the formula follows from the case with no roots. ■

Corollary 38.11. *Assume f is analytic in the closed disk of radius R and $f(0) \neq 0$. Let $\nu(r)$ denote the number of zeros of f in the disk of radius $r \leq R$. Then*

$$(38.24) \quad \int_0^R \frac{\nu(x)}{x} dx \leq \ln \max_{|z|=R} |f(z)| - \ln |f(0)|$$

(Note that for some $\varepsilon > 0$, $\nu(x) = 0$ for $x < \varepsilon$.) Clearly, $\nu(x)$ is an increasing (discontinuous) function of x .

Proof. Note that

$$\ln(R/|a_i|) = \int_{|a_i|}^R \frac{dx}{x} = \int_0^R \chi([|a_i|, x]) \frac{dx}{x}$$

Thus

$$\sum_{i=1}^n \ln \left(\frac{R}{|a_i|} \right) = \int_0^R \sum_{i=1}^n \chi([|a_i|, x]) \frac{dx}{x} = \int_0^R \frac{\nu(x)}{x} dx$$

The rest follows immediately from (38.19). ■

38.5. Entire functions of finite order. Let f be an entire function. We denote

$$\|f\|_R = \sup_{|z| \leq R} |f(z)| = \sup_{|z|=R} |f(z)|$$

A function is of **order** $\leq \rho$ if for all R large enough we have

$$(38.25) \quad \forall \varepsilon > 0 \exists c_\varepsilon \text{ s.t. } \|f\|_R \leq \exp(c_\varepsilon R^{\rho+\varepsilon})$$

or equivalently

$$(38.26) \quad \forall \varepsilon > 0 \ln \|f\|_R = O(R^{\rho+\varepsilon})$$

Note 38.12. *We can always check the condition for $R \in \mathbb{N}$ large enough since $(N+1)^\rho = O(N^\rho)$.*

The function f has **strict order** $\leq \rho$ if there is some $c > 0$ such that for all R large enough we have

$$(38.27) \quad \|f\|_R \leq \exp(cR^\rho)$$

A function has **order equal** ρ if

$$\rho = \inf\{\rho' : \forall \varepsilon > 0 \exists c_\varepsilon \text{ s.t. } \|f\|_R \leq \exp(c_\varepsilon R^{\rho'+\varepsilon})\}$$

A function has **strict order equal** ρ if

$$\rho = \inf\{\rho' : \forall R > 0, \|f\|_R \leq \exp(cR^{\rho'})\}$$

Exercise 38.13. * Check that $\cosh z^{1/2}$ has strict order $1/2$ and $\cos z^{1/4} + \cosh z^{1/4}$ has strict order $1/4$. Variations of this construction lets you find entire functions of any rational order. But can you find an entire function of exact order π ?

Proposition 38.14. Assume $f(z)$ is entire, and for large $|z|$ there are positive constants C , c and ρ such that $|f(z)| \leq Ce^{c|z|^\rho}$. Then, for $R > 0$ there is a $c_2 > 0$ such that

$$(38.28) \quad \nu(R) \leq c_2 R^\rho$$

Indeed, the zeros at zero do not change the shape of the inequality, and we can thus assume $f(0) \neq 0$. Then,

$$(38.29) \quad c|R|^\rho \geq \int_0^R \frac{\nu(x)}{x} dx \geq \int_{R/2}^R \frac{\nu(x)}{x} dx \geq \frac{\nu(R/2)}{R} \frac{R}{2}$$

and the rest is immediate. The constants in the inequality can be optimized by choosing R/τ , $\tau > 1$ instead of $R/2$ and finding the best τ .

Theorem 38.15. Let f be entire of strict order $\leq \rho$ and let $\{z_n\}$ be its nonzero zeros, repeated according to their multiplicity and ordered increasingly by their absolute value. Then for any $\varepsilon > 0$, $\{|z_n|^{-1}\}_{n \in \mathbb{N}} \in \ell^{\rho+\varepsilon}(\mathbb{N})$, i.e., the series

$$(38.30) \quad \sum_{n=1}^{\infty} \frac{1}{|z_n|^{\rho+\varepsilon}}$$

is convergent.

Proof. We can obviously discard the roots with $|z_i| \leq 1$ which are in finite number. Without loss of generality we assume there are none. We have, with $N \in \mathbb{N}$ and estimating the sum by annuli,

$$(38.31) \quad \sum_{|z_n| \leq N} \frac{1}{|z|^{\rho+\varepsilon}} \leq \sum_{k=1}^N \frac{\nu(k+1) - \nu(k)}{k^{\rho+\varepsilon}}$$

we can now use the method of Abel summation by parts. We write

$$(38.32) \quad \frac{\nu(k+1) - \nu(k)}{k^{\rho+\varepsilon}} = \nu(k+1) \left(\frac{1}{k^{\rho+\varepsilon}} - \frac{1}{(k+1)^{\rho+\varepsilon}} \right) + \left(\frac{\nu(k+1)}{(k+1)^{\rho+\varepsilon}} - \frac{\nu(k)}{k^{\rho+\varepsilon}} \right)$$

and note that by summation, the terms in the last parenthesis cancel out to

$$\frac{\nu(N+1)}{(N+1)^{\rho+\varepsilon}} - \nu(1)$$

Note that by the mean value theorem we have for some $\gamma = \gamma(k)$

$$(38.33) \quad \frac{\nu(k+1)}{k^{\rho+\varepsilon}} - \frac{\nu(k+1)}{(k+1)^{\rho+\varepsilon}} = \frac{(\rho+\varepsilon)\nu(k+1)}{(k+\gamma)^{\rho+\varepsilon+1}} \leq \frac{Ck^\rho(\rho+\varepsilon)}{k^{\rho+\varepsilon+1}}$$

and the sum converges. $\blacksquare$

38.6. Estimating analytic functions by their real part.

Theorem 38.16 (Borel-Carathéodory). *Let $R > 0$ and assume $f = u + iv$ is analytic $\mathbb{D}_R$. Let $A_R = \max_{|z|=R} u(z)$. Then for $r < R$ we have*

$$(38.34) \quad \max_{|z| \leq r} |f(z)| \leq \frac{2rA_R}{R-r} + \frac{R+r}{R-r} |f(0)|$$

Note that if f is entire and nonconstant, then $\|f\|_R \rightarrow \infty$ as $R \rightarrow \infty$. Then, since $|u| \leq |f|$, the theorem above shows that

$$(38.35) \quad \lim_{R \rightarrow \infty} \frac{\max_{|z| \leq R} u(z)}{\max_{|z| \leq R} |f(z)|} \geq \frac{1}{2}$$

Proof. Assume first that $f(0) = 0$. Then $u(0) = 0$ and by the mean value theorem $A_R \geq 0$. If $A_R = 0$ then by the same argument $u \equiv 0$ on $\partial\mathbb{D}_R$ and by Poisson's formula $u \equiv 0$ in $\mathbb{D}_R$. Then $v \equiv \text{const} = 0$ since $f(0) = 0$, thus $f \equiv 0$ and the formula holds trivially.

We now assume $A_R > 0$. Since the maximum of a harmonic function is reached on the boundary, we have $2A_R - u \geq u$ in $\mathbb{D}_R$ and the inequality is strict in the interior. Also note that if at some point $u < 0$, then again $2A_R - u = 2A_R + |u| \geq |u|$. In $\mathbb{D}_R$ we have

$$(38.36) \quad |2A_R - f| = \sqrt{(2A_R - u)^2 + v^2} > \sqrt{u^2 + v^2} = |f|$$

Hence the function

$$(38.37) \quad g(z) = \frac{1}{2A_R - f(z)} \frac{f(z)}{z}$$

is holomorphic in $\mathbb{D}_R$ and we have, by passing to the limit $|z| \rightarrow R$,

$$(38.38) \quad |g(z)| = \left| \frac{1}{2A_R - f(z)} \frac{f(z)}{z} \right| \leq \frac{1}{R}$$

hence

$$(38.39) \quad \left| \frac{f(z)}{z} \right| \leq \frac{1}{R} |2A_R - f(z)| \leq \frac{1}{R} (2A_R + |f(z)|)$$

Solving for $|f(z)|$ we get

$$(38.40) \quad |f(z)| \leq \frac{2|z|A_R}{R-|z|}$$

as claimed. The general case is obtained by applying this inequality to $f(z) - f(0)$ (exercise). ■

Corollary 38.17. *Assume $\rho \geq 0$, $f = u + iv$ is entire and as $|z| \rightarrow \infty$ we have*

$$(38.41) \quad |u(z)| \leq C|z|^\rho$$

Then f is a polynomial of degree at most ρ .

Proof. From (38.35), for any $\varepsilon > 0$ we have, for large R $|f(z)| \leq (2C + \varepsilon)|z|^\rho$ and the rest is a simple exercise. ■

39. HADAMARD'S THEOREM

Let $\rho > 0$ and let k_ρ be the smallest integer *strictly* greater than ρ , $k_\rho = \lfloor \rho \rfloor + 1$. We consider again the truncates of the series of $-\ln(1-z)$, namely, with $k = k_\rho$,

$$(39.1) \quad P_k(z) = z + \frac{z^2}{2} + \cdots + \frac{z^{k-1}}{k-1}$$

Theorem 39.1 (Hadamard). *Let f be entire of order ρ , let z_n be its nonzero zeros and let $k = k_\rho$. Then, with $m \geq 0$ the order of the zero of f at zero, there is a polynomial h of degree $\leq \rho$ such that*

$$(39.2) \quad f(z) = e^{h(z)} z^m \prod_{n=1}^{\infty} \left[\left(1 - \frac{z}{z_n} \right) e^{P_k(z/z_n)} \right] = e^{h(z)} E(z)$$

The proof of this important theorem requires a number of intermediate results, notably the *minimum modulus theorem* proved in the following section, a very useful result in its own right.

Lemma 39.2. *Let ε be such that $\lambda := \rho + \varepsilon < k_\rho := k$. There is a $c > 0$ such that*

$$(39.3) \quad |(1 - \zeta) \exp P_k(\zeta)| \leq \exp(c|\zeta|^\lambda)$$

Proof. For $|\zeta| \leq 1/2$ we have

$$(39.4) \quad \ln(1 - \zeta) + P_k(\zeta) = \sum_{n=k}^{\infty} \frac{\zeta^n}{n} = \zeta^k C_k; \quad |C_k| \leq \sum_{n=k}^{\infty} 2^{-n} \leq 2$$

$$(39.5) \quad \Rightarrow (1 - \zeta) e^{P_k(\zeta)} \leq e^{2|\zeta|^k} \leq e^{2|\zeta|^\lambda}$$

For $|\zeta| \in [1/2, 1]$ we have

$$(39.6) \quad |(1 - \zeta) \exp P_k(\zeta)| \leq \frac{1}{2} \exp \left[|\zeta|^k \left(\frac{1}{|\zeta|^{k-1}} + \cdots + \frac{1}{|\zeta|^{k-1}} \right) \right] \\ \leq \frac{1}{2} \exp(2^k |\zeta|^k) \leq \frac{1}{2} \exp(2^k |\zeta|^\lambda)$$

For $|\zeta| > 1$ we have

$$(39.7) \quad |(1 - \zeta) \exp P_k(\zeta)| \leq |1 - \zeta| \exp \left[|\zeta|^{k-1} \left(\frac{1}{k-1} + \cdots + \frac{1}{|\zeta|^{k-2}} \right) \right] \\ \leq \exp(k|\zeta|^{k-1} + \ln|1 + |\zeta||) \leq \exp(k|\zeta|^\lambda + \ln|1 + |\zeta||) \leq \exp(C_2 |\zeta|^\lambda)$$

for some C_2 independent of ζ , $|\zeta| > 1$. This is because $t^{-\lambda} \ln(1 + t)$ is continuous on $[1, \infty)$ and goes to zero at infinity (fill in the details). $\blacksquare$

39.1. Canonical products. Take any sequence $\{z_n\}_n$ where the terms are ordered by absolute value, with the property that for some $\rho > 0$ and any $\varepsilon > 0$ we have

$$(39.8) \quad \sum_{n=1}^{\infty} \frac{1}{|z_n|^{\rho+\varepsilon}} < \infty$$

Definition 39.3. The canonical product determined by the sequence $\{z_n\}$, denoted by $E^{(k)}(z, \{z_n\})$ or simply $E(z)$ is defined by

$$(39.9) \quad E(z) = \prod_{n=1}^{\infty} \left(1 - \frac{z}{z_n} \right) \exp[P_k(z/z_n)]$$

Theorem 39.4. $E(z)$ is an entire function of order $\leq \rho$.

Proof. Take again any ε be such that $\lambda := \rho + \varepsilon < k_\rho$. Then, by Lemma 39.2 we have

$$(39.10) \quad |E(z)| \leq \prod_{n=1}^{\infty} \exp(c|z/z_n|^\lambda) = \exp \left(c|z|^\lambda \sum_{k=1}^{\infty} |z_n|^{-\lambda} \right) \leq \exp(c_1 |z|^\lambda)$$

proving, in the process, uniform convergence of the product. $\blacksquare$

40. THE MINIMUM MODULUS THEOREM; END OF PROOF OF
THEOREM 39.1

This important theorem tells us, roughly, that if a function does not grow too fast it cannot decrease too quickly either, aside from zeros. More precisely we have

Theorem 40.1 (Minimum modulus theorem). *Let f be an entire function of order $\leq \rho$. As before, let $\{z_n\}$ be its zeros with $|z_i| > 1$, repeated according to their multiplicity and let $\varepsilon > 0$. At every root, remove a disk $\mathbb{D}_{r_n}(z_n)$ with $r_n = |z_n|^{-\rho-\varepsilon}$. $U = \mathbb{C} \setminus \bigcup_n \mathbb{D}_{r_n}(z_n)$. Then, there is a constant c such that for all $z \in U$ we have*

$$(40.1) \quad |f(z)| \geq \exp(-c|z|^{\rho+\varepsilon}) \quad \text{or} \quad \frac{1}{|f(z)|} = O(\exp(|z|^{\rho+\varepsilon}))$$

Proof. We start with the case when the entire function is a canonical product. We take $|z| = r$ and write

$$(40.2) \quad E(z) = \prod_{|z_n| < 2r} E_k(z, z_n) \prod_{|z_n| \geq 2r} E_k(z, z_n)$$

and estimate the two terms separately. We note that in the second product, all ratios $\zeta =: \zeta_n = z/z_n$ have the property $|\zeta| \leq 1/2$. Taking one term of the product, we have to estimate *below*

$$(40.3) \quad E(\zeta) = (1 - \zeta)e^{P(\zeta)}$$

Since $|\zeta| \leq 1/2$, $\ln(1 - \zeta)$ exists; we take the principal branch and write

$$(40.4) \quad \begin{aligned} |(1 - \zeta)e^{P(\zeta)}| &= |e^{\ln(1-\zeta)+P(\zeta)}| = \left| \exp \left(- \sum_{n=k}^{\infty} \frac{\zeta^n}{n} \right) \right| \\ &= \exp \left(- \operatorname{Re} \sum_{n=k}^{\infty} \frac{\zeta^n}{n} \right) \geq \exp \left(- \left| \sum_{n=k}^{\infty} \frac{\zeta^n}{n} \right| \right) \geq \exp \left(- \sum_{n=k}^{\infty} \frac{|\zeta|^n}{n} \right) \\ &\geq \exp \left(- |\zeta|^k \sum_{n=0}^{\infty} \frac{(1/2)^n}{n+k} \right) \geq e^{-2|\zeta|^k} \geq e^{-2|\zeta|^{\rho+\varepsilon}} \end{aligned}$$

Thus for the second product in (40.2) we have

$$(40.5) \quad \prod_{|z_n| \geq 2r} E_k(z, z_n) \geq \exp \left(-2|z|^{\rho+\varepsilon} \sum_{|z_n| \geq 2r} \frac{1}{|z_n|^{\rho+\varepsilon}} \right) \geq e^{-c|z|^{\rho+\varepsilon}}$$

since the infinite sum converges by Theorem 38.15.

We split the remaining region into $z_k \in [1, r]$ and $z_k \in (r, 2r)$. Here the factors $(1 - z/z_k)$ have to be bounded from below, and it is here that we use the conditions on the removed disks. We have, on $[1, r]$,

$$(40.6) \quad \prod_{|z_n| \leq r} |1 - z/z_n| = \frac{|z - z_n|}{|z_n|} \geq \prod_{|z_n| \leq r} |z_n|^{-\rho-\varepsilon-1} \geq \prod_{|z_n| \leq r} r^{-\rho-\varepsilon-1} = (r^{-\rho-\varepsilon-1})^{\nu(r)} \gtrsim e^{-r^{\rho+\varepsilon} \ln r(\rho+\varepsilon+1)} \geq e^{-c_1 r^{\rho+\varepsilon'}}$$

for some C_1 and $\varepsilon' > \varepsilon$, since $(r^{\varepsilon-\varepsilon'} \ln r \rightarrow 0$ as $r \rightarrow \infty$.

On $(r, 2r)$ we have

$$(40.7) \quad |1 - z/z_n| = \frac{|z - z_n|}{|z_n|} \geq |z_n|^{-\rho-\varepsilon-1} \geq (2r)^{-\rho-\varepsilon-1}$$

and thus

$$(40.8) \quad \prod_{|z_n| < 2r} |1 - z/z_n| \geq [(2r)^{-\rho-\varepsilon-1}]^{\nu(2r)} = e^{-\nu(2r)(\rho+\varepsilon+1) \ln(2r)} \geq e^{-c_6 r^{\rho+\varepsilon'}}$$

for any $\varepsilon' > \varepsilon$ if r is large enough.

We now examine the convergence improving factors, for $|z_n| < 2r$.

$$(40.9) \quad \left| \sum_{|z_n| < 2r} P_k(z/z_n) \right| \leq \left| \sum_{r < |z_n| < 2r} P_k(z/z_n) \right| + \left| \sum_{|z_n| \leq r} P_k(z/z_n) \right|$$

For the first term on the right we note that when $|z/z_n| = |\zeta_n| =: |\zeta| < 1$ and we have

$$(40.10) \quad \left| \sum_{n=1}^{k-1} \frac{\zeta^n}{n} \right| \leq \sum_{n=1}^{k-1} \frac{1}{n} =: c_1$$

and thus

$$(40.11) \quad \sum_{r < |z_n| < 2r} |P_k(z/z_n)| \leq \nu(2r) c_1 \leq c_2 r^{\rho+\varepsilon}$$

For the second term on the right of (40.9) we note that $|z/z_n| \geq 1$ and thus, with $\zeta = z/z_n$ we have

$$(40.12) \quad \left| \sum_{n=1}^{k-1} \frac{\zeta^n}{n} \right| \leq |\zeta|^{k-1} \sum_{n=1}^{k-1} \frac{1}{n} =: c_1 |\zeta|^{k-1} = c_1 r^{k-1} |z_n|^{-k+1}$$

We use Abel summation by parts (we are careful that r is not necessarily an integer)

$$\begin{aligned}
(40.13) \quad \sum_{|z_n| \leq r} |z_n|^{-k+1} &\leq \sum_{m \leq r} \frac{\nu(m+1) - \nu(m)}{m^{k-1}} \\
&= \sum_{m \leq r} \nu(m+1) \left(\frac{1}{m^{k-1}} - \frac{1}{(m+1)^{k-1}} \right) + \frac{\nu(r+1)}{r^{k-1}} - \nu(1) \\
&\leq \sum_{m \leq r} \nu(m+1) \left(\frac{1}{m^{k-1}} - \frac{1}{(m+1)^{k-1}} \right) + \frac{\nu(r+1)}{r^{k-1}} \\
&\leq \sum_{m \leq r} \frac{kCm^{\rho+\varepsilon}}{m^k} + c_3 r^{\rho+\varepsilon-k+1} \\
&\leq C_1 \sum_{m \leq r} m^{\rho+\varepsilon-k} + c_3 r^{\rho+\varepsilon-k+1} \leq C_3 r^{\rho+\varepsilon-k+1}
\end{aligned}$$

where we majorized the sum by an integral in the usual way. Multiplying by $c_1 r^{k-1}$ we get that the second term on the right of (40.9) is bounded by

$$(40.14) \quad C_4 r^{\rho+\varepsilon}$$

We now finish the proof of Theorem 39.1.

Proof. We take $\varepsilon > 0$ and $s = \rho + \varepsilon$. We order the roots nondecreasingly by $|z_n|$. For each root z_n we consider the annulus $A_n = \{z : |z| \in [|z_n| - 2|z_n|^{-s}, |z_n| + 2|z_n|^{-s}]\}$ (for the purpose of this argument, we can as well assume that all roots are on $\mathbb{R}^+$, since the angular position is irrelevant). Consider $J^c := \mathbb{R}^+ \setminus J$ where J is the union of all intersections of the A_n with $\mathbb{R}^+$. Since the Lebesgue measure of J does not exceed $4 \sum_n |z_n|^{-s} < \infty$, there exist arbitrarily large numbers in the complement J^c . Take r be any number in J^c and consider the circle $\partial \mathbb{D}_r$. Consider the function $g = f/E$. g is clearly an entire function with no zeros. Then, by Theorem 38.6, $g = e^h$ with h entire. Since $\operatorname{Re} h \leq (C_f + C_E)r^{\rho+\varepsilon}$ for some $C_f + C_E$ independent of r in $\mathbb{D}_r$ for arbitrarily large r (check), we have by Corollary 38.17 that h is a polynomial of degree at most $\rho + \varepsilon$. Since ε is arbitrary, h is a polynomial of degree at most ρ . ■

To finish the proof of the minimum modulus principle, we use Hadamard's theorem and the fact that e^{-h} satisfies the required bounds. (Exercise: fill in the details.) ■

Example 40.2. Let us show that $f(z) = e^z - z$ has infinitely many roots in $\mathbb{C}$. Indeed, first note that $f(z)$ has order 1 since $|z| \leq e^{|z|}$ for all z . Suppose f had finitely many zeros. Then

$$(40.15) \quad e^z - z = P(z)e^{h(z)}$$

where $P(z)$ is a polynomial and $h(z)$ is a polynomial of degree one, and without loss of generality we can take $h(z) = cz$, $c = \alpha + i\beta$. As $z = t \rightarrow +\infty$ we have

$$(40.16) \quad P(t)e^{(c-1)t} = 1 - te^{-t} \rightarrow 1$$

In particular $|P(t)|e^{(\alpha-1)t} \rightarrow 1$ which is only possible if $\alpha = 1$. But then $|P(t)| \rightarrow 1$ which is only possible if $P(t) = \text{const} = e^{i\phi}$. We are then left with

$$(40.17) \quad e^{i(\phi+t\beta)} = \cos(t\beta + \phi) + i \sin(t\beta + \phi) \rightarrow 1 \quad \text{as } t \rightarrow +\infty$$

which clearly implies $\beta = 0$. Then $e^{i\phi} = 1$. We are left with the identity

$$(40.18) \quad e^z - z = e^z \quad \forall z$$

which is obviously false.

Exercise 40.3. * Let $P \not\equiv 0$ be a polynomial. Show that the equation $e^z = P(z)$ has infinitely many roots in $\mathbb{C}$.

Exercise 40.4. ** (i) Rederive formula (38.12) using Hadamard's theorem.

(ii) Write down a product formula of the function

$$f(z) = \sin z + 3 \sin(3z) + 5 \sin(5z) + 7 \sin(7z)$$

The final formula should be explicit except for arcsins of roots of a cubic polynomial.

40.1. Some applications.

Corollary 40.5 (Borel). Assume that ρ is not an integer and f has strict order ρ . Then f takes every value in $\mathbb{C}$ infinitely many times.

Proof. It suffices to show that such a function has infinitely many zeros, since f and $f - z_0$ have the same strict order. Assume to get a contradiction f had finitely many zeros. Then $g(z) = f(z) \prod_{i=1}^n (z - z_i)^{-1}$ would be entire, with no zeros, and as it is easy to check, of strict order ρ . Then $g = e^h$ with h a polynomial whose degree can only be an integer. ■

Definition 40.6. Let $\exp_n$ be the exponential composed with itself n times.

Corollary 40.7 (A weak form of Picard's theorem). *A nonconstant entire function which is bounded by $\exp_n(C|z|)$ for some n and large z takes every value with at most one exception.*

Proof. We prove this by induction on n . We first show that a non-constant entire function of finite order takes every value with at most one exception. Assume a is an exceptional (*lacunary*) value. Then $f(z) - a$ is entire with no zeros, thus of the form e^h with h a polynomial, $f = e^h - a$. If the degree of h is zero, then f is a constant. Otherwise, we must show that $e^h - a$ takes all values with at most one exception ($-a$ of course), or, which is the same, e^h takes all values with at most one exception. The equation $e^h = b$, $b \neq 0$ is solved if $h - \ln b$ has roots, which is true by the fundamental theorem of algebra.

Assume now the property holds for $n \leq k-1$ and we wish to prove it for $n = k$. Let f be an entire function bounded by $\exp^{(n)}(C|z|)$ which avoids the value a . Then $f - a$ is entire with no zeros, $f - a = e^h$ with h entire. It is easy to show that h is bounded by $\exp^{(n-1)}(C|z|)$. Thus it avoids at most one value, by the induction hypothesis. The equation $e^h - a = b$, for $b \neq -a$ always has a solution. Indeed, if $\ln(b + a)$ is not an avoided value of h this is obvious. On the other hand, if $\ln(b + a)$ is avoided by h , then again by the induction hypothesis $\ln(b + a) + 2\pi i$ is not avoided. ■

Exercise 40.8. ** Show that the equation

$$(40.19) \quad \cos(z) = z^4 + 5z^2 + 13$$

has infinitely many roots in $\mathbb{C}$.

Exercise 40.9. ** (*Bonus*) Show that the error function

$$(40.20) \quad \operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$$

takes every complex value infinitely many times. (Hint: using L'Hospital show that $\operatorname{erf}(is)/(e^{s^2}/s) \rightarrow \text{const.}$ as $s \rightarrow +\infty$.)

We will return to the error function later and use asymptotic methods to locate, for large x , these special points.

41. THE PHRAGMÉN-LINDELÖF THEOREM

Theorem 41.1 (Phragmén-Lindelöf). *Let U be the open sector between two rays from the origin, forming an angle π/β , $\beta > \frac{1}{2}$. Assume f is analytic in U , and continuous in $\bar{U}$, and for some $C_1, C_2, M > 0$ and $\alpha \in (0, \beta)$ it satisfies the estimates*

$$(41.1) \quad |f(z)| \leq C_1 e^{C_2|z|^\alpha}; \quad z \in U; \quad |f(z)| \leq M; \quad z \in \partial U$$

Then

$$(41.2) \quad |f(z)| \leq M; \quad z \in U$$

Proof. By a rotation we can make $U = \{z : 2|\arg z| < \pi/\beta\}$. Making a cut in the complement of U we can define an analytic branch of the log in U and, with it, an analytic branch of z^β . By taking $g = f(z^{1/\beta})$, we can assume without loss of generality that $\beta = 1$ and $\alpha \in (0, 1)$ and then $U = \{z : |\arg z| < \pi/2\}$. Let $\alpha' \in (\alpha, 1)$, $\varepsilon > 0$ and consider the analytic function

$$(41.3) \quad h(z) = e^{-\varepsilon z^{\alpha'}} f(z)$$

Since $|e^{-\varepsilon z^{\alpha'}}| < 1$ in U (check) and $|e^{-\varepsilon z^{\alpha'} + C_2 z^\alpha}| \rightarrow 0$ as $|z| \rightarrow \infty$ on the half circle $|z| = R$, $\operatorname{Re} z \geq 0$ (check), the usual maximum modulus principle shows that $|h| < M$ in $\overline{U}$. The proof is completed by taking $\varepsilon \rightarrow 0$. ■

41.1. An application to Laplace transforms. We will study Laplace and inverse Laplace transforms in more detail later. For now let $F \in L^1(\mathbb{R})$. Then by Fubini and dominated convergence, the Laplace transform

$$(41.4) \quad \mathcal{L}F := \int_0^\infty e^{-px} F(p) dp$$

is well defined and continuous in x in the closed $\mathbb{H}^+$ and analytic in the open RHP. (Obviously, we could allow $F e^{-|\alpha|p} \in L^1$ and then $\mathcal{L}F$ would be defined for $\operatorname{Re} x > |\alpha|$.) F is uniquely defined by its Laplace transform, as seen below.

Lemma 41.2 (Uniqueness). *Assume $F \in L^1(\mathbb{R}^+)$ and $\mathcal{L}F = 0$ for a set of x with an accumulation point in $\mathbb{H}^+$. Then $F = 0$ a.e. on $[0, \infty)$.*

Proof. By analyticity, $\mathcal{L}F = 0$ in the open RHP and by continuity, for $s \in \mathbb{R}$, $\mathcal{L}F(is) = 0 = \hat{\mathcal{F}}F$ where $\hat{\mathcal{F}}F$ is the Fourier transform of F (extended by zero for negative values of p). Since $F \in L^1$ and $0 = \hat{\mathcal{F}}F \in L^1$, by the known Fourier inversion formula [8], $F = 0$ a.e. ■

More however can be said. We can draw interesting conclusions about F just from the rate of decay of $\mathcal{L}F$.

Proposition 41.3 (Lower bound on decay rates of Laplace transforms). *Assume $F \in L^1(\mathbb{R}^+)$ and for some $\varepsilon > 0$ we have*

$$(41.5) \quad \mathcal{L}F(x) = O(e^{-\varepsilon x}) \quad \text{as } x \rightarrow +\infty$$

Then $F = 0$ a.e. on $[0, \varepsilon]$.

Proof. We write

$$(41.6) \quad \int_0^\infty e^{-px} F(p) dp = \int_0^\varepsilon e^{-px} F(p) dp + \int_\varepsilon^\infty e^{-px} F(p) dp$$

we note that

$$(41.7) \quad \left| \int_\varepsilon^\infty e^{-px} F(p) dp \right| \leq e^{-\varepsilon x} \int_\varepsilon^\infty |F(p)| dp \leq e^{-\varepsilon x} \|F\|_1 = O(e^{-\varepsilon x})$$

Therefore

$$(41.8) \quad g(x) = \int_0^\varepsilon e^{-px} F(p) dp = O(e^{-\varepsilon x}) \quad \text{as } x \rightarrow +\infty$$

The function g is entire. Let $h(x) = e^{\varepsilon x} g(x)$. Then by assumption h is entire and uniformly bounded for $x \in \mathbb{R}$ (since by assumption, for some x_0 and all $x > x_0$ we have $h \leq C$ and by continuity $\max |h| < \infty$ on $[0, x_0]$). The function is also manifestly bounded for $x \in i\mathbb{R}$ (by $\|F\|_1$). By Phragmén-Lindelöf (first applied in the first quadrant and then in the fourth quadrant, with $\beta = 2, \alpha = 1$) h is bounded in the closed RHP. Now, for $x = -s < 0$ we have

$$(41.9) \quad e^{-s\varepsilon} \int_0^\varepsilon e^{sp} F(p) dp \leq \int_0^\varepsilon |F(p)| dp \leq \|F\|_1$$

Again by Phragmén-Lindelöf (again applied twice) h is bounded in the closed $\mathbb{H}_l$ thus bounded in $\mathbb{C}$, and it is therefore a constant. But, by the Riemann-Lebesgue lemma, $h \rightarrow 0$ for $x = is$ when $s \rightarrow +\infty$. Thus $h \equiv 0$. But then, with χ_A the characteristic function of A ,

$$(41.10) \quad \int_0^\varepsilon F(p) e^{-isp} dp = \hat{\mathcal{F}}(\chi_{[0, \varepsilon]} F) = 0$$

for all $s \in \mathbb{R}$ entailing the conclusion. $\blacksquare$

Corollary 41.4. *Assume $F \in L^1$ and $\mathcal{L}F = O(e^{-Ax})$ as $x \rightarrow +\infty$ for all $A > 0$. Then $F = 0$.*

Proof. This is straightforward. $\blacksquare$

As we see, uniqueness of the Laplace transform can be reduced to estimates. Also, no two different $L^1(\mathbb{R}^+)$ functions, real-analytic on $(0, \infty)$, can have Laplace transforms within exponentially small corrections of each-other. This will play an important role later on.

41.2. A Laplace inversion formula.

Theorem 41.5. Assume $c \geq 0$, $f(z)$ is analytic in the closed half plane $H_c := \{z : \operatorname{Re} z \geq c\}$. Assume further that $\sup_{c' \geq c} |f(c' + it)| \leq g(t)$ with $g(t) \in L^1(\mathbb{R})$. Let

$$(41.11) \quad F(p) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} f(x) dx =: (\mathcal{L}^{-1}F)(p)$$

Then for any $x \in \{z : \operatorname{Re} z > c\}$ we have

$$(41.12) \quad \mathcal{L}F = \int_0^\infty e^{-px} F(p) dp = f(x)$$

Proof. Note that for any $x' = x'_1 + iy'_1 \in \{z : \operatorname{Re} z > c\}$

$$(41.13) \quad \int_0^\infty dp \int_{c-i\infty}^{c+i\infty} \left| e^{p(s-x')} f(s) \right| |d|s| \leq \int_0^\infty dp e^{p(c-x'_1)} \|g\|_1 \leq \frac{\|g\|_1}{x'_1 - c}$$

and thus, by Fubini we can interchange the orders of integration:

$$(41.14) \quad \begin{aligned} U(x') &= \int_0^\infty e^{-px'} \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{px} f(x) dx \\ &= \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} dx f(x) \int_0^\infty dp e^{-px' + px} = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \frac{f(x)}{x' - x} dx \end{aligned}$$

Since $g \in L^1$ there must exist subsequences $\tau_n, -\tau'_n$ tending to ∞ such that $|g(\tau_n)| \rightarrow 0$. Let $x' > \operatorname{Re} x = x_1$ and consider the box $B_n = \{z : \operatorname{Re} z \in [x_1, x'], \operatorname{Im} z \in [-\tau'_n, \tau_n]\}$ with positive orientation. We have

$$(41.15) \quad \int_{B_n} \frac{f(s)}{x' - s} ds = -f(x')$$

while, by construction,

$$(41.16) \quad \lim_{n \rightarrow \infty} \int_{B_n} \frac{f(s)}{x' - s} ds = \int_{x'-i\infty}^{x'+i\infty} \frac{f(s)}{x' - s} ds - \int_{c-i\infty}^{c+i\infty} \frac{f(s)}{x' - s} ds$$

On the other hand, by dominated convergence, we have

$$(41.17) \quad \int_{x'-i\infty}^{x'+i\infty} \frac{f(s)}{x' - s} ds \rightarrow 0 \quad \text{as } x' \rightarrow \infty$$

■

41.3. Abstract Stokes phenomena. This theorem shows that if an analytic function decays rapidly along some direction, then it increases “correspondingly” rapidly along a complementary direction. The following is reminiscent of a theorem by Carlson [13].

Theorem 41.6. *Assume $f \not\equiv 0$ is analytic in the closed $\mathbb{H}^+$ and that for all $a > 0$ we have $f(t) = O(e^{-at})$ for $t \in \mathbb{R}^+, t \rightarrow \infty$. Then, for all $b > 0$ the function*

$$(41.18) \quad e^{-bz}f(z)$$

is unbounded in the closed $\mathbb{H}^+$.

Proof. Assume that for some $b > 0$ we had $|e^{-bz}f(z)| < M$ in the closed RHP. Then, the function

$$(41.19) \quad \psi(z) = \frac{e^{-bz}f(z)}{(z+1)^2}$$

satisfies the assumptions of Theorem 41.5. But then $\psi(z) = \mathcal{L}\mathcal{L}^{-1}\psi(z)$ satisfies the assumptions of Corollary (41.4) and $\psi \equiv 0$. ■

Let $\alpha > 2$.

Corollary 41.7. *Assume $f \not\equiv 0$ is analytic in the closed sector $S = \{z : 2|\arg z| \leq \pi/\alpha\}$, $\alpha > \frac{1}{2}$ and that $f(t) \leq Ce^{-t^\beta}$ with $\beta > \alpha$ for $t \in \mathbb{R}^+$. Then for any $\beta' < \beta$ there exists a subsequence $z_n \in S$ such that*

$$(41.20) \quad \left| f(z_n)e^{-z_n^{\beta'}} \right| \rightarrow \infty \quad \text{as } n \rightarrow \infty$$

Proof. This follows from Theorem 41.6 by simple changes of variables. ■

Exercise 41.8. * Carry out the details of the preceding proof.

42. ELLIPTIC FUNCTIONS

This material is based on [3] and [12], with some simplifications and additions.

Elliptic functions are doubly periodic function, and play an important role in analysis, algebra, and number theory. We have already encountered an example of an elliptic function, sn , in §34.1.

42.1. The period module. We have already established (see §36.1) that, for a function f to have two distinct periods ω_1 and ω_2 , we must have $\omega_1/\omega_2 \notin \mathbb{R}$. If ω_1 and ω_2 are periods of f , so is, as you can check, any element of $\Lambda = \Lambda_{\omega_1, \omega_2}$, and this lattice is called *the period module* of f .

We will only analyze doubly periodic functions which are *meromorphic*.

42.2. General properties of elliptic functions. In general, we will not necessarily assume that Λ contains all the periods of f . The equivalence relation $z_1 = z_2 \pmod{\Lambda}$ means, naturally, that $z_1 - z_2 \in \Lambda$. Since the values of f only depend on congruence classes, we can regard it as a function defined on these congruence classes. One way to do that is to restrict f to a parallelogram P_a with vertices $a, a + \omega_1, a + \omega_2, a + \omega_1 + \omega_2$ for some a . To represent all the values of f , we need to include part of the boundary of the parallelogram in P_a , for instance, two adjacent sides. The exact choice of a is immaterial, but it is convenient to choose an a such that f has no poles on the boundary.

Theorem 42.9. *An elliptic function without poles is a constant.*

Proof. Such a function is entire and, by double periodicity, bounded.

■

By definition, poles of an analytic function are isolated, and thus there are only finitely many in a fundamental parallelogram. When we count poles or zeros of elliptic functions, we only count those in a fundamental parallelogram.

Theorem 42.10. *The sum of the residues of an elliptic function is zero.*

Proof. We choose as discussed a fundamental parallelogram P_a such that no poles lie on ∂P_a . The sum of the residues is given by

$$(42.21) \quad \frac{1}{2\pi i} \int_{\partial P_a} f(s) ds$$

This integral is zero since the integrals on opposite sides cancel each other.

■

Theorem 42.11. *If f is a non-constant elliptic function, then the number of poles equals the number of zeros of f .*

Proof. The function $g = f'/f$ is also meromorphic and doubly periodic, with poles of residue -1 for each pole of f and $+1$ for each zero. Now Theorem 42.10 follows from Theorem 42.10.

■

Note 42.12. *The three theorems above are due to Liouville.*

Corollary 42.13. *Any value of f is assumed equally many times.*

Proof. If $c \in \mathbb{C}$, then $f - c$ is elliptic, and has the same poles as f . ■

Theorem 42.14. *Let $a_1, \dots, a_n$ be the zeros of the elliptic function f and $b_1, \dots, b_n$ its poles in the period parallelogram. Then $\sum_{i=1}^n a_i = \sum_{i=1}^n b_i \pmod{\Lambda}$.*

Proof. Take a parallelogram as in Theorem 42.10 and note that

$$(42.22) \quad \sum_{i=1}^n a_i - \sum_{i=1}^n b_i = \frac{1}{2\pi i} \int_{\partial P_a} \frac{sf'(s)}{f(s)} ds$$

Note also that the values of s , and thus of the integrand, on opposite sides of the parallelogram differ by the corresponding period, ω_i of f , yielding

$$(42.23) \quad \frac{1}{2\pi i} \left(\int_a^{a+\omega_1} + \int_{a+\omega_1+\omega_2}^{a+\omega_2} \right) \frac{sf'(s)}{f(s)} ds = -\frac{\omega_2}{2\pi i} \int_a^{a+\omega_1} \frac{f'(s)}{f(s)} ds = -\frac{\omega_2}{2\pi i} N_1$$

where N_1 , an integer, is the winding number of 0 w.r.t. the *closed* curve $f([a, a + \omega_1])$. Repeating with the other sides, we see that

$$(42.24) \quad \sum_{i=1}^n a_i - \sum_{i=1}^n b_i = n_1 \omega_1 + n_2 \omega_2, \quad \text{for some } n_1, n_2 \in \mathbb{Z}$$

■

42.3. The Weierstrass elliptic functions. The starting point of the Weierstrass theory are elliptic functions with one double pole per parallelogram, that we convene to place at the origin. Since multiplication by a constant does not change the theory of such functions, we choose the singular part to be just z^{-2} . The function f must be even. Indeed, $f(z) - f(-z)$ is elliptic without poles, thus a constant, and since $f(\omega_1/2) = f(-\omega_1/2)$ the constant is zero. Since adding a constant to f is also irrelevant, we choose it so that the constant term in the Laurent expansion is zero. With this choice, we have $f = \wp$, the Weierstrass elliptic function:

$$(42.25) \quad \wp(z) = z^{-2} + \sum_{k \geq 1} a_k z^{2k}$$

However, all of the above is contingent on the existence of such a function which we must prove:

Theorem 42.15. *The function*

$$(42.26) \quad \wp(z) = \frac{1}{z^2} + \sum_{0 \neq \omega \in \Lambda_{\omega_1, \omega_2}} \left(\frac{1}{(z - \omega)^2} - \frac{1}{\omega^2} \right)$$

is doubly periodic and has a Laurent expansion of the form (42.25) at the origin.

Proof. Until we finish the proof, we write $f(z)$ for the sum on the right side of (42.26). Let us first check convergence of the sum on compact sets away from Λ . We note that $|n_1\omega_1 + n_2\omega_2| = |\omega_1||n_1 + n_2\tau|$. Since $\tau \notin \mathbb{R}$, for fixed z and large $|n_1| + |n_2|$ we have, for some constants C', C

$$(42.27) \quad \left| \frac{1}{(z - \omega)^2} - \frac{1}{\omega^2} \right| \leq \frac{C'|z|}{|\omega|^3} \leq \frac{C|z|}{(n_1^2 + n_2^2)^{3/2}}$$

Uniform and absolute convergence now follows from the integral test since

$$\int_{r \geq R} \int_0^{2\pi} \frac{1}{r^3} r dr d\phi$$

converges. Next we check periodicity. Due to the correction terms $-\omega^{-2}$ needed to ensure convergence, it is tedious to check this directly. Instead, we can differentiate term by term in (42.26) and get

$$(42.28) \quad f'(z) = -2 \sum_{\omega \in \Lambda} \frac{1}{(z - \omega)^3}$$

Clearly f' is doubly periodic; it follows that $f(z + \omega_1) - f(z) = c_1$ and $f(z + \omega_2) - f(z) = c_2$ for some constants c_1 and c_2 . Recall that f is even. Because of that, choosing $z_1 = -\omega_1/2$ and $z_2 = -\omega_2/2$ we see that $c_1 = c_2 = 0$. It is easy to check that the Laurent expansion at 0 of f is of the form (42.25). ■

42.3.1. The Weierstrass zeta function. Since $\wp$ has zero residues, its antiderivative is also a meromorphic function. We choose the antiderivative $-\zeta(z)$ which is *odd*. Explicitly,

$$(42.29) \quad \zeta(z) = \frac{1}{z} + \sum_{\omega \neq 0} \left(\frac{1}{z - \omega} + \frac{1}{\omega} + \frac{z}{\omega^2} \right)$$

The Maclaurin series in z of the term in parenthesis, equal to the expansion in ω^{-1} as $\omega \rightarrow \infty$, is

$$(42.30) \quad \frac{1}{z - \omega} + \frac{1}{\omega} + \frac{z}{\omega^2} = -\omega^{-1} \sum_{k \geq 2} (z/\omega)^k$$

from which convergence and parity follow. Summing over periods we get

$$(42.31) \quad \zeta(z) = \frac{1}{z} - \sum_{k \geq 2} G_k z^{2k-1}$$

where

$$(42.32) \quad G_k = \sum_{\omega \neq 0} \frac{1}{\omega^{2k}}$$

Since $\wp$ is doubly periodic, we must have $\zeta(z + \omega_i) - \zeta(z) = \eta_i$, $i = 1, 2$, where η_i are constants. Since ζ has one simple pole of residue one per parallelogram, we have

$$(42.33) \quad \frac{1}{2\pi i} \int_{\partial P_a} \zeta(s) ds = 2\pi i$$

By adding the contributions of the opposite sides of the parallelogram, using the relation $\zeta(z + \omega_i) - \zeta(z) = \eta_i$ and comparing with (42.33) we get *Legendre's relation*

$$(42.34) \quad \eta_1 \omega_2 - \eta_2 \omega_1 = 2\pi i$$

42.3.2. *The differential equation of $\wp$.* Using the definition $-\zeta' = \wp$ and (42.31) (or simply the definition of $\wp$) we get

$$(42.35) \quad \wp(z) = \frac{1}{z^2} + \sum_{k=2}^{\infty} (2k-1) G_k z^{2k-2}$$

From here, a calculation shows that the singular part of the combination $f = (\wp')^2 - 4\wp^3 + 60G_2\wp$ vanishes. Since f is a doubly periodic function, it must be a constant, equal to $f(0)$ which the same calculation shows it equals $-140G_3$. Writing $+60G_2 = g_2$ and $140G_3 = g_3$ we get the separable ODE

$$(42.36) \quad \wp'(z)^2 = 4\wp^3 - g_2\wp - g_3$$

with solution

$$(42.37) \quad z - z_0 = \int_{\wp(z_0)}^{\wp(z)} \frac{ds}{\sqrt{4s^3 - g_2s - g_3}}$$

along a path which avoids the zeros of the denominator.

We note that the roots of the cubic are distinct. We can see this in two ways. One is to note that the integral is elementary if two or three roots coincide. If three coincide, we get $\wp(s) = 4(s + C)^{-2} + 1$ whose inverse is not periodic, and if only two coincide, we get an elementary simply periodic function. But there is a more consequential way to show and use that.

42.4. The modular function λ . Let e_1, e_2, e_3 be the roots of $\wp$ and write its differential equation in the form

$$(42.38) \quad \wp'(z)^2 = 4(\wp(z) - e_1)(\wp(z) - e_2)(\wp(z) - e_3)$$

We see that the roots of $\wp$ are zeros of $\wp'$. Since $\wp$ is periodic and even, for any period ω of $\wp$ we have $\wp(\omega - z) = \wp(z)$ which implies $\wp'(\omega - z) = -\wp'(z)$ implying that $\wp'(\omega/2) = 0$. If we take for ω the values $\omega_1, \omega_2, \omega_1 + \omega_2$ we get the three roots of $\wp'$ which lie in the fundamental parallelogram and are distinct,

$$(42.39) \quad e_1 = \wp(\omega_1/2), \quad e_2 = \wp(\omega_2/2), \quad e_3 = \wp(\tfrac{1}{2}(\omega_1 + \omega_2))$$

If, say, $e_1 = e_2$ then we would have $\wp(\omega_1/2) = \wp(\omega_2/2)$; since $\omega_1/2$ is distinct from $\omega_2/2$ and $\wp$ takes each value exactly twice, this is a contradiction (otherwise $(\wp')^2$ cannot be of order 6).

Calculating e_1 from (42.26), we see that the e_i are homogeneous of degree -2 in ω_1, ω_2 , that is, replacing ω_1, ω_2 by $\lambda\omega_1, \lambda\omega_2$ we have $e_i \mapsto \lambda^{-2}e_i$. Based on this, we conclude that the λ function,

$$(42.40) \quad \lambda(\tau) = \frac{e_3 - e_2}{e_1 - e_2}$$

depends only on $\tau = \omega_2/\omega_1$ as indicated.

We next note, again from (42.26) that

$$(42.41) \quad \omega_1^2 e_1 = 4 + \sum_{\omega \neq 0} \left(\frac{1}{(\frac{1}{2} - n_1 - n_2 \tau)^2} - \frac{1}{(n_1 + n_2 \tau)^2} \right)$$

which is analytic for $\text{Im } \tau > 0$ (and also for $\text{Im } \tau < 0$). The same conclusion is reached for e_2, e_3 . Since $e_1 \neq e_2$ and $e_2 \neq e_3$, λ is analytic in the UHP and does not take the values 0 or 1.

42.5. The action of the $\text{PSL}(2, \mathbb{Z})$ on λ . Recall that $\text{PSL}(2, \mathbb{Z})$ is an automorphism of Λ , and since $\wp$ is periodic with any $\omega \in \Lambda$, if we take

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PSL}(2, \mathbb{Z})$$

and change basis to

$$(42.42) \quad \begin{pmatrix} \omega'_2 \\ \omega'_1 \end{pmatrix} = M \begin{pmatrix} \omega_2 \\ \omega_1 \end{pmatrix}$$

the roots of $\wp$ cannot change; they can only be permuted. Recall also that $\text{PSL}(2, \mathbb{Z})$ is generated by $M_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$, $M_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ corresponding to $\tau \mapsto \tau + 1$ and $\tau \mapsto -1/\tau$ resp. Choosing $M = M_1$ in (42.42) has the effect $\frac{1}{2}\omega'_2 = \frac{1}{2}(\omega_1 + \omega_2)$ and $\omega'_1 = \omega_1$. This means e_2

is fixed and e_3 and e_1 are interchanged. Choosing $M = M_2$ we get $\omega'_2/2 = -\omega_1/2$ and $\omega'_1/2 = -\omega_2/2$, in which case e_1 and e_2 are interchanged and e_3 is fixed. It follows that λ satisfies the functional equations

$$(42.43) \quad \lambda(\tau + 1) = \frac{\lambda(\tau)}{\lambda(\tau) - 1}; \quad \lambda(-1/\tau) = 1 - \lambda(\tau)$$

Both transformations on the right side are involutions. It follows that $\lambda(\tau + 2) = \lambda(\tau)$. In turn, this implies that λ is an analytic function of $e^{i\pi\tau}$.

Now, if the matrix $M \in PL(2, \mathbb{Z})$ is of the form

$$(42.44) \quad M := \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \pmod{2}$$

then

$$\lambda\left(\frac{a\tau + b}{c\tau + d}\right) = \lambda(\tau)$$

This is simply because $\omega'_1/2 = \omega_1/2 + \tilde{\omega}_1$ and $\omega'_2/2 = \omega_2/2 + \tilde{\omega}_2$ where $\tilde{\omega}_{1,2} \in \Lambda$. Hence, if $M_2 \equiv M_1 \pmod{2}$, then the modular transformations induced on λ by M_1 and M_2 are identical.

The subgroup of $PSL(2, \mathbb{Z})$ defined by the equivalence mod 2 is called the congruence subgroup mod 2.

Exercise 42.16. Using (42.43), check that

$$(42.45) \quad \lambda\left(\frac{a\tau + b}{c\tau + d}\right) = \begin{cases} \lambda(\tau), & (a, b, c, d) \equiv (1, 0, 0, 1) \pmod{2} \\ 1 - \lambda(\tau), & (a, b, c, d) \equiv (0, 1, 1, 0) \pmod{2} \\ \frac{1}{\lambda(\tau)}, & (a, b, c, d) \equiv (1, 0, 1, 1) \pmod{2} \\ \frac{1}{1 - \lambda(\tau)}, & (a, b, c, d) \equiv (0, 1, 1, 1) \pmod{2} \\ \frac{\lambda(\tau) - 1}{\lambda(\tau)}, & (a, b, c, d) \equiv (1, 1, 1, 0) \pmod{2} \\ \frac{\lambda(\tau)}{\lambda(\tau) - 1}, & (a, b, c, d) \equiv (1, 1, 0, 1) \pmod{2} \end{cases};$$

$$\tau \in \mathbb{H} \text{ and } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$$

We note that the transformations above generate the so-called **anharmonic group**:

$$\left\{ \lambda, 1 - \lambda, \frac{1}{\lambda}, \frac{1}{1 - \lambda}, \frac{\lambda - 1}{\lambda}, \frac{\lambda}{\lambda - 1} \right\}$$

Recall that by the definition of e_1, e_2, e_3 and the double periodicity of $\wp$ any transformation as in (42.45) can only permute the e_i . The table above can also be obtained based on this observation.

42.6. The conformal mapping of λ . We can without loss of generality normalize $\omega_1 = 1, \omega_2 = \tau$. In this case we have

$$(42.46) \quad \begin{aligned} e_3 - e_2 &= \sum_{m,n \in \mathbb{Z}} \left[\frac{1}{(m - \frac{1}{2} + (n + \frac{1}{2})\tau)^2} - \frac{1}{(m + (n - \frac{1}{2})\tau)^2} \right] \\ e_1 - e_2 &= \sum_{m,n \in \mathbb{Z}} \left[\frac{1}{(m - \frac{1}{2} + n\tau)^2} - \frac{1}{(m + (n - \frac{1}{2})\tau)^2} \right] \end{aligned}$$

As discussed, these functions are analytic in the UHP. Note also that the sums are even, implying that *both of them are real on $i\mathbb{R}^+$* . Proposition 37.1 gives

$$\frac{\pi^2}{\sin^2 \pi z} = \sum_{m \in \mathbb{Z}} \frac{1}{(z - m)^2}$$

and we see that

$$(42.47) \quad \begin{aligned} e_3 - e_2 &= \pi^2 \sum_{n \in \mathbb{Z}} \left(\frac{1}{\cos^2 \pi(n - \frac{1}{2})\tau} - \frac{1}{\sin^2 \pi(n - \frac{1}{2})\tau} \right) \\ e_1 - e_2 &= \pi^2 \sum_{n \in \mathbb{Z}} \left(\frac{1}{\cos^2 \pi n\tau} - \frac{1}{\sin^2 \pi(n - \frac{1}{2})\tau} \right) \end{aligned}$$

which converge uniformly in $\{z : \operatorname{Im} z \geq \delta > 0\}$. This implies that λ is real on $i\mathbb{R}^+$.


Theorem 42.17. *The function λ is a conformal map between the domain Ω bounded by the lines $i[0, \infty)$, the circle of radius $\frac{1}{2}$ centered at $\frac{1}{2}$ and the line $1 + i[0, \infty)$ and the UHP.*

Proof. We first follow the mapping of the boundary of Ω and then apply the argument principle. Dominated convergence shows that, as $\operatorname{Im} \tau \rightarrow \infty$, $e_3 - e_2 \rightarrow 0$ and $e_1 - e_2 \rightarrow \pi^2$ (the latter due to the term with $n = 0$). We have $\lambda(\tau) \rightarrow 0$ as $\operatorname{Im} \tau \rightarrow \infty$ uniformly in $\operatorname{Re} \tau$. More precisely, from (42.47) we see that

$$(42.48) \quad e_3 - e_2 = 2\pi^2 \left[\frac{4e^{\pi i\tau}}{(1 + e^{\pi i\tau})^2} + \frac{4e^{\pi i\tau}}{(1 - e^{\pi i\tau})^2} \right] (1 + h(\tau))$$

where $h \rightarrow 0$ as $\operatorname{Im} \tau \rightarrow \infty$. Hence

$$(42.49) \quad \lambda(\tau)e^{-i\pi\tau} \rightarrow 16 \quad \text{as } \operatorname{Im} \tau \rightarrow +\infty$$

FIGURE 45. The domain Ω

From (42.43), $\lambda(-1/\tau) = 1 - \lambda(\tau)$, we see that $\lambda \rightarrow 1$ as $\tau \rightarrow 0$ along $i\mathbb{R}^+$; hence 0 is mapped to 1.

The map $\frac{\tau}{\tau+1} \in \text{Aut}(\mathbb{H}^u)$ and takes $i[0, \infty)$ to the half circle in Fig. 45.

We have $\lambda(\frac{\tau}{\tau+1}) = \frac{1}{\lambda(\tau)}$. This implies that 0 is mapped to 1, as before and, as $\tau \rightarrow i\infty$ we have $\frac{\tau}{\tau+1} \rightarrow 1$, hence 1 is mapped to $+\infty$. Finally, we have

$$(42.50) \quad \lambda(1+it) = \frac{\lambda(it)}{\lambda(it)-1} = -\frac{1}{\lambda(\frac{i}{t})}$$

This means that $1+i0^+$ is mapped to $-\infty$ and $1+i\infty$ is mapped to 0. Hence the boundary of Ω is mapped to $\mathbb{R}$.

Now we use the argument principle and §19.1. By (42.50) λ takes any small value only once. Indeed, we cannot have $\lambda \rightarrow 0$ inside Ω since $\lambda \neq 0$. Near infinity λ is clearly injective, and the only point on $\partial\Omega$ where 0 is taken is at ∞ . Hence, by §19.1 every value in the UHP is taken exactly once. The same reasoning shows that λ takes no value in the LHP: Indeed, small values can only be taken if $\text{Im } \tau \rightarrow +\infty$ but there λ is in the UHP. ■

43. THE UNIFORMIZATION THEOREM

Theorem 43.18. *Every simply connected Riemann surface M is conformally equivalent to one of three Riemann surfaces: $\mathbb{D}$, $\mathbb{C}$, or $\hat{\mathbb{C}}$.*

(see [9]). Recall the hyperbolic plane. Since the UHP is conformally equivalent through a Cayley transform to $\mathbb{D}$, $\mathbb{D}$ also has a metric of constant negative curvature. The conformal map in Theorem 43.18 induces a constant curvature metric on any simply connected Riemann surface M .

If M is compact, then its *universal cover* is $\mathbb{D}$ iff it is a hyperbolic surface of genus greater than 1 with non-abelian fundamental group; its universal cover is $\mathbb{C}$ iff it has genus 1: the complex tori and the elliptic curves with fundamental group $\mathbb{Z}_2$; the universal cover is the Riemann sphere iff M has genus zero, meaning M is the Riemann sphere itself, with trivial fundamental group. Recall also that the only functions analytic on $\hat{\mathbb{C}}$ are rational functions. For a rational function to be injective, it has to be a LFT, and these belong to $\text{Aut}(\hat{\mathbb{C}})$.

If M is not simply connected, we have the following generalization (see [6], with a proof in [9])

Theorem 43.19. *Let M be a non-simply connected Riemann surface. Then, the following is a complete list of isomorphic classes:*

(A) *M is conformally isomorphic to the punctured plane, or compact and of genus 1, and isomorphic to $\mathbb{C}/\Lambda_{1,\tau}$, with $\text{Im } \tau > 0$.*

(B) *M is conformally isomorphic to $\mathbb{D}/G$ where $G \in \text{Aut}(\mathbb{D})$ acts freely and properly discontinuously on $\mathbb{D}$ and apart from the identity no LFT in G has fixed points in $\mathbb{D}$; furthermore, G is isomorphic to $\pi_1(M)$ as groups.*

Exercise 43.20 (Ahlfors). *Show that the Klein j -invariant*

$$j(\tau) = 256 \frac{(1-x)^3}{x^2}$$

where $x = \lambda(1-\lambda)$ is invariant under the modular group.

43.1. An example: M , the universal cover of $\hat{\mathbb{C}} \setminus \{0, 1, \infty\}$. We use the function λ to illustrate the uniformization of this Riemann surface on the upper half plane; of course, composition with a Cayley transform achieves uniformization on $\mathbb{D}$. The fundamental group $\pi_1(M)$ is the free group with two generators, a_0, a_1 . We think of M in terms of equivalence classes of curves over $\mathbb{C} \setminus \{0, 1\}$, winding around 0, 1 in the way prescribed by the associated word, e.g. $a_0 a_1 a_0^{-1} a_1, \dots$




FIGURE 46. Successive reflections

Exercise 43.21. *Show that a Schwarz reflection of a line in a circle is a line or a circle. Show that reflecting Ω successively across boundary arcs as shown in Fig. 47 covers the UHP.*

First we note that each domain obtained by Schwarz reflections as shown in Fig. 47 is mapped by λ conformally, alternatively, to the UHP and LHP. Since the $1 + i\mathbb{R}^+$ is mapped to $(-\infty, 0)$, it means that a reflection of Ω itself across its right boundary is a reflection of the UHP across the segment $(-\infty, 0)$. Likewise, a reflection across the left boundary corresponds to reflecting the UHP across $(0, 1)$ and reflection across the circle is a reflection of the UHP across $(1, \infty)$. Clearly, all these are analytic arcs, and it means that $\psi = \lambda^{-1}$ has analytic continuation through these arcs. Continuing all these reflections, we see that ψ has analytic continuation along any curve in the universal cover, and has singularities only at $\{0, 1, \infty\}$. Hence:

Proposition 43.22. *ψ has analytic continuation on M .*

Furthermore, the word $abc\dots$ corresponds uniquely to a set of successive reflections in the intervals $(-\infty, 0)$, $(0, 1)$ and $(1, \infty)$. Indeed, we see that two different words $abcd, \dots$ and $abcD\dots$ correspond to reflections across distinct arcs in the figure (here the difference is in the fact that d and D correspond to reflections across different arcs). Hence, two different points on M are mapped by ψ to different points in the upper half plane. Using the result in Exercise 43.21 we see that

Proposition 43.23. *ψ a conformal map from M to the UHP.*




FIGURE 47. A winding curve in $\hat{\mathbb{C}} \setminus \{0, 1, \infty\}$ becomes an open curve in the UHP.

Exercise 43.24. Clearly, λ is conformal from the UHP to M . Show that $\mathbb{R}$ is a natural boundary for λ . Use Exercises 42.16 and 36.21 to give a different proof of this fact.

44. THE LITTLE PICARD THEOREM

Theorem 44.25 (Little Picard theorem). *If $f : \mathbb{C} \rightarrow \mathbb{C}$ is entire and there are two points $a \neq b$ which are not in the range of f , then f is a constant.*

Points that are not in the range of f are called *lacunary values*.

Proof. Assume that f is entire and has two lacunary values. Without loss of generality, we can assume that these are 0 and 1. Let $h = C \circ \psi \circ f$ where C is the Cayley transform and ψ is as in Proposition 43.23. Since the only singularities of ψ are at 0 and 1, ψ takes M to the UHP and C is analytic in the UHP, h is entire with values in $\mathbb{D}$, meaning h , hence f are constants. ■

Theorem 44.26 (Montel's fundamental normality test). *Let $\mathcal{D}$ be a domain in $\mathbb{C}$, let α and β be two distinct complex numbers, and let $\mathcal{F}$ be the set of holomorphic functions in $\mathcal{D}$ whose range omits the two values α and β . Then $\mathcal{F}$ is a normal family in $\hat{\mathbb{C}}$.*

A stronger version, using the uniformization theorem is ([6])

Theorem 44.27. *Let M be a Riemann surface and let $\mathcal{F}$ be the set of holomorphic functions on M with values in $\hat{\mathbb{C}}$ whose range omits the three values α , β and γ . Then $\mathcal{F}$ is a normal family in $\hat{\mathbb{C}}$.*

Proof of Theorem 44.26. We first arrange the following: $\alpha = 0, \beta = 1$, and, since normality is a local property, that $\mathcal{D} = \mathbb{D}$.

With $\psi = \lambda^{-1}$, let $\eta = C \circ \psi$ where C is the Cayley transform; the function η is a conformal map from M , the universal cover of $\hat{\mathbb{C}} \setminus \{0, 1, \infty\}$ to $\mathbb{D}$.

Let $f \in \mathcal{F}$ and $a \in \mathbb{D}$ and choose any branch of $\eta(f(a))$. We note, as in the uniformization proof, that for that branch, $\tilde{f} = \eta \circ f$ defined by analytic continuation from a to $\mathbb{D}$ is analytic in $\mathbb{D}$.

To construct a convergent subsequence of a given sequence $\{f_n \in \mathcal{F}\}_{n \in \mathbb{N}}$ we first note that for any $a \in \mathbb{D}$, there is a subsequence of $\{f_n(a)\}$ which converges in $\hat{\mathbb{C}}$, which, by passing to this subsequence, we can assume is $\{f_n(a)\}_{n \in \mathbb{N}}$ itself. If for all a and any subsequence the limit is infinity, then f_n converges to the constant $\infty \in \mathbb{C}$ proving the result. Otherwise, there is a point, say $a = 0$ s.t. a subsequence of $\{f_n(0)\}_{n \in \mathbb{N}}$ converges to $\ell \in \mathbb{C}$. We assume first that $\ell \notin \{0, 1\}$.

Note now that $\tilde{f}_n, n \in \mathbb{N}$ have range in $\overline{\mathbb{D}}$, hence they are a normal family and by Theorem 29.28, they have a convergent subsequence, that by passing again to a subsequence, we can assume is $\{\tilde{f}_n\}_{n \in \mathbb{N}}$; let the limit be g . If g is a constant, then $g = \eta(\ell) \in \mathbb{D}$. Otherwise, since $\text{ran}(g) \subset \overline{\mathbb{D}}$, by the open mapping theorem, $\text{ran}(g) \subset \mathbb{D}$. Now we note that $\eta^{-1} \circ g$ is well-defined, and that $\lim_{n \rightarrow \infty} f_n = \lim_{n \rightarrow \infty} \eta^{-1} \tilde{f}_n = \eta^{-1} \circ g$, which finishes the proof if $\ell \notin \{0, 1\}$.

The remaining case is $\ell = 1$ since, if $\ell = 0$, then we can take the family $1 - f, f \in \mathcal{F}$. Since $f_n \neq 0$, there is a *holomorphic* branch of the square root such that $h_n = \sqrt{f_n} \rightarrow -1$. A holomorphic branch of the square root cannot take both values 1 and -1 , so h_n also avoid 1 and -1 . For h_n the proof above applies to extract a convergent subsequence, which by squaring gives the desired subsequence of f_n .

■

Note 44.28. The fundamental normality test plays an important role in complex dynamics, see [6].

We are now going to prove the celebrated Great Picard's theorem:

Theorem 44.29. *If an analytic function f has an essential singularity at a point z_0 , then on any punctured neighborhood of z_0 , f takes on all possible complex values, with at most a single exception, infinitely often.*

Proof. Without loss of generality, we take $z_0 = 0$, assume that f is analytic in $\mathbb{D}^* = \mathbb{D} \setminus \{0\}$ and assume, to get a contradiction, that f omits the values 0 and 1. Let $\{z_n\}_{n \in \mathbb{N}}$ be any sequence converging to 0 and $f_n(z) = f(z_n z)$. By Theorem 44.26 the family $\{f_n\}_{n \in \mathbb{N}}$ is

normal. Hence we can extract a subsequence $\{f_{n_k}\}_{k \in \mathbb{N}}$ convergent on compact sets in $\mathbb{D}_r \setminus \{0\}$ to, say g which is analytic in $\mathbb{D}_r \setminus \{0\}$ with values in $\hat{\mathbb{C}}$. Assume that g is not the constant $\infty \in \hat{\mathbb{C}}$. Note that, for s in the annulus between the circle of radius $12|z_{n_k}|$ and the circle of radius $\frac{1}{4}|z_n|$, we have $f(s) - g(s') \rightarrow 0$ as $n \rightarrow \infty$ and s' is in the annulus between the circle of radius $\frac{1}{2}$ and the circle of radius $\frac{1}{4}$. Hence f is bounded along any sequence converging to zero, contradiction. If $g = \infty$ then $1/f$ has a removable singularity at 0, thus f has a pole at zero, again a contradiction. ■

45. RIEMANN-HILBERT PROBLEMS: AN INTRODUCTION

An impressive number of problems coming from integrable world can be reduced to so-called Riemann Hilbert problems, and for many of them the only known way to get a closed form solution is via the associated Riemann-Hilbert problem.

Problems which can be solved with R-H techniques include

(1) integrable models such as the transcendental Painlevé equations e.g. $y'' = 6y^2 + x$ (P_I) and many others;

(2) integrable PDEs s.a. nonlinear initial value problem for the KdV (Korteweg–deVries) equation

(45.1)

$$u_t + u_{xxx} + uu_x = 0; \quad u(x, 0) = u_0(x), \quad u \rightarrow 0 \text{ as } |x| \rightarrow \infty \quad (x \in \mathbb{R}, t \in \mathbb{R}^+)$$

(3) inverse scattering problems: find, from the scattering data the potential $q(x)$ in the time-independent Schrödinger equation

$$(45.2) \quad -\psi_{xx} + (k^2 + q(x))\psi = 0$$

(4) questions in orthogonal polynomials, random matrices, and combinatorial probability;

(5) integral equations of the type

$$(45.3) \quad f(t) + \int_0^\infty \alpha(t - t')f(t')dt' = \beta(t)$$

(under suitable integrability conditions);

(6) finding the inverse Radon transform, a transform which is measured in tomography.

45.1. A simple Riemann-Hilbert problem. Perhaps the simplest R-H problem is: given a simple smooth contour $\mathcal{C}$ and $f(t)$ a suitably regular function on $\mathcal{C}$, find analytic functions Φ^+, Φ^- , defined to the left and right of $\mathcal{C}$ such that the limits of $\Phi^\pm$ on $\mathcal{C}$ exist and satisfy

$$(45.4) \quad \Phi^+(z) - \Phi^-(z) = f(z)$$

Note 45.1. Φ is clearly determined up to an entire function. To hope for a unique solution one needs to impose more conditions on Φ , such as behavior at infinity and other special points.

46. CAUCHY TYPE INTEGRALS

We recall that a function is Hölder continuous of order β on a smooth curve $\mathcal{C}$ if

$$(46.1) \quad \exists \beta > 0 \text{ and } C > 0 \text{ s.t. } \forall x, y \in \mathcal{C}, |f(x) - f(y)| \leq C|x - y|^\beta$$

The condition implies continuity if $\beta > 0$ and it is nontrivial if $\beta \leq 1$ (if $\beta > 1$ then $df/ds = 0$).

Let $\mathcal{C}$ for now be a compact smooth curve and ϕ be Hölder continuous on $\mathcal{C}$. Then the function

$$(46.2) \quad \Phi(z) = \frac{1}{2\pi i} \int_{\mathcal{C}} \frac{\phi(s)}{s - z} ds$$

is manifestly analytic for $z \notin \mathcal{C}$ (you can check this by Morera's theorem using Fubini, and in a good number of other ways).

Revisit §10.2 and Definition 30.10.

46.1. Asymptotic behavior of $\Phi(z)$ for large z .

Exercise 46.1. Assume $\mathcal{C}$ is compact. Show that $\Phi(z)$ is analytic at infinity in z and that

$$(46.3) \quad \Phi(z) = - \left(\frac{1}{2\pi i} \int_{\mathcal{C}} \phi(s) ds \right) \frac{1}{z} (1 + g(1/z)) \quad \text{as } z \rightarrow \infty$$

as where $g(1/z) \rightarrow 0$ as $z \rightarrow \infty$.

46.2. Regularity and singularities. Let us first take a simple example, in which ϕ is analytic in a neighborhood $\mathcal{N}$ of $(0, 1)$ (note that this does not exclude multi-valued functions singular at zero and one s.a. $\ln z(1 - z)$).

Then Φ has analytic continuation from $\mathbb{H}^u$ and $\mathbb{H}_l$ through $\mathcal{C} \setminus \{c_i\}$ in $\mathcal{N}$, where c_i are endpoints of $\mathcal{C}$ (if any). (These analytic continuations are in general different.) Indeed, to perform analytic continuation in $z \in \mathbb{H}^u \cup (\mathcal{N} \setminus \{c_i\})$, we can first deform the contour as shown in deform as shown in Fig. 48. To calculate the values of Φ on $\mathcal{C}$, we can use Exercise 10.61 to see that $A(a) = \pm \frac{1}{2} \phi(a) + \frac{1}{2\pi i} PV \int_0^1 \phi(s)(s - t)^{-1} ds$.

The same, interestingly, holds more generally:




FIGURE 48. Analyticity of Φ when ϕ is analytic in a neighborhood of $\mathcal{C}$.

Theorem 46.2 (Plemelj's formulas). *Assume ϕ is Hölder continuous of exponent β on the simple smooth curve $\mathcal{C}$. Let t be an interior point of $\mathcal{C}$. Then,*

(i)

$$(46.4) \quad PV \int \frac{\phi(s)}{s-t} ds$$

(cf. Definition 10.62) exists.

(ii) Let $\{z_n\}_{n \in \mathbb{N}}$ be a sequence approaching t from the left (right). If $\mathcal{C}$ is not bounded, assume also that $\phi \in L^1(\mathcal{C})$. Then, with the $\pm$ sign being $+$ for left limit and $-$ for right limit,

$$(46.5) \quad \lim_{n \rightarrow \infty} \Phi(z_n) = \Phi^\pm(t)$$

where

$$(46.6) \quad \Phi^\pm(t) = \pm \frac{1}{2} \phi(t) + \frac{1}{2\pi i} PV \int \frac{\phi(s)}{s-t} ds$$

and

Note 46.3. (i) The properties in the theorem are local; it is enough to prove them for compact pieces of $\mathcal{C}$.

(ii) A similar statement can be obviously made when t is approached along a curve, since all limits along subsequences coincide.

We leave it as an exercise to extend the proof from the case when $\mathcal{C}$ is a piece of $\mathbb{R}$, say $[a, b]$ with $t = 0 \in (a, b)$, approached from above, to a more general smooth bounded curve (open or not): parametrize the curve.

Proof. We can assume without loss of generality that $0 \in [a, b]$, $t = 0$ and $|a| > |b|$.

(i) We write $\phi(s) = (\phi(s) - \phi(0)) + \phi(0)$ and show that both PV integrals exist. By symmetry,

$$(46.7) \quad PV \int_{-b}^b s^{-1} \phi(0) ds = 0$$

implying

$$(46.8) \quad PV \int_a^b \frac{\phi(0)}{s} ds = \int_a^{-b} \frac{\phi(0)}{s} ds$$

We then note that $|\phi(s) - \phi(0)|/|s| = (|\phi(s) - \phi(0)|s|^{-\beta})|s|^{\beta-1} \in L^1([a, b])$, hence

$$(46.9) \quad PV \int_a^b \frac{\phi(s) - \phi(0)}{s} ds = \int_a^b \frac{\phi(s) - \phi(0)}{s} ds$$

(ii) Take $\varepsilon > 0$ small enough, n large enough, and $0 < c \in (a, b)$, change variable to $u = s - x_n$ and break the interval of integration into $[a - x_n, -c]$, $[-c, c]$ and $[c, b - x_n]$.

$$(46.10) \quad \lim_{n \rightarrow \infty} \left(\int_{a-x_n}^{-c} + \int_c^{b-x_n} \right) \frac{\phi(u + x_n) du}{u - iy_n} = \left(\int_a^{-c} + \int_c^b \right) \frac{\phi(u) du}{u}$$

We decompose the integral on $[-c, c]$ into

$$(46.11) \quad \int_{-c}^c \frac{\phi(u + x_n) - \phi(x_n)}{|u|^\beta} \frac{|u|^\beta du}{u - iy_n} + \phi(x_n) \int_C \frac{du}{u}$$

where in the last integral we homotopically deformed the contour into a half-circle of radius c in the LHP centered at zero. For all n , the integrand in the first integral is bounded in absolute value, up to a constant, by $|u|^{\beta-1} \in L^1([-c, c])$ while the second integral evaluates to πi . The result now follows by using dominated convergence and the symmetry argument in (i). ■

The following result follows immediately.

Theorem 46.4 (Existence). *Under the conditions of Theorem 46.2, the function in (46.2) solves the Riemann-Hilbert problem in §45.1.*

Note 46.5. *The Hölder condition can be replaced with the weaker Dini condition, with the same proof.*

Note 46.6. (a) The function defined by the Cauchy type integral (46.2) is called **sectionally analytic**. With the convention about the sides of the curve mentioned before, functions that are boundary values of Cauchy type integrals are sometimes denoted $\oplus$ and $\ominus$ functions, respectively.

(b) Branch jumps of analytic functions are an essential ingredient in Sato's theory of hyperfunctions.

46.3. Examples.

46.3.1. *A very simple example.* Find a function Φ analytic in $\mathbb{C} \setminus \partial\mathbb{D}$ such that along $\partial\mathbb{D}$ we have

$$(46.12) \quad \Phi^+(t) - \Phi^-(t) = 1$$

Note that for this problem the set of analyticity of Φ is a union of two disjoint domains. Since Φ is discontinuous across $\partial\mathbb{D}$, we are dealing with two separate analytic functions. Plemelj's formula reads

$$(46.13) \quad \Phi(z) = \frac{1}{2\pi i} \int_{\partial\mathbb{D}} \frac{1}{s-z} ds$$

Clearly, if $z \in \mathbb{D}$ (which is to the left of $\partial\mathbb{D}$ oriented positively), we have

$$(46.14) \quad \Phi_{\text{in}}(z) = \frac{1}{2\pi i} \int_{\partial\mathbb{D}} \frac{1}{s-z} ds = 1$$

Likewise, if z is outside $\mathbb{D}$ we have

$$(46.15) \quad \Phi_{\text{out}}(z) = \frac{1}{2\pi i} \int_{\partial\mathbb{D}} \frac{1}{s-z} ds = 0$$

Both Φ_{in} and Φ_{out} are analytic, but not analytic continuations of each other, so in this case our sectionally analytic function is really a pair of distinct analytic functions. We leave the question of uniqueness to the next subsection when the contour is open and which leads to a more interesting discussion.

46.3.2. *Another simple example.* Find a function Φ analytic in $\mathbb{C} \setminus [-1, 1]$ such the upper and lower limits across $[-1, 1]$ satisfy

$$(46.16) \quad \Phi^+(z) - \Phi^-(z) = 1$$

46.3.3. *A solution.* According to Plemelj's formulas a function satisfying (46.16) is given by

$$(46.17) \quad \Phi(z) = \frac{1}{2\pi i} \int_{-1}^1 \frac{1}{s-z} ds$$



FIGURE 49. Contour used for applying Morera's theorem.

This formula shows Φ is analytic on the universal cover of $\hat{\mathbb{C}} \setminus \{-1, 1\}$ (Φ is analytic at infinity). We get explicitly

$$(46.18) \quad \Phi(z) = -\frac{1}{2\pi i} (\ln(z-1) - \ln(z+1))$$

where the log is determined by the condition $\Phi(z) < 0$ for $z > 1$. It is useful to check directly on (46.18) that $\Phi^+ - \Phi^- = 1$ on $(-1, 1)$

46.3.4. *Calculating principal value integrals.* Plemelj's formulas help us calculate principal values integrals as well, sometimes in a simpler way. Let $\mathcal{C}$ be a simple smooth closed curve and assume that $f(z)$ is analytic in $\text{Int}(\mathcal{C})$ and Hölder continuous in the closure of $\text{Int}(\mathcal{C})$. Then,

$$(46.19) \quad \frac{1}{2\pi i} PV \int_{\mathcal{C}} \frac{f(s)}{s-t} ds = \frac{1}{2} f(t)$$

a "limiting case" of a Cauchy formula.

46.3.5. *Uniqueness issues.* What obvious freedoms do we have in solving the problem in §46.3.2? As mentioned, we can add to Φ any entire function. More generally, we can add any function analytic in $\mathbb{C} \setminus \{-1, 1\}$ (note that this means single-valued). The following theorem shows that these are all the freedoms.

Theorem 46.7 (Uniqueness). *Consider the problem in §45.1 with the following further conditions:*

- (1) $\Phi(z) \rightarrow 0$ as $|z| \rightarrow \infty$;
- (2) $(1 \mp z)\Phi(z) \rightarrow 0$ as $z \rightarrow \pm 1$

Then the solution is unique, namely (46.2).

Proof. The Φ in (46.2) satisfies this condition, as it is easy to verify. Assume Φ_1 is another solution with the same properties. Then $f = \Phi - \Phi_1$ is analytic in $\mathbb{C} \setminus [-1, 1]$ and continuous on $(-1, 1)$, entailing continuity in $\mathbb{C} \setminus \{-1, 1\}$. Show that the contour integral of $\Phi - \Phi_1$ on a circle of radius ε around 1 and -1 is zero, hence it vanishes on any

closed contour in $\mathbb{C}$ (see Fig. 49) implying, by Morera, that $\Phi - \Phi_1$ is entire. Since $\lim_{z \rightarrow \infty} (\Phi(z) - \Phi_1(z)) = 0$, $\Phi - \Phi_1 = 0$. ■

Exercise 46.8. Check using Plemelj's formulas that any simple smooth curve $\mathcal{C}$ in $\mathbb{C}$ is the natural boundary of a large class analytic of functions, and furthermore the limit of such functions on $\mathcal{C}$ can be as smooth as we want, short of analytic. Use the Riemann mapping theorem to show that any simple Jordan curve is a natural boundary of a large class of analytic functions.

We see from

46.3.6. *Degree of a function at infinity.* By definition Φ has degree k at infinity if for some $C \neq 0$ we have

$$(46.20) \quad \Phi(z) = Cz^k + O(z^{k-1}) \quad \text{as } z \rightarrow \infty$$

The function Φ has finite degree at infinity if $\Phi = o(z^m)$ for some m .

47. MORE GENERAL SCALAR R-H PROBLEMS

47.1. Scalar homogeneous R-H problems.

47.1.1. *Scalar homogeneous R-H problems.* This is a problem of the type

$$(47.1) \quad \Phi^+ = g\Phi^- \quad \text{on } \mathcal{C}$$

where $\mathcal{C}$ is a smooth simple closed contour, g nonzero on $\mathcal{C}$ and satisfying a Hölder condition on $\mathcal{C}$. We are looking for solutions of degree m at infinity.

Solution to (47.1) First we note that if Φ is a solution and H is entire, then by homogeneity ΦH is also a solution.

Assume that the *index of g w.r.t. $\mathcal{C}$* is k . Without loss of generality we assume $0 \in \text{Int}(\mathcal{C})$. We can rewrite the problem as

$$(47.2) \quad \Phi^+(t) = (t^{-k}g(t))(t^k\Phi^-) \quad \text{on } \mathcal{C}$$

Formally for now, taking the log of both sides, we get

$$(47.3) \quad \ln \Phi^+(t) = \ln(t^{-k}g(t)) + \ln(t^k\Phi^-) \quad \text{on } \mathcal{C}$$

or finally, with obvious notation,

$$(47.4) \quad \Gamma^+(t) = f(t) + \Gamma^-(t) \quad \text{on } \mathcal{C}$$

The reason we formed the combination $t^{-k}g(t)$, where we choose k to be the index of ϕ w.r.t $\mathcal{C}$, is to ensure Hölder continuity of f . Otherwise, since $\arg \phi$ changes by $2k\pi$ upon traversing $\mathcal{C}$, f would have a jump discontinuity somewhere on $\mathcal{C}$.

A solution to (47.4) is given by Plemelj's formulas:

$$(47.5) \quad \Gamma(z) = \frac{1}{2\pi i} \int_{\mathcal{C}} \frac{f(s)}{s-z} ds$$

The exterior of $\mathcal{C}$ corresponds to the negative sign and we have $\Gamma^- = O(z^{-1})$ for large z , and thus $\exp(\Gamma^-) = z^k \Phi^- = 1 + o(1)$ for large z , hence $\Phi^- = z^{-k} + o(z^{-k})$. We finally get a solution of degree m at infinity by multiplying by P_{m+k} , a polynomial of degree $m+k$. The solution of (47.1) is then

$$(47.6) \quad \Phi = \begin{cases} P_{m+k}(z) \exp(\Gamma(z)), & z \text{ inside } \mathcal{C} \\ z^{-k} P_{m+k}(z) \exp(\Gamma(z)), & z \text{ outside } \mathcal{C} \end{cases}$$

We will not, for reasons of space, discuss uniqueness issues here.

47.1.2. *Inhomogeneous R-H problems.* These are equations of the form

$$(47.7) \quad \Phi^+ = g\Phi^- + f$$

Where f and g are Hölder continuous. These can be brought to Plemelj's formulas in the following way. We first solve the homogeneous problem

$$(47.8) \quad \Psi^+ = g\Psi^-$$

which we dealt with in §47.1.1, and look for a solution of (47.7) in the form $\Phi = X\Psi$. We get

$$(47.9) \quad X^+\Psi^+ = gX^-\Psi^- + f \Rightarrow X^-g\Psi^+ = X^-g\Psi^- + f \Rightarrow \Psi^+ - \Psi^- = \frac{f}{gX^-}$$

which is of the form we already solved.

47.2. Applications.

47.2.1. *Inhomogeneous singular integral equations.* These are equations of the form

$$(47.10) \quad a(t)\phi(t) + b(t)PV \int_{\mathcal{C}} \frac{\phi(s)}{s-z} ds = c(t)$$

with a, b, c Hölder continuous and the further condition $i\pi a(t) \pm b(t) \neq 0$. We attempt to write, guided by Plemelj's formulas

$$(47.11) \quad \phi(t) = \Phi^+(t) - \Phi^-(t)$$

and

$$(47.12) \quad \Phi(z) = \frac{1}{2\pi i} \int_{\mathcal{C}} \frac{\phi(s)}{s-z} ds$$

and then

$$(47.13) \quad PV \int_{\mathcal{C}} \frac{\phi(s)}{s-t} ds = i\pi [\Phi^+(t) + \Phi^-(t)]$$

where, of course ϕ is still unknown. The equation becomes

$$(47.14) \quad a(t) [\Phi^+(t) - \Phi^-(t)] + b(t)i\pi [\Phi^+(t) + \Phi^-(t)] = c(t)$$

or

$$(47.15) \quad \Phi^+(t)(a(t) + b(t)i\pi) + \Phi^-(t)(b(t)i\pi - a(t)) = c(t)$$

or, finally,

$$(47.16) \quad \Phi^+(t) = \frac{a(t) - b(t)i\pi}{a(t) + b(t)i\pi} \Phi^-(t) + \frac{c(t)}{a(t) + b(t)i\pi}$$

which is of the form (47.7) which we addressed already. Care must be taken that the chosen solution Φ is such that $\Phi^+ + \Phi^-$ has the behavior (46.3) at infinity. Then the substitution is a posteriori justified. We did not discuss whether there are other solutions of the integral equation. A complete discussion of this and related equations can be found in [11].

We choose one of the applications in [1], the solution of the Dirichlet problem for the Laplacian in the upper half plane, with condition f on the boundary, $\mathbb{R}$. The problem can be reformulated as a Riemann-Hilbert problem of the form (47.16), see [1], but in this case, the solution is obtained easily from Plemelj's formulas.

For this purpose we look for an analytic function in $\mathbb{H}^u$ generated by u . As we know, this is

$$(47.17) \quad \Phi^+ = u + iv$$

where v is the harmonic conjugate of u , unique up to a constant. Now note that the function

$$(47.18) \quad \Phi(z) = \frac{1}{\pi i} \int_{\mathbb{R}} \frac{f(s)}{s-z} ds$$

is analytic in the upper half plane and, if f is Hölder continuous, then by Plemelj's formulas we have

$$(47.19) \quad \lim_{z \downarrow t \in \mathbb{R}} \Phi(z) = f(t) + \frac{1}{\pi i} PV \int_{\mathbb{R}} \frac{f(s)}{s-t} ds$$

In particular $u = \operatorname{Re} \Phi$ is harmonic and has the limit $f(x, y)$ as $(x, y) \rightarrow (t, 0)$. Now, simply writing $z = x + iy$ and taking the real part we get the solution in the Poisson kernel form,

$$(47.20) \quad u(x, y) = \frac{y}{\pi} \int_{-\infty}^{\infty} \frac{f(\tau) d\tau}{(\tau - x)^2 + y^2}$$

The condition that f is Hölder can be relaxed to mere continuity as follows. To better adapt to the limit $y \rightarrow 0$ we change variable to $t = x + \beta y$ and obtain

$$(47.21) \quad u(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{f(x + \beta y)}{\beta^2 + 1} d\beta$$

which, by dominated convergence, has $f(x)$ as the limit when $y \rightarrow 0$.

48. ASYMPTOTIC SERIES

We have seen in the Schwarz-Christoffel section that the behavior of analytic functions near a point of nonanalyticity can be given by a series in noninteger powers of the distance to the singularity. The behavior can be more complicated, containing exponentially small corrections, logarithmic terms and so on. The series themselves may have zero radius of convergence. It is not the purpose of this part of the course to classify these behaviors, but it can be done for a fairly large class of functions. Here we look how simple behaviors can be determined for relatively simple functions.

Example 48.1. *Consider the following integral related to the so-called error function*

$$F(z) = e^{z^{-2}} \int_0^z s^{-2} e^{-s^{-2}} ds$$

It is clear that the integral converges at the origin, if the origin is approached through real values (see also the change of variable below).

Definition of $F(z)$. *We define the integral to $z \in \mathbb{C}$ as being taken on a curve γ with $\gamma'(0) > 0$, and define $F(0) = 0$.*

Check that this is a consistent definition and the resulting function is analytic except at $z = 0$ (this is essentially the contents of Exercise 48.3 below).

What about the behavior at $z = 0$? It depends on the direction in which 0 is approached! Let's look more carefully. Replace z by $1/x$, make a corresponding change of variable in the integral and you are led to

$$(48.1) \quad E(x) = e^{x^2} \int_x^\infty e^{-s^2} ds =: \frac{\sqrt{\pi}}{2} e^{x^2} \operatorname{erfc}(x)$$

Let us take x (and thus z) real and integrate by parts m times

$$(48.2) \quad \begin{aligned} E(x) &= \frac{1}{2x} - \frac{e^{x^2}}{2} \int_x^\infty \frac{e^{-s^2}}{s^2} ds = \frac{1}{2x} - \frac{1}{4x^3} + \frac{3e^{x^2}}{4} \int_x^\infty \frac{e^{-s^2}}{s^4} ds = \dots \\ &= \sum_{k=0}^{m-1} \frac{(-1)^k}{2\sqrt{\pi}} \frac{\Gamma(k + \frac{1}{2})}{x^{2k+1}} + \frac{(-1)^m e^{x^2} \Gamma(m + \frac{1}{2})}{\sqrt{\pi}} \int_x^\infty \frac{e^{-s^2}}{s^{2m}} ds \end{aligned}$$

On the other hand, we have, by L'Hospital

$$(48.3) \quad \left(\int_x^\infty \frac{e^{-s^2}}{s^{2m}} ds \right) \left(\frac{e^{-x^2}}{x^{2m+1}} \right)^{-1} \rightarrow \frac{1}{2} \text{ as } x \rightarrow \infty$$

and the last term in (48.2) is $O(x^{-2m-1})$ as well. On the other hand it is also clear that the series in (48.2) is alternating and thus

$$(48.4) \quad \sum_{k=0}^{m-1} \frac{(-1)^k}{2\sqrt{\pi}} \frac{\Gamma(k + \frac{1}{2})}{x^{2k+1}} \leq E(x) \leq \sum_{k=0}^m \frac{(-1)^k}{2\sqrt{\pi}} \frac{\Gamma(k + \frac{1}{2})}{x^{2k+1}}$$

if m is even.

Remark 48.2. Using (48.3) and Exercise 48.13 below we conclude that $F(z)$ has a Taylor series at zero,

$$(48.5) \quad \tilde{F}(z) = \sum_{k=0}^{\infty} \frac{(-1)^k}{2\sqrt{\pi}} \Gamma(k + \frac{1}{2}) z^{2m+1}$$

that $F(z)$ is C^∞ on $\mathbb{R}$ and analytic away from zero.

Exercise 48.3. ** Show that $z = 0$ is an isolated singularity of $F(z)$. Using Remark 48.2, show that F is unbounded as 0 is approached along some directions in the complex plane.

Notes (1) It is *not* the Laurent series of f at 0 that we calculated! Laurent series converge. Think carefully about this distinction and why the positive index coefficients do not coincide.

(2) The rate of convergence of the Laurent series is slower as 0 is approached, quickly becoming numerically useless. By contrast, the precision gotten from (48.4) near zero is such that for $z = 0.1$ the error in calculating f is of order 10^{-45} ! However, of course (48.4) is divergent and it cannot be used to calculate *exactly* for any nontrivial value of z .

(3) We have illustrated here a simple method of evaluating the behavior of integrals, the method of integration by parts.

48.1. More general asymptotic series. Classical asymptotic analysis typically deals with the qualitative and quantitative description of the behavior of a function close to a point, usually a singular point of the function. This description is provided in the form of an *asymptotic expansion*. The expansion certainly depends on the point studied and, as we have noted, often on the direction along which the point is approached (in the case of several variables, it also depends on the relation between the variables as the point is approached). If the direction matters, it is often convenient to change variables to place the special point at infinity.

Asymptotic expansions are formal series¹⁹ of simpler functions f_k ,

$$(48.6) \quad \tilde{f} = \sum_{k=0}^{\infty} f_k(t)$$

in which each successive term is much smaller than its predecessors (one variable is assumed for clarity). For instance if the limiting point is t_0 approached from above along the real line this requirement is written

$$(48.7) \quad f_{k+1}(t) = o(f_k(t)) \quad \text{or} \quad f_{k+1}(t) \ll f_k(t) \quad \text{as } t \downarrow t_0$$

denoting

$$(48.8) \quad \lim_{t \rightarrow t_0^+} f_{k+1}(t)/f_k(t) = 0$$

We will often use the variable x when the limiting point is $+\infty$ and z when the limiting point is zero. Simple examples are the Taylor series, e.g.

$$\sin z + e^{-\frac{1}{z}} \sim z - \frac{z^3}{6} + \dots \quad (z \rightarrow 0^+)$$

and the expansion in the Stirling formula

$$\ln \Gamma(x) \sim x \ln x - x - \frac{1}{2} \ln x + \frac{1}{2} \ln(2\pi) + \sum_{n=1}^{\infty} \frac{B_{2n}}{2n(2n-1)x^{2n-1}}, \quad x \rightarrow +\infty$$

where B_k are the Bernoulli numbers.

¹⁹That is, there are no convergence requirements. More precisely, they are defined as sequences $\{f_k\}_{k \in \mathbb{N} \cup \{0\}}$, the operations being defined in the same way as if they represented convergent series; see also §48.2.

(The asymptotic expansions in the examples above are the formal sums following the “ $\sim$ ” sign, the meaning of which will be explained shortly.)

Examples of expansions that are *not* asymptotic expansions are

$$\sum_{k=0}^{\infty} \frac{x^k}{k!} \quad (x \rightarrow +\infty)$$

which converges to $\exp(x)$, but it is not an asymptotic series for large x since it fails (48.7); another example is the series

$$(48.9) \quad \sum_{k=0}^{\infty} \frac{x^{-k}}{k!} + e^{-x} \quad (x \rightarrow +\infty)$$

(because of the exponential terms, this is not an ordered *simple series* satisfying (48.7)). Note however expansion (48.9), *does* satisfies all requirements in the *left* half plane, if we write e^{-x} in the first position.

We also note that in this particular case the first series is convergent, and if we replace (48.9) by

$$(48.10) \quad e^{1/x} + e^{-x}$$

then (48.10) *is* a valid asymptotic expansion, of a very simple kind, with two nonzero terms. Since convergence is relative to a topology, this elementary remark will play a crucial role when we will speak of Borel summation.

Functions asymptotic to a series, in the sense of Poincaré. The relation $f \sim \tilde{f}$ between an actual function and a formal expansion is defined as a sequence of limits:

Definition 48.4. *A function f is asymptotic to the formal series $\tilde{f}$ as $t \rightarrow t_0^+$ if*

$$(48.11) \quad f(t) - \sum_{k=0}^N \tilde{f}_k(t) =: f(t) - \tilde{f}^{[N]}(t) = o(\tilde{f}_N(t)) \quad (\forall N \in \mathbb{N})$$

We note that condition (48.11) can then be also written as

$$(48.12) \quad f(t) - \sum_{k=0}^N \tilde{f}_k(t) = O(\tilde{f}_{N+1}(t)) \quad (\forall N \in \mathbb{N})$$

where $g(t) = O(h(t))$ means $\limsup_{t \rightarrow t_0^+} |g(t)/h(t)| < \infty$. Indeed, this follows from (48.11) and the fact that $f(t) - \sum_{k=0}^{N+1} \tilde{f}_k(t) = o(\tilde{f}_{N+1}(t))$.

48.2. Asymptotic power series. In many instances the functions f_k are exponentials, powers and logarithms. This is not simply a matter of choice or an accident, but reflects some important fact about the relation between asymptotic expansions and functions which will be clarified later.

A special role is played by power series which are series of the form

$$(48.13) \quad \tilde{S} = \sum_{k=0}^{\infty} c_k z^k, \quad z \rightarrow 0^+$$

With the transformation $z = t - t_0$ (or $z = x^{-1}$) the series can be centered at t_0 (or $+\infty$, respectively).

Remark. If a c_k is zero then Definition 48.4 fails trivially in which case (48.13) is not an asymptotic series. This motivates the following definition.

Definition 48.5 (Asymptotic power series). *A function possesses an asymptotic power series if*

$$(48.14) \quad f(z) - \sum_{k=0}^N c_k z^k = O(z^{N+1}) \quad (\forall N \in \mathbb{N})$$

We use the boldface notation $\sim$ for the stronger asymptoticity condition in (48.11) when confusion is possible.

Example Check that the Taylor series of an analytic function at zero is its asymptotic series there.

In the sense of (48.14), the asymptotic power series at zero of e^{-1/x^2} is the zero series. It is however surely not the case that e^{-1/x^2} behaves like zero as $x \rightarrow 0$ on $\mathbb{R}$. Rather, in this case, the asymptotic *behavior* of e^{-1/x^2} is e^{-1/x^2} itself (only exponentials and powers involved).

Asymptotic power series form an algebra; addition of asymptotic power series is defined in the usual way:

$$A \sum_{k=0}^{\infty} c_k z^k + B \sum_{k=0}^{\infty} c'_k z^k = \sum_{k=0}^{\infty} (Ac_k + Bc'_k) z^k$$

while multiplication is defined as in the convergent case

$$\left(\sum_{k=0}^{\infty} c_k z^k \right) \left(\sum_{k=0}^{\infty} c'_k z^k \right) = \sum_{k=0}^{\infty} \left(\sum_{j=0}^k c_j c'_{k-j} \right) z^k$$

Remark 48.6. *If the series $\tilde{f}$ is convergent and f is its sum (note the ambiguity of the “sum” notation) $f = \sum_{k=0}^{\infty} c_k z^k$ then $f \sim \tilde{f}$.*

The proof of this remark follows directly from the definition of convergence.

Lemma 48.7. *(Uniqueness of the asymptotic series to a function) If $f(z) \sim \tilde{f} = \sum_{k=0}^{\infty} \tilde{f}_k z^k$ as $z \rightarrow 0$ then the $\tilde{f}_k$ are unique.*

Proof. Assume that we also have $f(z) \sim \tilde{F} = \sum_{k=0}^{\infty} \tilde{F}_k z^k$. We then have (cf. (48.11))

$$\tilde{F}^{[N]}(z) - \tilde{f}^{[N]}(z) = o(z^N)$$

which is impossible unless $g_N(z) = \tilde{F}^{[N]}(z) - \tilde{f}^{[N]}(z) = 0$, since g_N is a polynomial of degree N in z . ■

Corollary 48.8. *The asymptotic series at the origin of an analytic function is its Taylor series at zero. More generally, if F has a Taylor series at 0 then that series is its asymptotic series as well.*

The proof of the following lemma is immediate:

Lemma 48.9. *(Algebraic properties of asymptoticity to a power series)*

If $f \sim \tilde{f} = \sum_{k=0}^{\infty} c_k z^k$ and $g \sim \tilde{g} = \sum_{k=0}^{\infty} d_k z^k$ then

$$(i) \quad Af + Bg \sim A\tilde{f} + B\tilde{g}$$

$$(ii) \quad fg \sim \tilde{f}\tilde{g}$$

Sometimes it is convenient to check a formally weaker condition of asymptoticity:

Lemma 48.10. *Let $\tilde{f} = \sum_{n=0}^{\infty} a_n z^n$. If f is such that there exists a sequence $p_n \rightarrow \infty$ such that*

$$\left(\forall n \exists p_n \right) \text{ s.t. } f(z) - \tilde{f}^{[p_n]}(z) = o(z^n) \text{ as } z \rightarrow 0$$

then $f \sim \tilde{f}$.

Proof. We let m be arbitrary and choose $n > m$ such that $p_n > m$. We have

$$f(z) - \tilde{f}^{[m]} = (f(z) - \tilde{f}^{[p_n]}) + (\tilde{f}^{[p_n]} - \tilde{f}^{[m]}) = o(z^n) \quad (z \rightarrow 0)$$

by assumption and since $\tilde{f}^{[p_n]} - \tilde{f}^{[m]}$ is a polynomial for which the smallest power is z^{m+1} (we are dealing with truncates of the same series). ■

48.3. Integration and differentiation of asymptotic power series. While asymptotic power series can be safely integrated term by term as the next proposition shows, differentiation is more delicate. In suitable spaces of functions and expansions, we will see the asymmetry largely disappears if we are dealing with analytic functions in suitable regions.

Anyway, for the moment note that the function $e^{-1/z} \sin(e^{1/z^2})$ is asymptotic to the zero power series as $z \rightarrow 0^+$ although the derivative is unbounded and thus not asymptotic to the zero series.

Proposition 48.11. *Assume f is integrable near $z = 0$ and that*

$$f(z) \sim \tilde{f}(z) = \sum_{k=0}^{\infty} \tilde{f}_k z^k$$

Then

$$\int_0^z f(s) ds \sim \int \tilde{f} := \sum_{k=0}^{\infty} \frac{\tilde{f}_k}{k+1} z^{k+1}$$

Proof. This follows from the fact that $\int_0^z o(s^n) ds = o(z^{n+1})$ as can be seen by immediate estimates. ■

Asymptotic power series of analytic function, if they are valid in wide enough regions can be differentiated.

Asymptotics in a strip. Assume $f(x)$ is analytic in the strip $S_a = \{x : |x| > R, |\operatorname{Im}(x)| < a\}$. Let $\alpha < a$ and $S_\alpha = \{x : |x| > R, |\operatorname{Im}(x)| < \alpha\}$ and assume that

$$(48.15) \quad f(x) \sim \tilde{f}(x) = \sum_{k=0}^{\infty} c_k x^{-k} \quad (|x| \rightarrow \infty, x \in S_\alpha)$$

It is assumed that the limits implied in (48.15) hold uniformly in the given strip.

Proposition 48.12. *If (48.15) holds, then, for $\alpha' < \alpha$ we have*

$$f'(x) \sim \tilde{f}'(x) := \sum_{k=0}^{\infty} -\frac{k c_k}{x^{k+1}} \quad (|x| \rightarrow \infty, x \in S_{\alpha'})$$

Proof. We have $f(x) = \tilde{f}^{[N]}(x) + g_N(x)$ where clearly g is analytic in S_a and $|g_N(x)| \leq \text{Const.} |x|^{-N-1}$ in S_α . But then, for $x \in S_{\alpha'}$ and $\delta = \frac{1}{2}(\alpha - \alpha')$ we get

$$\begin{aligned}
|g'_N(x)| &= \frac{1}{2\pi} \left| \oint_{|x-s|=\delta} \frac{g_N(s)ds}{(s-x)^2} \right| \leq \frac{1}{\delta} \frac{\text{Const.}}{(|x| - |\delta|)^{N+1}} \\
&= O(x^{-N-1}) \quad (|x| \rightarrow \infty, x \in S_{\alpha'})
\end{aligned}$$

By Lemma 48.10, the proof follows. ■

Exercise 48.13. ** Show that if $f(x)$ is continuous on $[0, 1]$ and differentiable on $(0, 1)$ and $f'(x) \rightarrow L$ as $x \downarrow 0$, then f is differentiable to the right at zero and this derivative equals L . Use this fact, Proposition 48.12 and induction to show that the Taylor series at the origin of $F(z)$ is indeed given by (48.5).

48.4. Watson's Lemma. In many instances integral representations of functions are amenable to Laplace transforms

$$(48.16) \quad (\mathcal{L}F)(x) := \int_0^\infty e^{-xp} F(p) dp$$

The behavior of $\mathcal{L}F$ for large x relates to the behavior for small p of F .

It is shown in the later parts of this book that solutions of generic analytic differential equations, under mild assumptions can be conveniently expressed in terms of Laplace transforms.

For the error function note that

$$\int_N^\infty e^{-s^2} ds = N \int_1^\infty e^{-N^2 u^2} du = \frac{\sqrt{x} e^{-x}}{2} \int_0^\infty \frac{e^{-xp}}{\sqrt{p+1}} dp; \quad x = N^2$$

For the Gamma function,

$$(48.17) \quad \Gamma(n+1) = \int_0^\infty e^{-t} t^n dt = n^{n+1} \int_0^\infty e^{n(-s+\ln s)} ds$$

writing $\int_0^\infty = \int_0^1 + \int_1^\infty$ we can make the substitution $t - \ln t = p$ in each of the two integrals and obtain

$$(48.18) \quad \Gamma(n+1) = n^{n+1} e^{-n} \int_0^\infty e^{-np} G(p) dp;$$

$$G(p) = \frac{1}{1 + W(0, -e^{-u-1})} - \frac{1}{1 + W(-1, -e^{-u-1})}$$

where W is the Lambert function, the inverse function of xe^x ; $W(0, \cdot)$ is the principal branch of W and $W(-1, \cdot)$ is the -1 branch.

Watson's Lemma

This important tool states that the asymptotic series at infinity of $(\mathcal{L}F)(x)$ is obtained by formal term-by-term integration of the asymptotic series of $F(p)$ for small p , provided F has such a series.

Lemma 48.14. *Let $F \in L^1(\mathbb{R}^+)$ and assume $F(p) \sim \sum_{k=0}^{\infty} c_k p^{k\beta_1 + \beta_2 - 1}$ as $p \rightarrow 0^+$ for some constants β_i with $\operatorname{Re}(\beta_i) > 0$, $i = 1, 2$. Then*

$$\mathcal{L}F \sim \sum_{k=0}^{\infty} c_k \Gamma(k\beta_1 + \beta_2) x^{-k\beta_1 - \beta_2}$$

along any ray ρ in the open right half plane H .

Proof. Induction, using the simpler version, Lemma 48.15, proved below. $\square$

Lemma 48.15. *Let $F \in L^1(\mathbb{R}^+)$, $x = \rho e^{i\phi}$, $\rho > 0$, $\phi \in (-\pi/2, \pi/2)$ and assume*

$$F(p) \sim p^\beta \quad \text{as } p \rightarrow 0^+$$

with $\operatorname{Re}(\beta) > -1$. Then

$$\int_0^\infty F(p) e^{-px} dp \sim \Gamma(\beta + 1) x^{-\beta-1} \quad (\rho \rightarrow \infty)$$

Proof. If $U(p) = p^{-\beta} F(p)$ we have $\lim_{p \rightarrow 0} U(p) = 1$. Let χ_A be the characteristic function of the set A and $\phi = \arg(x)$. We choose C and a positive so that $|F(p)| < C|p^\beta|$ on $[0, a]$. Since

$$(48.19) \quad \left| \int_a^\infty F(p) e^{-px} dp \right| \leq e^{-|x|a \cos \phi} \|F\|_1$$

we have by dominated convergence, and after the change of variable $s = p|x|$,

$$(48.20) \quad \begin{aligned} x^{\beta+1} \int_0^\infty F(p) e^{-px} dp &= e^{i\phi(\beta+1)} \int_0^\infty s^\beta U(s/|x|) \chi_{[0,a]}(s/|x|) e^{-se^{i\phi}} ds \\ &\quad + O(|x|^{\beta+1} e^{-|x|a \cos \phi}) \rightarrow \Gamma(\beta + 1) \quad (|x| \rightarrow \infty) \end{aligned}$$

48.5. The Gamma function.

Remark 48.16. *The function $G(p)$ is an analytic function in $\sqrt{p}$ and thus $G'(p)$ has a convergent Puiseux series*

$$\sum_{k=-1}^{\infty} c_k p^{k/2} = \sqrt{2} p^{-1/2} + \frac{\sqrt{2}}{6} p^{1/2} + \frac{\sqrt{2}}{216} p^{3/2} - \frac{139\sqrt{2}}{97200} p^{5/2} + \dots$$

Thus, by Watson's Lemma, for large n we have

$$(48.21) \quad n! \sim \sqrt{2\pi n} n^n e^{-n} \left(1 + \frac{1}{12n} + \frac{1}{288n^2} - \frac{139}{51840n^3} + \dots \right)$$

Proof. It is easy to calculate this expansion directly from the definition, $s - \ln s = p$, without using the Lambert function.

We write $s = 1 + S$ and $t = 1 + p$ and the equation $s - \ln(s) = t$ becomes $S - \ln(1 + S) = p$. Note that $S - \ln(1 + S) = S^2 U(S)/2$ where $U(0) = 1$ and $U(S)$ is analytic for small S ; with the natural branch of the square root, $\sqrt{U(S)} = H(S)$ is also analytic. We rewrite $S - \ln(1 + S) = p$ as $SH(S) = \pm\sqrt{2}\sigma$ where $\sigma^2 = p$. Since $(SH(S))'_{|S=0} = 1$ the implicit function theorem ensures the existence of two functions $S_{1,2}(\sigma)$ (corresponding to the two choices of sign) which are analytic in σ . The concrete expansion may be gotten by implicit differentiation in $SH(S) = \pm\sqrt{2}\sigma$, for instance. ■

48.6. Lebesgue's Dominated Convergence Theorem. Let $\{f_n\}_{n \in \mathbb{N}}$ be a sequence of complex-valued measurable functions on a measure space (S, Σ, μ) . (Frequently, in our applications the measure is of the form $d\mu = h(x) dx$ where h is some “nice” weight function.)

Suppose that the sequence converges pointwise to a function f , i.e.

$$\lim_{n \rightarrow \infty} f_n(x) = f(x), \quad \forall x \in S.$$

Assume moreover that the sequence $\{f_n\}$ is dominated by some integrable function $g \in L^1(S, \mu)$, i.e.

$$|f_n(x)| \leq |g|(x), \quad \forall x \in S, \forall n,$$

with $\int_S |g| d\mu < \infty$.

Then $f_n, f \in L^1(S, \mu)$ and

$$\lim_{n \rightarrow \infty} \int_S f_n d\mu = \int_S \lim_{n \rightarrow \infty} f_n d\mu = \int_S f d\mu.$$

In fact, more is true:

$$\lim_{n \rightarrow \infty} \int_S |f_n - f| d\mu = 0.$$

48.7. Appendix to Chapter 8.

48.8. Some facts about the topology of $\mathbb{C}$. From a topological point of view, $\mathbb{C}$ is isomorphic to $\mathbb{R}^2$: we identify the point $z = x + iy$ with $(x, y) = (\operatorname{Re} z, \operatorname{Im} z)$ and the length $|z|$ with $|(x, y)| = \sqrt{x^2 + y^2}$. This induces the metric

$$d(z_1, z_2) := |z_1 - z_2|.$$




FIGURE 50. Open disk $\mathbb{D}(z_0, r) = \{z : |z - z_0| < r\}$: boundary dashed to indicate exclusion.

A region $O \subset \mathbb{C}$ is *open* if with each $z_0 \in O$ it contains some nonempty disk centered at z_0 . Examples include the open disk $\mathbb{D}(z_0, r)$,

the punctured disk $\mathbb{D}_p(z_0, r)$, the upper half-plane $\mathbb{H}^u$, $\mathbb{C}$ itself, and $\emptyset$. By contrast, the *closed disk*

$$\overline{\mathbb{D}}(z_0, r) := \{z \in \mathbb{C} : |z - z_0| \leq r\}$$

is not open.




FIGURE 51. Closed disk $\overline{\mathbb{D}}(z_0, r)$: solid boundary indicates inclusion.

Finite intersections of open sets are open, and arbitrary unions (finite or infinite) of open sets are open.

Definition 48.17. A **topology** on a set X is a family $\mathcal{O}$ of subsets, called *open sets*, such that: (1) $X, \emptyset \in \mathcal{O}$; (2) $O_1, O_2 \in \mathcal{O} \implies O_1 \cap O_2 \in \mathcal{O}$; (3) Arbitrary unions of sets in $\mathcal{O}$ are also in $\mathcal{O}$.

Complements of open sets are called *closed sets*. Note that X and $\emptyset$ are both open and closed. For $\mathbb{C}$, the topology can be taken as the collection of all unions of open disks.

The *boundary* of $S \subset \mathbb{C}$, denoted ∂S , consists of points which can be approached both from within S and from outside S . For example,

$$\partial \mathbb{D}(z_0, r) = \{z \in \mathbb{C} : |z - z_0| = r\}.$$




FIGURE 52. Boundary of the disk: the circle $|z - z_0| = r$.

Since

$$\max(|x|, |y|) \leq \sqrt{x^2 + y^2} \leq |x| + |y|,$$

a sequence $\{z_n\}$ is Cauchy in $\mathbb{C}$ **iff** both $\{x_n\}$ and $\{y_n\}$ are Cauchy in $\mathbb{R}$. Similarly, $z_n \rightarrow 0$ in $\mathbb{C}$ iff $x_n \rightarrow 0$ and $y_n \rightarrow 0$ in $\mathbb{R}$. In $\mathbb{R}^n$, Cauchy sequences are exactly the convergent ones, hence $\mathbb{C}$ is complete.

Consequently, if a series $\sum_{n=0}^{\infty} c_n z_0^n$ converges, then $c_n z_0^n \rightarrow 0$ as $n \rightarrow \infty$. This implies that the real-valued series $\sum_{n=0}^{\infty} |c_n| |z_1|^n$ converges for every $|z_1| < |z_0|$, i.e. $\sum_{n=0}^{\infty} c_n z_1^n$ converges uniformly and absolutely for $|z| < |z_0|$. This is Abel's theorem in the complex setting: the domain of convergence is a disk rather than an interval.

A region is *closed* iff it contains all of its boundary points; it is *open* iff it contains none of them.

48.8.1. *Connected sets.* An open set $O \subset \mathbb{C}$ is *connected* if it is not the union of two disjoint nonempty open sets. More generally, $X_1 \subset X$ is connected if it is not the disjoint union of two nonempty sets open in the induced topology. Equivalently, there is no proper subset of X_1 that is simultaneously open and closed in the induced topology. For example, any disk or punctured disk in $\mathbb{C}$ is connected.




FIGURE 53. Annulus $\{z : r < |z - z_0| < R\}$: region between two excluded boundary circles.

Exercise 48.18. *Is the annulus $\{z \in \mathbb{C} : r \leq |z| < R\}$ open? closed? connected? What is its boundary?*

Proposition 48.19. *Domains $\mathcal{D} \subset \mathbb{C}$ are path connected. Moreover, the connecting path can be chosen polygonal.*

49. SOAP FILMS AND THE MINIMAL SURFACE EQUATION

Let $\Gamma \subset \mathbb{R}^3$ be a simple smooth closed curve (the wire). An equilibrium soap film spanning Γ minimizes the area among admissible




FIGURE 54. Any two points in a domain can be joined by a polygonal path (Proposition 48.19).

surfaces with boundary Γ (Plateau's problem). The Euler–Lagrange equation of the area functional says that the *mean curvature* vanishes:

$$H \equiv 0 \quad \text{on the film.}$$

Equivalently, the *mean curvature vector* $\mathbf{H} = H \mathbf{n}$ is zero.

Graph formulation. If the film can be written locally as a graph $z = u(x, y)$ over a planar domain $\Omega \subset \mathbb{R}^2$, then u solves the *minimal surface equation*

$$(49.2) \quad \operatorname{div} \left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}} \right) = 0 \quad \text{in } \Omega, \quad u|_{\partial\Omega} = u_{\partial\Omega},$$

where the boundary data $u_{\partial\Omega}$ encodes the wire (after projection to the plane).

Parametric (intrinsic) formulation. If $X(u, v) : \Sigma \rightarrow \mathbb{R}^3$ is a smooth immersion with $X(\partial\Sigma) = \Gamma$, the area functional is

$$\mathcal{A}[X] = \iint_{\Sigma} |X_u \times X_v| \, du \, dv.$$

Critical points satisfy $H \equiv 0$. In *isothermal (conformal) parameters* one has the equivalent harmonic map form

$$\Delta X = 0 \quad \text{on } \Sigma, \quad X|_{\partial\Sigma} = \Gamma,$$

where Δ is the flat Laplacian in the (u, v) -coordinates (the conformality constraints ensure $H = 0$).

Remark (bubbles vs. films). A *soap film* has negligible pressure difference across it, hence $H = 0$. A *soap bubble* encloses a volume and has constant mean curvature:

$$H = \frac{\Delta p}{2\sigma} \quad (\text{Laplace–Young law}),$$

with surface tension σ and pressure jump Δp .

REFERENCES

- [1] Mark J. Ablowitz and Athanassios S. Fokas, Complex variables, 2nd edition, Cambridge University Press, 2003.
- [2] M Abramowitz, I A Stegun, Handbook of mathematical functions with formulas, graphs, and mathematical tables New York : Wiley-Interscience (1984).
- [3] L. Ahlfors, Complex Analysis, McGraw-Hill (1979).
- [4] E.A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations*, McGraw-Hill, New York, (1955). (2009).
- [5] O. Forster, Lectures on Riemann Surfaces, Springer New York (1981).
- [6] J. Milnor, Complex Dynamics, Annals of Mathematics Studies, Third Edition, (2006).
- [7] O. Costin and M. Huang, Geometric construction and analytic representation of Julia sets of polynomial maps, Nonlinearity 24, 1311–1327 (2011)
- [8] Rudin, Real and Complex Analysis
- [9] W. Schlag, A Course in Complex Analysis and Riemann Surfaces, Graduate Studies in Mathematics vol. 154, American Mathematical Society, 2014.
- [10] S. Lang, Complex Analysis, fourth edition, (1999).
- [11] N.I. Muskhelishvili, Singular integral equations, Dover (2008).
- [12] Z. Nehari, Conformal Mapping, Dover (1975).
- [13] E. C. Titchmarsh, The theory of functions, Oxford University Press, 1964.
- [14]
- [15] Alexander Hulpke - Own work based on: ModularGroup-FundamentalDomain-01.png,
<https://commons.wikimedia.org/w/index.php?curid=59963451>
- [16] G. Pólya & G. Szegő, Problems and Theorems in Analysis II –see Problem IV.4.8 and its discussion.