

### ALGEBRA I (MATH 6111 AUTUMN 2020) - HOMEWORK 3

*Assumption:* All groups considered in this problem set are finite.

**Problem 1.** Let  $G$  be a group and  $H < G$  be a subgroup. Let  $P < H$  be a Sylow  $p$ -subgroup of  $H$ . Prove that there exists a Sylow  $p$ -subgroup of  $G$  (call it  $Q$ ), satisfying  $P = Q \cap H$ .

**Problem 2.** Let  $G$  be a group and  $N \triangleleft G$  be a normal subgroup. Consider the natural surjection  $\pi: G \rightarrow G/N$ .

(i) Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Prove that  $P \cap N$  is a Sylow  $p$ -subgroup of  $N$ .

(ii) Prove that  $\pi(P)$  is a Sylow  $p$ -subgroup of  $G/N$ .

**Problem 3.** Let  $P < G$  be a Sylow  $p$ -subgroup of  $G$ , and let  $N = N_G(P)$  be the normalizer of  $P$  (that is,  $N = \{g \in G: gPg^{-1} = P\}$ ). Prove that for every  $L < G$  containing  $N$ ,  $N_G(L) = L$ .

**Problem 4.** Let  $H \triangleleft G$  be a normal subgroup of a group  $G$ . Let us assume that  $|H| = p$ . Prove that  $H$  is contained in every Sylow  $p$ -subgroup of  $G$ .

**Problem 5.** Let  $K \triangleleft G$  be a normal subgroup of a group  $G$ , and let  $P$  be a Sylow  $p$ -subgroup of  $K$ . Prove that  $G = KN_G(P) := \{kh: k \in K, h \in N_G(P)\}$ .

**Problem 6.** Let  $G = GL_n(\mathbb{Z}/p\mathbb{Z})$  where  $n \in \mathbb{Z}_{\geq 2}$ .

(i) Show that  $|G| = p^r m$  with  $(m : p) = 1$  and  $r = n(n-1)/2$ .

(ii) Show that the set  $H$  of upper triangular matrices with 1's along the diagonal is a Sylow  $p$ -subgroup of  $G$ .

(iii) Show that the normalizer of  $H$  is the set of all invertible upper triangular matrices. (*Hint:* Use convenient elementary matrices in  $H$  to show any  $A \in G$  with  $AHA^{-1} = H$  must be upper triangular.)

(iv) Conclude from this that the number  $n_p$  of Sylow  $p$ -groups of  $G$  equals:

$$n_p = \prod_{k=1}^n (p^{k-1} + p^{k-2} + \dots + 1) =: [n!]_p.$$

**Problem 7.** Assume that  $G$  is a non-abelian group of order  $p^3$ .

(i) Prove that  $Z(G) \simeq \mathbb{Z}/p\mathbb{Z}$ .

- (ii) Prove that  $G/Z(G) \simeq (\mathbb{Z}/p\mathbb{Z}) \times (\mathbb{Z}/p\mathbb{Z})$ .
- (iii) If  $H < G$  is a subgroup of  $G$  of order  $p^2$ , show that  $H$  is normal and contains  $Z(G)$ .

**Problem 8.** Let  $p, q$  be two prime numbers with  $p < q$ . Fix a group  $G$  of order  $pq$ .

- (i) Prove that  $G$  is not simple.
- (ii) Further assume that  $q \not\equiv 1 \pmod{p}$ . Prove that  $G$  is cyclic.

**Problem 9.** Prove that there is no simple group of order  $p^2q$ , where  $p, q$  are prime numbers (not necessarily distinct).

**Problem 10.** Let  $a \in \{1, 2, \dots, p-1\}$  and  $k \in \mathbb{Z}_{\geq 1}$ . Prove that there is no simple group of order  $p^ka$ .

**Problem 11.** For each of the following numbers, prove that there is no simple group of that order: (a) 12; (b) 40; (c) 216.

**Problem 12.** Describe a Sylow 2-subgroup of the Dihedral group  $D_{10}$  of order 20.

**Problem 13.** Let  $G$  be a simple group of order 60 (assume it exists!). How many Sylow  $p$ -subgroups are there in  $G$ , for  $p = 2, 3$  and 5?

**Problem 14. (Key idea in Sylow's original proof of Thm (A))**

Consider  $H < G$  two groups, and fix a prime  $p > 0$  with  $p \mid |H|$ . Assume  $G$  has a Sylow  $p$ -group, called  $P$ . We aim to prove that the same is true for  $H$ . We write  $|H| = p^s m$  with  $(m : p) = 1$  and  $|G| = p^r m'$  with  $(m' : p) = 1$  and  $s \leq r$ .

- (i) Consider the left cosets  $X := G/P$ . Show that left multiplication defines an action of  $G$  on  $X$ .
- (ii) Consider the action of  $H$  on  $X$  inherited from the action on item (i). Show that there is a left coset  $gP$  whose  $H$ -orbit has size coprime to  $p$ .
- (iii) Show that  $p^s$  divides  $|\text{Stab}_H(gP)|$ .
- (iv) Show that  $\text{Stab}_G(gP) = gPg^{-1}$ . Conclude that  $|\text{Stab}_G(gP)| = |P|$  and, therefore,  $\text{Stab}_G(gP)$  is a Sylow  $p$ -subgroup of  $G$ .
- (v) Conclude from (iii) and (iv) that  $|\text{Stab}_H(gP)| = p^s$ , so it is a Sylow  $p$ -subgroup of  $H$ .

Sylow's original proof realizes  $G$  as a subgroup of  $S_n$  for  $n = |G|$  via the action of  $G$  on itself by left multiplication. In turn, given a prime  $p$  with  $p \mid n$ ,  $S_n$  is viewed as a subgroup of

$\mathrm{GL}_n(\mathbb{Z}/p\mathbb{Z})$  sending a permutation  $\sigma$  to the permutation matrix  $P_\sigma$ , where

$$(P_\sigma)_{ij} = \begin{cases} 1 & \text{if } \sigma(i) = j \\ 0 & \text{otherwise.} \end{cases}$$

By Problem 6,  $\mathrm{GL}_n(\mathbb{Z}/p\mathbb{Z})$  has a  $p$ -Sylow subgroup. Then, item (v) shows that the same is true for  $G$ .

**Problem (Bonus).** The objective of this problem is to prove the following stronger version of Sylow's Theorem (A).

**Theorem.** Given  $G$  of order  $n := p^r m$  with  $(m : p) = 1$ , we can find subgroups  $H_i < G$  with  $|H_i| = p^i$  for all  $i = 0, 1, \dots, r$ .

To prove the statement we will show that given  $i < r$  and a subgroup  $H$  with  $|H| = p^i$  we can find a  $p$ -subgroup  $H'$  of  $G$  with  $H \subset H'$  and  $(H' : H) = p$ . Then,  $H'$  will satisfy  $|H'| = p^{i+1}$ .

(i) Show that the normalizer  $N_G(H)$  satisfies  $H < N_G(H)$  and  $(N_G(H) : H) \equiv (G : H) \pmod{p}$  (*Hint:* Use the action of  $H$  on  $X := G/H$  by left multiplication.)

(ii) Show the group  $N_G(H)/H$  satisfies  $p \mid |N_G(H)/H|$ .

(iii) Assuming Sylow Theorem (A), show that  $N_G(H)/H$  has an element  $\sigma H$  of order  $p$ .

(iv) Conclude that  $H' = \langle \sigma, H \rangle$  has the desired properties.

*Note:* item (iii) can be proved without assuming Sylow Theorem (A). This result is known as *Cauchy's Theorem*: Every finite group  $G$  with  $p \mid |G|$  contains an element of order  $p$ . It can be proven for abelian groups by induction on the order of the group. In the general case, the result follows by working with either  $Z(G)$  (if  $p \mid |Z(G)|$ ) or a suitable centralizer

$$C_G(g) := \{h \in G : hgh^{-1} = g\}$$

where  $g \notin Z(G)$  and  $p$  does not divide the size of the conjugacy class of  $g$ .