

When are two Coxeter orbifolds diffeomorphic?

Michael W. Davis *

February 9, 2014

Abstract

One can define what it means for a compact manifold with corners to be a “contractible manifold with contractible faces.” Two combinatorially equivalent, contractible manifolds with contractible faces are diffeomorphic if and only if their 4-dimensional faces are diffeomorphic. It follows that two simple convex polytopes are combinatorially equivalent if and only if they are diffeomorphic as manifolds with corners. On the other hand, by a result of Akbulut, for each $n \geq 4$, there are smooth, contractible n -manifolds with contractible faces which are combinatorially equivalent but not diffeomorphic. Applications are given to rigidity questions for reflection groups and smooth torus actions.

AMS classification numbers. Primary: 57R18, 57R55, 20F55, 20F65

Secondary: 57S15, 57R91

Keywords: orbifold, manifold with corners, homology sphere, Coxeter orbifold, quasitoric manifold, polyhedral homology manifold

1 Introduction

More than once during the past few years¹ I have been asked, “Are combinatorially equivalent Coxeter orbifolds diffeomorphic?” By “Coxeter orbifold” the questioner means something like the fundamental polytope of a geometric reflection group. The underlying space of such an orbifold has the

*I thank the Mathematical Sciences Center at Tsinghua University for the opportunity to visit there during the spring of 2013. I began writing this paper during that period. I also was partially supported by NSF grant DMS 1007068.

¹Taras Panov and Mikiya Masuda asked me this at a 2010 conference on toric topology in Banff. More recently, Suhyoung Choi asked me the same question in connection with projective representations of Coxeter groups, cf. [5, Question 3.3].

structure of a manifold with corners. Usually, the questioner also wants to require that the underlying space is diffeomorphic, as a manifold with corners, to a simple convex polytope. In this context the answer to the question is affirmative although the proof is not obvious (see Wiemeler [28, Corollary 5.3] and Corollary 1.3, below). However, if one weakens the definition by only requiring the strata to be compact contractible manifolds, there are counterexamples (cf. Theorem 1.4). The problem is caused by the 4-dimensional strata.

An orbifold Q is *reflection type* if its local models are finite linear reflection groups on \mathbb{R}^n . (These were called “reflectofolds” in [11].) Since the orbit space of a finite linear reflection group is the product of a Euclidean space with a simplicial cone, a smooth orbifold of reflection type naturally has the structure of a smooth manifold with corners. One can label each codimension two stratum by an integer $m \geq 2$ to indicate that the local dihedral group along the stratum has order $2m$. So, Q is a smooth manifold with corners and the labeling is such that it determines a *finite* Coxeter group of rank k on each codimension k stratum. Conversely, if Q is a smooth manifold with corners with such a labeling on its codimension two strata, then it can be given the structure of a smooth orbifold of reflection type and this structure is unique up to isotopy (cf. [9, Section 17]). Let us suppose that each codimension k stratum of Q is the intersection of k codimension one strata (rather than, say, a component of such an intersection). There are several possibilities for when Q should be called a “Coxeter orbifold.” Here are three successively stronger definitions:

- (I) Each stratum of Q is a compact contractible manifold.
- (II) Each stratum of Q is homeomorphic to a disk.
- (III) Q is isomorphic (as a manifold with corners) to a simple convex polytope.

Any Q satisfying (I) is a *Coxeter orbifold*. It is *type* (II) or (III) if it satisfies the corresponding condition. When Q is smooth, note that if a stratum is homeomorphic to D^k , then, for $k \neq 4$, it is diffeomorphic to D^k , but, for $k = 4$, this implication is equivalent to the smooth 4-dimensional Poincaré Conjecture.

Under any of the above definitions, there is a simplicial complex dual to Q , called its *nerve* and denoted $N(Q)$. The vertices of $N(Q)$ are the codimension one strata of Q . A set of vertices spans a simplex of $N(Q)$ if and only if the intersection of the corresponding set of codimension one

faces is nonempty. For Q of type (III), $N(Q)$ is the boundary complex of the simplicial polytope that is dual to Q . For type (II), $N(Q)$ is a triangulation of S^{n-1} such that the link of each k -simplex of $N(Q)$ is homeomorphic to S^{n-k-2} . For a general Q of type (I), $N(Q)$ is a *generalized homology $(n-1)$ -sphere*, (or a GHS^{n-1} for short), where this means a polyhedral homology $(n-1)$ -manifold which has the same homology as S^{n-1} (cf. [10, p.192]). The labels on the codimension two strata of Q become labels on the edges of $N(Q)$. A *combinatorial equivalence* between Coxeter orbifolds Q and Q' is a label-preserving simplicial isomorphism $N(Q) \rightarrow N(Q')$.

The 1-skeleton of a simplicial complex N together with a labeling of its edges by integers ≥ 2 determines a Coxeter system (W, S) (cf. [10, Ex. 7.1.6]). The set of generators S can be identified with the vertex set of N . The labeling is *proper* if for each simplex σ of N the special subgroup $W_{S(\sigma)}$, generated by the vertices of σ , is finite. First, there is the question of existence. If N is a GHS^{n-1} with a proper labeling of its edges, is there a Coxeter orbifold Q with $N(Q) = N$? This was addressed in [9] where the following result was proved.

Theorem 1.1. ([9, Theorems 12.2 and 17.1]). *Suppose N is a GHS^{n-1} with a proper labeling of its edges. Then there is topological Coxeter orbifold Q with $N(Q) = N$. Noting that each 3-dimensional link in N is a PL homology 3-sphere (and therefore, has a unique smooth structure), we have that the orbifold Q admits a smooth structure if and only if each 3-dimensional link bounds a smooth contractible 4-manifold.*

The proof is based on the fact that every homology sphere of dimension $\neq 3$ has a smooth structure in which it smoothly bounds a contractible manifold (cf. Lemma 2.3 below).

This paper concerns the uniqueness question: if two Coxeter orbifolds are combinatorially equivalent, are they isomorphic (as orbifolds)? In the topological category the answer is yes. This is a consequence of the facts that the Poincaré Conjecture and the topological h-cobordism Theorem hold in all dimensions.

In Section 4 (Definition 4.1) we define what it means for two combinatorially equivalent, smooth Coxeter orbifolds to have “diffeomorphic 4-dimensional faces.” Our goal is the following theorem.

Theorem 1.2. *Suppose Q and Q' are combinatorially equivalent, smooth Coxeter orbifolds with diffeomorphic 4-dimensional faces. Then Q and Q' are diffeomorphic.*

Corollary 1.3. (Wiemeler [28]). *Combinatorially equivalent Coxeter orbifolds of type (III) are diffeomorphic.*

On the other hand, according to the following theorem, without the hypothesis of diffeomorphic 4-dimensional faces, Theorem 1.2 is false. As we will explain in Section 4, this is a consequence of a result of Akbulut [1].

Theorem 1.4. *In each dimension $n \geq 4$, there are smooth Coxeter orbifolds that are combinatorially equivalent but not diffeomorphic. Moreover, these orbifolds can be chosen to be aspherical.*

The questions we are dealing with in Theorems 1.1, 1.2 and 1.4 have nothing to do with the labelings of the codimension two strata of the reflection type orbifold, rather we are only concerned with its underlying structure as a manifold with corners – the Coxeter group is extraneous. For this reason, we will reformulate our results in what follows in terms of “resolutions” of cones on generalized homology spheres by “contractible manifolds with contractible faces.”

Sections 5 and 6 give applications of Theorems 1.2 and 1.4 to the theory of reflection groups and to the theory of locally standard torus actions. Mikiya Masuda suggested that I add to the original version of this paper some details in the section on torus actions concerning smoothness questions. On the basis of his very helpful comments, I have added Propositions 6.2, 6.4 and 6.8 which address questions of equivariant diffeomorphism versus equivariant homeomorphism.

After completing the first version of this paper, I learned of the recent work [28] by M. Wiemeler on torus manifolds, with which this paper has substantial overlap. In particular, Wiemeler proves the result for simple polytopes (Corollary 1.3), as well as its application in Section 6 to quasitoric manifolds of type (III) (Theorem 6.6). My thanks also go to the referee for finding some minor errors.

2 Existence

Manifolds with faces. A topological n -manifold with boundary is a *smooth manifold with corners* if it is locally differentiably modeled on the simplicial cone, $[0, \infty)^n \subset \mathbb{R}^n$. If $(x_1, \dots, x_n): U \rightarrow [0, \infty)^n$ are local coordinates in a neighborhood of a point x , then the number $c(x)$ of indices i with $x_i = 0$, is independent of the chart. A component of $\{x \mid c(x) = m\}$ is a *pure stratum* of codimension m ; a *stratum* is the closure of a pure stratum. One also can define the notion of a *topological manifold with corners* by requiring

the overlap maps to be strata-preserving (cf. [10, p. 180]). A manifold with corners is *nice* if each stratum of codimension two is contained in the closure of exactly two strata of codimension one. Niceness implies that the closure of a codimension k -stratum is also a manifold with corners.

A nice manifold with corners P will be called a *manifold with faces*. A *face* is an intersection of codimension one strata. The k -*skeleton* of P is the union of faces of dimension $\leq k$. Two manifolds with faces, P and P' , are *topologically isomorphic* if there is a strata-preserving homeomorphism $P \rightarrow P'$. (Of course, a diffeomorphism of smooth manifolds with corners is automatically strata-preserving.) Parallel to our earlier definitions in the Introduction, consider the following conditions on a manifold with faces P :

- (I) Each face of P is a compact contractible manifold.
- (II) Each face of P is homeomorphic to a disk.
- (III) P is isomorphic to a simple convex polytope.

In case (I), P is a *contractible manifold with contractible faces*. In case (II), it is a *cell with cellular faces*. If we replace the word “contractible” by “acyclic,” we get the notion of an *acyclic manifold with acyclic faces*.

One defines $N(P)$, the *nerve* of P , as before: it is the simplicial complex with vertex set the set of codimension one faces and with a $k - 1$ -simplex for each face of codimension k . Two manifolds with faces, P and P' , are *combinatorially equivalent* if there is a simplicial isomorphism $N(P) \rightarrow N(P')$. If P is an acyclic n -manifold with acyclic faces, then $N(P)$ is a GHS^{n-1} .

Example 2.1. A simple convex polytope is naturally a smooth cell with cellular faces. More generally, if N is a triangulation of S^{n-1} and if the link of each k -simplex is homeomorphic to S^{n-k-2} (e.g., if the triangulation is PL), then $\text{Cone}(N)$ has a dual cell structure giving it the structure of a (topological) cell with cellular faces.

Resolutions. Suppose N is a GHS^{n-1} . A *resolution* of $\text{Cone}(N)$ is a contractible manifold with contractible faces P such that $N(P) = N$. To define *acyclic resolution* replace the word “contractible” by “acyclic.” A resolution is *smooth* if P is a smooth manifold with corners. (The use of the term “resolution” will be discussed in Remark 2.4 below.)

The following existence result essentially was proved in [9]. It implies Theorem 1.1 of the Introduction.

Theorem 2.2. (cf. [9, Thm. 17.2]). *Suppose N is a GHS^{n-1} . Then the following statements are true.*

- (1) *Cone(N) has a topological resolution.*
- (2) *Cone(N) has a smooth acyclic resolution.*
- (3) *Cone(N) has a smooth resolution if and only if each of its 3-dimensional links bounds a smooth (or PL) contractible 4-manifold.*

We will recall the proof of this theorem at the end of this section since it is based on the same obstruction theory which we will need for the uniqueness result.

Homology spheres and homotopy spheres. Let Θ_k be the Kervaire-Milnor group of h -cobordism classes of smooth structures on S^k (cf. [17]). Let Θ_k^H be the group of homology cobordism classes of smooth homology k -spheres. The next lemma is well-known.

- Lemma 2.3.** (1) *Any homology k -sphere is the boundary of a topological contractible $(k+1)$ -manifold.*
- (2) *For $k \neq 3$, any PL homology k -sphere is the boundary of a PL contractible $(k+1)$ -manifold.*
- (3) *Suppose M^k is a smooth homology k -sphere. For $k \geq 4$, there is a smooth homotopy k -sphere Σ^k , possibly with an exotic smooth structure, such that $M^k \# \Sigma^k$ bounds a smooth contractible $(k+1)$ manifold. (Here $\#$ is used for the connected sum operation.)*

Statement (2) is not true for every homology 3-sphere because of Rokhlin's Theorem.

Sketch of proof of Lemma 2.3. We note that special arguments are needed to prove (1) when $k = 3$ or 4. We first prove (3). Suppose $k > 4$ and that M^k is smooth. As in [16, Theorem 5.6] one can add 1- and 2-handles to $M^k \times [0,1]$ to get a simply connected, homology cobordism W from M^k to a smooth homotopy k -sphere. If Σ^k denotes this homotopy sphere with orientation reversed, then $M^k \# \Sigma^k$ bounds a contractible manifold. When $k = 4$, one argues that M^4 bounds a framed manifold W^5 (since it represents 0 in framed bordism) and then that one can do surgery to make W^5 contractible. This proves (3). Since every PL homology sphere is smoothable, it also proves (2).

As for (1), when $k = 4$, it is not known if M^4 admits a PL structure. However, since we can do simply connected topological surgery, we can add 1- and 2-handles to $M^4 \times [0, 1]$ as before to get a simply connected homology cobordism W from M^4 to a homotopy 4-sphere, and by Freedman's proof of the Poincaré Conjecture in [13] this is homeomorphic to the standard S^4 . Gluing on D^5 gives the desired contractible 5-manifold. When $k = 3$, Freedman [13] proved that (1) holds in the topological category. \square

Remark 2.4. (On resolutions). A *polyhedral homology m -manifold* means a simplicial complex N such that the link of any k -simplex has the same homology as S^{m-k-1} . The “dual cell” of such a k -simplex in N is the cone on its link, i.e., it is a contractible polyhedral homology $(m - k)$ -manifold with boundary. General resolutions for polyhedral homology manifolds (rather than just for generalized homology spheres) were considered by Cohen [6] and Sullivan [27] in the early 1970s. They developed an obstruction theory for finding a resolution of N by a manifold by replacing each dual cell by an acyclic manifold with boundary. One proceeds by induction on the dimension of the dual cell. By Lemma 2.3 (2), the only obstruction to finding a PL acyclic resolution of a polyhedral homology manifold N lies in $H^4(N; \Theta_3^H)$. A version of this obstruction theory is used in the proof of Theorem 2.2 which we sketch below. In the topological category, it follows from Lemma 2.3 (1) that contractible resolutions always exist.

Sketch of proof of Theorem 2.2. We build the faces of P by induction on dimension. Start with a 0-dimensional face for each $(n - 1)$ -simplex of N . Assume by induction that we have constructed a face F_τ for each simplex τ of codimension $< k$ and let σ be a simplex of codimension k . Define $\partial F_\sigma := \bigcup_{\tau > \sigma} F_\tau$ where the union is over all $\tau > \sigma$ of codimension $(k - 1)$ in N . Then ∂F_σ is a homology $(k - 1)$ -sphere and the problem is to fill it in with a k -dimensional contractible face. By Lemma 2.3 (1) we can always do this in the topological category; hence, statement (1).

To get a smooth resolution one encounters a problem when trying to construct the 4-dimensional faces. If each face of dimension ≤ 3 is a disk, then the boundary of a potential 4-dimensional face is dual to the corresponding 3-dimensional link. So, the condition that this homology 3-sphere smoothly bounds a contractible manifold is sufficient for constructing the 4-dimensional face. *A priori*, this condition might not be necessary since the homology 3-sphere would be indeterminate if 3-dimensional faces could be fake 3-disks. However, since the 3-dimensional Poincaré Conjecture is true, fake 3-disks do not exist. Therefore, a smooth 4-dimensional face exists if

and only if the homology 3-sphere bounds a contractible 4-manifold. For $k \geq 4$, the obstruction to filling in the smooth $(k + 1)$ -dimensional strata lies in $H^{k+1}(\text{Cone}(N); \Theta_k)$. This group is 0 since $\text{Cone}(N)$ is acyclic. This proves (3).

What about (2)? As explained in [9, pp. 322–323], there is an obstruction cocycle to filling in the boundaries of potential 4-dimensional strata with smooth acyclic 4-manifolds. This cocycle takes values in Θ_3^H . However, this cocycle is indeterminate, since we can alter our construction of 3-dimensional faces by taking connected sum with elements of Θ_3^H . In other words, the cocycle can be varied by a coboundary giving us a well-defined obstruction in $H^4(\text{Cone}(N); \Theta_3^H) = 0$. After filling in the 4-dimensional faces, we can continue as in the proof of (3) to fill in the faces of higher dimension. \square

Uniqueness in the topological category. The next lemma follows from the Poincaré Conjecture and the h-cobordism Theorem.

Lemma 2.5. *Suppose C and C' are compact contractible k -manifolds and that $\varphi : \partial C \rightarrow \partial C'$ is a homeomorphism, then φ extends to a homeomorphism $\Phi : C \rightarrow C'$.*

Suppose P and P' are topological contractible n -manifolds with contractible faces and $N(P) = N(P') = N$. It follows from Lemma 2.5 that P and P' are topologically isomorphic as manifolds with corners. In other words, *the topological resolution of $\text{Cone}(N)$ is unique.*

3 More facts about homology spheres and contractible manifolds

Pseudo-isotopies on S^n . Given a smooth manifold M and M' , denote by $\text{Diff}(M, M')$ the set of diffeomorphisms from M to M' . Let $\text{Diff}(M) := \text{Diff}(M, M)$, be the topological group of self-diffeomorphisms of M . If M is oriented, $\text{Diff}^+(M)$ denotes the subgroup of orientation-preserving self-diffeomorphisms. If M is a manifold with boundary, then $\text{Diff}_\partial(M)$ is the subgroup of self-diffeomorphisms which are the identity on ∂M .

Let Γ_{n+1} denote the group of pseudo-isotopy classes of orientation preserving, self diffeomorphisms of S^n . Let $\iota : \text{Diff}(D^{n+1}) \rightarrow \text{Diff}(S^n)$ be the natural homomorphism which takes $\varphi \in \text{Diff}(D^{n+1})$ to $\varphi|_{S^n}$. Here are three other equivalent definitions of Γ_{n+1} :

- (i) $\Gamma_{n+1} = \text{Diff}(S^n)/\iota(\text{Diff}(D^{n+1}))$. In other words, a diffeomorphism $\varphi \in \text{Diff}(S^n)$ represents 0 in Γ_{n+1} if and only if it extends to a self-diffeomorphism of D^{n+1} .
- (ii) $\Gamma_{n+1} = \pi_0(\text{Diff}^+(S^n))$, the group of isotopy classes of orientation preserving diffeomorphisms of S^n .
- (iii) $\Gamma_{n+1} = \Theta_{n+1}$, the abelian group (under connected sum) of h -cobordism classes of smooth structures on S^{n+1} . (It is a coincidence that $\Theta_4 = 0 = \Gamma_4$.)

Version (i) is obvious. As for (ii), in dimension 4 Cerf [2] proved that $\Gamma_4 = 0$ and then Hatcher [14] showed that $\pi_0(\text{Diff}^+(S^3))$ is also 0. Cerf [3] also showed that for simply connected manifolds of dimension ≥ 5 , pseudo-isotopic diffeomorphisms are isotopic. Hence, for $n \geq 5$, $\Gamma_{n+1} = \pi_0(\text{Diff}^+(S^n))$. It follows that Γ_{n+1} is a finite abelian group.

The next lemma is well-known and not difficult to prove.

Lemma 3.1. *Let $f \in \text{Diff}^+(S^n)$. Let D' be a closed, smooth n -disk in S^n and let D^n be the closure of complementary n -disk. Then f is isotopic to a diffeomorphism $f' \in \text{Diff}^+(S^n)$ such that $f'(D') = D'$ and $f'|_{\partial D'}$ is the identity. Consequently, any element of Γ_{n+1} can be represented by a diffeomorphism in $\text{Diff}_{\partial}(D^n)$.*

Contractible manifolds and homology spheres. The next two lemmas are well-known consequences of the smooth h -cobordism Theorem.

Lemma 3.2. (cf. [18, Theorem on p.183]). *Suppose C and C' are smooth, compact contractible $(n+1)$ -manifolds with homeomorphic boundaries. If $n \neq 3$ or 4, then C and C' are diffeomorphic. In particular, if $n \neq 4$, ∂C and $\partial C'$ are diffeomorphic.*

Proof. Suppose $\varphi : \partial C \rightarrow \partial C'$ is a homeomorphism. Then $C \cup_{\varphi} C'$ is homeomorphic to S^{n+1} . Hence, it has a smooth structure in which it is diffeomorphic to S^{n+1} . The union of a collared neighborhood of ∂C in C with a collared neighborhood of $\partial C'$ in C' , gives a smooth h -cobordism from ∂C to $\partial C'$; hence, for $n \neq 4$, they are diffeomorphic. Filling in $C \cup_{\varphi} C'$ with D^{n+2} we get a smooth h -cobordism from C to C' which restricts to the given one from ∂C to $\partial C'$. \square

Lemma 3.3. *Suppose $n > 4$. Let M be a homology n -sphere and let $\Theta_n(M)$ be the set of oriented diffeomorphism classes of smooth structures on M .*

Then the action of Θ_n on $\Theta_n(M)$ by connected sum is simply transitive. In particular, if M_0 denotes the “standard” smooth structure on M induced from the contractible manifold which it bounds, then there is a bijection $\Theta_n \rightarrow \Theta_n(M)$ defined by $\Sigma \mapsto M_0 \# \Sigma$.

Proof. The induced action of Θ_n on Θ_n^H is simply transitive (cf. [18, p. 183]). \square

Suppose that M and M' are smooth, closed n -manifolds, that $D^n \subset M$ and $D^n \subset M'$ are embedded disks and that $g \in \text{Diff}(M, M')$ is such that $g|_{D^n} = \text{id}_{D^n}$. Given $f \in \text{Diff}_\partial(D^n)$, let $f \cdot g : M \rightarrow M'$ be the diffeomorphism defined by extending f via $g|_{M-D^n}$. When $M' = M$ and $g = \text{id}_M$, this gives a homomorphism $f \mapsto f \cdot \text{id}$ from $\text{Diff}_\partial(D^n)$ to $\text{Diff}(M)$ which descends to a homomorphism $\lambda : \Gamma_{n+1} \rightarrow \pi_0(\text{Diff}(M))$.

Next suppose A, A' are smooth compact, acyclic $(n+1)$ -manifolds with boundary with $n+1 > 4$ and that $\partial A = \partial A'$. We want to define left inverse for the action of Γ_{n+1} on $\pi_0(\text{Diff}(\partial A, \partial A'))$. Suppose $\varphi : \partial A \rightarrow \partial A'$ is a diffeomorphism. Then $A \cup_\varphi A'$ is a smooth homology $(n+1)$ -sphere. By Lemma 3.3, there is an exotic sphere $\Sigma \in \Theta_{n+1}$ so that $(A \cup_\varphi A') \# (-\Sigma)$ bounds a contractible manifold. Hence, we can find another diffeomorphism $\varphi_0 : \partial A \rightarrow \partial A'$ so that $(A \cup_{\varphi_0} A')$ bounds this contractible manifold, i.e.,

$$(A \cup_\varphi A') = (A \cup_{\varphi_0} A') \# \Sigma.$$

So, for $n+1 > 4$, we get a well-defined element $\gamma(\varphi)$ in $\Theta_{n+1} = \Gamma_{n+1}$,

$$\gamma(\varphi) = \Sigma. \tag{3.1}$$

From its definition, γ is a left inverse for the action of Γ_{n+1} defined by λ . Indeed, if $f \in \text{Diff}_\partial(D^n)$ and $\lambda(f) = f \cdot \varphi_0 : \partial A \rightarrow \partial A$, then since $\lambda(f)$ is supported on an n -disk, $(A \cup_{\lambda(f)} A') = (A \cup_{\varphi_0} A') \# \Sigma(f)$, where $\Sigma(f) \in \Theta_{n+1}$ is the exotic sphere corresponding to f , i.e., $\gamma(\lambda(f))$ is the class of f in Γ_{n+1} .

Remark 3.4. In dimension 3, we have $\Gamma_3 = 0 = \Theta_3$, and $\Theta_3^H \neq 0$. In dimension 4, $\Gamma_4 = \Theta_4^H = 0$, and Θ_4 is unknown. So, when $n+1 = 3$ or 4, if A and A' are contractible, then $A \cup_\varphi A'$ bounds a contractible manifold and one can require $\gamma(\varphi)$ to lie in Γ_{n+1} (which is $= 0$).

We will sometimes identify $A \cup_\varphi A'$ with the smooth homology sphere:

$$H(\varphi) := A \cup M_\varphi \cup A', \tag{3.2}$$

where M_φ is the mapping cylinder of the diffeomorphism $\varphi : \partial A \rightarrow \partial A'$.

Since A is acyclic, so is $A \times [0, 1]$. Hence, when $A' = A$ we have that $H(\text{id}) = \partial(A \times [0, 1])$ represents 0 in Θ_{n+1}^H . So, if φ extends to a diffeomorphism $\Phi : A \rightarrow A'$, we get a diffeomorphism from $A \cup_{\text{id}} A$ to $A \cup_{\varphi} A'$. In other words, if φ extends to a diffeomorphism, then $\gamma(\varphi) = 0$. For $n + 1 > 4$ and for A and A' contractible, the converse is a consequence of the h-cobordism Theorem as we show in the next lemma.

Lemma 3.5. *Suppose C and C' are smooth, compact contractible $(n + 1)$ -manifolds and that $\varphi : \partial C \rightarrow \partial C'$ is a diffeomorphism.*

- (1) $C \cup_{\varphi} C'$ smoothly bounds a contractible $(n + 2)$ -manifold if and only if $\gamma(\varphi) = 0$.
- (2) For $n + 1 \neq 4$, φ extends to a diffeomorphism $\Phi : C \rightarrow C'$ if and only if $\gamma(\varphi) = 0$.

Proof. (1) When $n + 1 = 3$, the 3-dimensional Poincaré Conjecture (cf. [23]) implies that $C \cup_{\varphi} C' = S^3 = \partial D^4$. For $n + 1 = 4$, by Lemma 2.3 (3), any smooth structure on S^4 bounds a contractible 5-manifold. For $n + 1 > 4$, by Lemma 3.3, $\gamma(\varphi) = 0$ if and only if $C \cup_{\varphi} C'$ bounds a contractible manifold.

(2) As explained in the paragraph preceding this lemma, if φ extends to Φ , then $\gamma(\varphi) = 0$. Conversely, suppose $\gamma(\varphi) = 0$. Then, since $H(\varphi) = S^{n+1}$, we can fill in $H(\varphi)$ with an $(n + 2)$ -disk to obtain an h-cobordism W from C to C' . By the h-cobordism Theorem (which is true for $n + 1 \neq 4$), there is a diffeomorphism $\Phi : C \times [0, 1] \rightarrow W$ such that $\Phi(C \times 0) = C$, $\Phi(C \times 1) = C'$, and $\Phi(\partial C \times [0, 1]) = M_{\varphi}$. Moreover, $\Phi|_{C \times 0} = \text{id}$ and $\Phi|_{\partial C \times [0, 1]}$ is the natural identification onto M_{φ} . Thus, $\Phi|_{\partial C \times 1} = \varphi$ and $\Phi|_{C \times 1}$ is the desired extension. \square

In the next section we will need the following lemma to establish that an obstruction cochain is a cocycle.

Lemma 3.6. *Let E be a smooth, compact contractible $(n + 1)$ -manifold and let F_1, \dots, F_m be disjoint compact contractible submanifolds of codimension 0 in ∂E . Let $X^n = \partial E - \bigcup_{i=1}^m F_i^{\circ}$, where F_i° denotes the interior of F_i . Suppose $f : X^n \rightarrow X^n$ is a diffeomorphism which takes each ∂F_i to itself. Let $\gamma_i \in \Gamma_{n+1}$ be the obstruction to extending $f|_{\partial F_i}$ to F_i . Then $\gamma_1 + \dots + \gamma_m = 0$.*

Proof. Choose disjoint embedded paths (“edges”) connecting the ∂F_i so that the union of the edges with the F_i is simply connected (giving a “tree of contractible n -manifolds” in ∂E). We can arrange that f restricts to the identity map on a tubular neighborhood of each edge. Let A denote

the complement of these neighborhoods of edges in X^n . In other words, A is the complement of a regular neighborhood of the tree of contractible n -manifolds. It follows that A is a compact acyclic n -manifold, that ∂A is the connected sum, $\partial F_1 \# \cdots \# \partial F_m$ and that f takes A to itself. The restriction of f to ∂A represents the element $\gamma_1 + \cdots + \gamma_m \in \Gamma_{n+1}$. Since f extends over A , the paragraph preceding Lemma 3.5, implies that this element is 0. \square

Remark 3.7. For $n + 1 = 4$, Lemma 3.5 (2) is unknown even when C and C' are both homeomorphic to D^4 . (This is equivalent to the smooth 4-dimensional Poincaré Conjecture.) When ∂C and $\partial C'$ are allowed to be homology 3-spheres with nontrivial fundamental groups, Lemma 3.5 (2) does not hold. Indeed, Akbulut [1] has shown that there exist compact contractible smooth 4-manifolds C and C' with $\partial C = \partial C'$, such that C and C' are not diffeomorphic rel ∂C (cf. Theorem 4.4 below).

4 Uniqueness

A k -dimensional face F of an n -dimensional manifold with faces has a tubular neighborhood of the form $F \times [0, \infty)^{n-k}$, where $[0, \infty)^{n-k}$ denotes the standard simplicial cone in \mathbb{R}^{n-k} . This is an easy consequence of the Colored Neighborhood Theorem for manifolds with boundary. Given a smooth n -dimensional manifold with faces P , let $\hat{\partial}P$ denote its topological boundary. (N.B. We write $\hat{\partial}P$ instead of ∂P to indicate that the corners have not been rounded.) The tubular neighborhoods of the faces can be fit together compatibly to give a neighborhood $e(P)$ of $\hat{\partial}P$. If P is smooth, we can push $\hat{\partial}P$ into the interior of $e(P)$ to get a smooth manifold ∂P (∂P is $\hat{\partial}P$ with corners rounded). The manifold ∂P separates P into two pieces, one is a neighborhood of $\hat{\partial}P$ and the other is homeomorphic to P . It should not cause confusion to continue to denote the second piece by P .

Suppose that N is a GHS^{n-1} and that P, P' are smooth contractible manifolds with contractible faces with $N(P) = N(P') = N$. For each $\sigma \in N$, let F_σ and F'_σ be the corresponding faces of P and P' . We will try to construct a diffeomorphism $\varphi : P \rightarrow P'$ one face at a time. Since the faces of dimension ≤ 3 are cells, we can define φ on a neighborhood of the 3-skeleton of P . It is unique up to isotopy. Suppose σ is a simplex of codimension 4. Then we have a diffeomorphism $\varphi : \partial F_\sigma \rightarrow \partial F'_\sigma$ (which we can assume is the identity). The problem is to extend it. According to Remark 3.7, Lemma 3.5 (2) does not apply. This leads us to the following.

Definition 4.1. Suppose, as above, P and P' are smooth contractible manifolds with contractible faces and $N(P) = N(P') = N$. Then P and P' have *diffeomorphic 4-dimensional faces* if for each simplex σ of codimension 4 in N , F_σ and F'_σ are diffeomorphic rel boundary.

Our goal is to prove the following, which is equivalent to Theorem 1.2.

Theorem 4.2. *Suppose P and P' are smooth contractible manifolds with contractible faces and $N(P) = N(P') = N$. Then P and P' are diffeomorphic if and only if they have diffeomorphic 4-dimensional faces.*

Once we have constructed a φ taking a $(k-1)$ -face F to the corresponding $(k-1)$ -face F' of P' , we extend it to a map of tubular neighborhoods, $F \times [0, \infty)^{n-k} \rightarrow F' \times [0, \infty)^{n-k}$, which is linear on the $[0, \infty)^{n-k}$ factors. So, suppose φ has been defined on the $(k-1)$ -skeleton of P and that we want to extend it to a map on a k -face, $F \rightarrow F'$. Since we have extended to tubular neighborhoods, we have a diffeomorphism $\varphi : e(F) \rightarrow e(F')$ restricting to a diffeomorphism $\partial F \rightarrow \partial F'$. The problem is to extend this to a diffeomorphism $F \rightarrow F'$. There might be an obstruction.

Proof of Theorem 4.2. The faces of P and P' of dimension ≤ 2 are standard cells. Since the 3-dimensional Poincaré Conjecture is true (cf. [23], [22]), the 3-dimensional faces are also standard. It follows that we can define φ on a neighborhood of the 3-skeleton and this definition is unique up to isotopy. The hypothesis that the 4-dimensional faces of P and P' are diffeomorphic implies that φ can be extended over a neighborhood of the 4-skeleton.

Suppose, by induction, that we have constructed φ on a regular neighborhood of the $(k-1)$ -skeleton, with $k-1 \geq 4$. One needs to choose orientations for the faces of P in order to define the cellular cochains on P . Let F be an oriented k -face. Then $\varphi|_{\partial F} : \partial F \rightarrow \partial F'$ is defined. As in the paragraph preceding Lemma 3.5 there is an element $c(F) \in \Gamma_k$ defined by

$$c(F) = \gamma(\varphi) = [F \cup_\varphi F'] \in \Theta_k = \Gamma_k.$$

By Lemma 3.5 (2), $\varphi|_{\partial F}$ extends across F if and only if $c(F)$ vanishes. The assignment $F \mapsto c(F)$ is a cellular cochain $c \in C^k(P; \Gamma_k)$. It is the obstruction to extending φ across the k -skeleton. We have to check two points in order for this obstruction theory to work:

- (1) We are free to vary the construction of φ on the $(k-1)$ -skeleton. If we fix φ on the $(k-2)$ -skeleton and vary its extension over the $(k-1)$ -faces, then c should change by a coboundary.

(2) c should be a cocycle.

To check (1) suppose G is an oriented $(k-1)$ -face and $\gamma \in \Gamma_k$. Let $d_{G,\gamma} \in C^{k-1}(P; \Gamma_k)$ be the cochain which assigns γ to G and 0 to all other $(k-1)$ -faces. Next we want to alter φ on the $(k-1)$ -skeleton by changing $\varphi|_G$ by the element γ (where γ is thought of as a diffeomorphism of D^{k-1} which is the identity on ∂D^{k-1}). Denote the new map on the $(k-1)$ -skeleton by ψ . So, we have two cochains $c_\varphi, c_\psi \in C^k(P; \Gamma_k)$. How are they related? Let F be an oriented k -face such that G occurs in $\hat{\partial}F$ with coefficient ± 1 . Denote this coefficient by $[G:F]$ (and call it the *incidence number*). Then c_φ and c_ψ are related by the formula

$$c_\psi(F) = c_\varphi(F) + [G:F] d_{G,\gamma}.$$

That is to say, $c_\psi = c_\varphi + \delta(d_{G,\gamma})$, where $\delta : C^{k-1}(P; \Gamma_k) \rightarrow C^k(P; \Gamma_k)$ is the coboundary map. This establishes (1).

As for (2) we want to check that c ($= c_\varphi$) is a cocycle. Let E be a $(k+1)$ -face and let F_1, \dots, F_m the k -faces of $\hat{\partial}E$, oriented by outward-pointing normals. (This means that all incidence numbers $[F_i : E]$ are $= 1$.) Apply Lemma 3.6 with X a regular neighborhood in E of the $(k-1)$ -skeleton of E and with $\gamma_i = c(F_i)$ to get:

$$\delta(c)(E) = c(\hat{\partial}E) = \sum_{i=1}^m c(F_i) = 0.$$

Hence, c is a cocycle.

We use (1) and (2) to complete the proof. Suppose $c \in C^k(P; \Gamma_k)$ is the obstruction cochain. Since c is a cocycle and P is acyclic, we have $c = \delta(d)$ for some $d \in C^{k-1}(P; \Gamma_k)$. If we use $-d$ to alter φ on the $(k-1)$ -skeleton, the new c will be identically 0. Hence, we can extend over the k -skeleton. \square

Nonuniqueness. In the language of resolutions, Theorem 1.4 can be restated as follows.

Theorem 4.3. *For each $n \geq 4$ there is a generalized homology $(n-1)$ -sphere N with two distinct smooth resolutions.*

This is more or less an immediate consequence of the following important result of Akbulut [1].

Theorem 4.4. (Akbulut [1]). *There exist two smooth, compact contractible 4-manifolds C and C' with the same boundary so that C and C' are not diffeomorphic rel boundary.*

In Akbulut's construction C is the Mazur manifold. (This was the first example of a compact, contractible 4-manifold whose boundary is a homology sphere that is not simply connected.)

Here are a few remarks about the proof of 4.3. First suppose $n = 4$. Let N be a triangulation of the boundary of the Mazur manifold. The dual cell structure gives a resolution of N . We can get two different smooth resolutions P and P' of $\text{Cone}(N)$ by filling in C and C' , respectively. Since the diffeomorphism $\partial P \rightarrow \partial P'$ is isotopic to the identity, it cannot be extended to a diffeomorphism $P \rightarrow P'$.

For $n > 4$, let N be any GHS^{n-1} which has ∂C as the link of a simplex of codimension 4. For example, one could take N to be the join $\partial C * S^{n-5}$ (or any subdivision of this). By filling in each 4-dimensional face with copy of C or C' one obtains distinct resolutions.

5 Reflection groups

Smooth equivariant rigidity of reflection groups. Associated to a Coxeter orbifold Q , there is a Coxeter system (W, S) , where the set of generators S corresponds to the set of codimension one faces of Q . The group W is the orbifold fundamental group of Q . If $L(W, S)$ denotes the nerve of the Coxeter system, then properness of the labeling means that $N(Q) \subset L(W, S)$. If $L(W, S) = N(Q)$, then the universal cover \tilde{Q} of Q is contractible (cf. [9] or [10]), i.e., Q is aspherical. Since each face of a Coxeter orbifold is contractible, it follows that $\tilde{Q} = \underline{EW}$, the universal space for proper W -actions and hence, that the fixed point set of any spherical special subgroup is a contractible submanifold of \tilde{Q} .

Now suppose Q' is another aspherical Coxeter orbifold with associated Coxeter system (W', S') and with $W \cong W'$. By the universal properties of \underline{EW} and \underline{EW}' , there is a W -homotopy equivalence $\tilde{Q} \rightarrow \tilde{Q}'$. It is proved in [4] that $L(W, S)$ and $L(W', S')$ are isomorphic, i.e., that Q and Q' are combinatorially equivalent. From this we get the rigidity theorem of Prassidis-Spieler in [24]: \tilde{Q} and \tilde{Q}' are W -equivariantly homeomorphic. When combined with Theorem 1.2, this gives the following.

Theorem 5.1. (cf. Prassidis-Spieler [24]). *Suppose Q and Q' are aspherical Coxeter orbifolds with isomorphic Coxeter groups. Then \tilde{Q} and \tilde{Q}' are equivariantly diffeomorphic if and only if corresponding 4-dimensional faces of Q and Q' are diffeomorphic.*

In other words, the equivariant, smooth version of the Borel Conjecture

holds for aspherical Coxeter orbifolds if and only if their 4-dimensional faces are diffeomorphic.

Remark 5.2. If we only assume that the orbifold Q'' is the quotient of a cocompact W -action on a contractible manifold, then Q'' will still be a manifold with faces; however, the proper faces need only be acyclic. If at least one face is not contractible, then Q'' will not be homeomorphic to Q . (There is an equivariant map $\tilde{Q}'' \rightarrow \underline{E}W$ which is a homotopy equivalence; however, it will not be a W -homotopy equivalence.)

Theorem 1.4 yields the following.

Theorem 5.3. *For each $n \geq 4$, there are n -dimensional Coxeter orbifolds Q and Q' which are combinatorially equivalent but not diffeomorphic (hence, Q and Q' are not equivariantly diffeomorphic).*

There is no problem in arranging for Q and Q' to be aspherical. For example, if N is a GHS^{n-1} discussed in the last two paragraphs of the previous section, we can insure asphericity by replacing N by its barycentric subdivision and labeling all edges 2.

The following question remains open.

Question. *Are there fake smooth closed aspherical 4-manifolds?*

If Q and Q' are aspherical 4-dimensional orbifolds as in Theorem 5.3, then by passing to a subgroup of finite index in the Coxeter group, we obtain smooth aspherical 4-manifolds M and M' with the same fundamental group. They might or might not be diffeomorphic. A variation of this would be to change only one chamber of M from C to C' . It seems plausible that some such construction could yield homeomorphic but not diffeomorphic aspherical manifolds.

6 Torus actions

The moment angle manifold. There is a standard linear action of the m -torus T^m on \mathbb{C}^m . The orbit space is \mathbb{R}_+^m , where $\mathbb{R}_+ = [0, \infty)$. The orbit map $p : \mathbb{C}^m \rightarrow \mathbb{R}_+^m$ can be defined by $p(z_1, \dots, z_m) = (|z_1|^2, \dots, |z_m|^2)$.

Suppose P is an n -dimensional smooth manifold with faces with m faces of codimension one (“facets”), F_1, \dots, F_m . Let $f : P \rightarrow \mathbb{R}_+^m$ be a map such that the inverse image of the coordinate hyperplane $x_i = 0$ is F_i . Moreover, f is required to be transverse to each of these hyperplanes. Next we will construct a smooth $(n + m)$ -manifold Z_P with a smooth T^m -action and

with orbit space P . The *moment-angle manifold* Z_P corresponding to P is defined by the pullback diagram:

$$\begin{array}{ccc} Z_P & \longrightarrow & \mathbb{C}^m \\ \downarrow & & \downarrow p \\ P & \xrightarrow{f} & \mathbb{R}_+^m \end{array} \quad (6.1)$$

In other words, $Z_P = f^*(\mathbb{C}^m) := \{(x, z) \in P \times \mathbb{C}^m \mid f(x) = p(z)\}$. By the transversality hypothesis, Z_P inherits the structure, as a subset of $P \times \mathbb{C}^m$, of a smooth $(n + m)$ -manifold. Moreover, the T^m -action on the second factor induces a smooth T^m -action on Z_P . As we will see in Proposition 6.2 below, up to an equivariant diffeomorphism inducing the identity on P , the moment angle polytope is independent of the choice of f .

The T^m -action on Z_P is *modeled on the standard representation* in the sense that its orbit types and normal representations occur among those of T^m on \mathbb{C}^m . The principal orbits (i.e., the maximal orbits) are isomorphic to T^m . The principal orbit bundle is trivial (since it is the pullback of the principal orbit bundle for T^m on \mathbb{C}^m).

Remarks 6.1. For G a compact Lie group, the question of pulling back smooth G -manifolds, which are modeled on certain types of representations, from their linear models was the topic of my Ph.D. thesis (cf. [7] and [8]). (In these papers I was mainly concerned with cases such as $G = O(n)$, $U(n)$ or $Sp(n)$ where the linear model was a multiple of the standard representation.) The case of the standard T^m -action on \mathbb{C}^m is probably the simplest special case of the general theory developed in [8]. Here are a few remarks concerning the general theory.

a) Suppose a smooth G -manifold M is modeled on a linear representation of G on a vector space V . An equivariant map between G -manifolds is *isovariant* if it preserves isotropy subgroups and *transverse isovariant* if its differential induces isomorphisms between normal representations to the strata. A transverse isovariant map $F : M \rightarrow V$ induces a map $f : M/G \rightarrow V/G$ of orbit spaces and an equivariant diffeomorphism, $M \cong f^*(V)$, where $f^*(V)$ is defined as in (6.1). Thus, M is a pullback of its linear model if and only if it admits a transverse isovariant map to its linear model.

b) For $B = M/G$, consider maps $B \rightarrow V/G$ which are strata-preserving and “transverse” to the strata in some obvious sense (for short, *stratified maps*). Let f_0 and f_1 be two maps from B to V/G which are homotopic through stratified maps. It is a consequence of G. Schwarz’s Covering Homotopy Theorem that $f_0^*(V) \cong f_1^*(V)$ (cf. [25]).

c) Suppose α is the “normal orbit type” corresponding to (H, E) where H is an isotropy subgroup and E is the normal representation to the stratum of G/H orbits. The normal bundle to this stratum is a bundle over the corresponding stratum B_α in the orbit space. The fiber of this bundle over B_α is $G \times_H E$. After choosing a metric on the normal bundle, the structure group reduces to the group $S_\alpha := N_H(G \times O(E))/H$ (where $O(E)$ is the orthogonal group of E and $N_H(G \times O(E))$ is the normalizer of H in $G \times O(E)$). Let S_{prin} be the structure group for the principal orbit bundle. For some linear models, S_α is a subgroup of S_{prin} . If the principal orbit bundle of M is trivial, we get a map $B_\alpha \rightarrow S_{\text{prin}}/S_\alpha$ called the α -twist invariant of M . If, as in the situations of interest in [8], the α -stratum of M/G is homotopy equivalent to S_{prin}/S_α , the twist invariants can be used to provide a transverse isovariant map $M \rightarrow V$.

Proposition 6.2. *Suppose Y^{m+n} is a smooth T^m -manifold, modeled on the standard representation and that the bundle of principal orbits is trivial. Let $P = Y/T^m$ and let F_i be the codimension one face of P whose isotropy subgroup is the coordinate circle T_i . Then Y is the pullback of the linear model via a stratified map $f : P \rightarrow \mathbb{R}_+^m$ which takes F_i into the hyperplane $x_i = 0$. Hence, Y is T^m -equivariantly diffeomorphic to the moment angle manifold Z_P defined in (6.1).*

Sketch of Proof. In the case at hand, each $S_\alpha = T^m = S_{\text{prin}}$, so the range of each twist invariant is a point. Since each stratum of \mathbb{R}_+^m is contractible, the twist invariants can be used to construct a map $Y \rightarrow \mathbb{C}^m$ inducing $P \rightarrow \mathbb{R}_+^m$ (cf. Remark 6.1 c)). For the same reason, any two maps to \mathbb{R}_+^m are homotopic through stratified maps; so, by Remark 6.1 b), any two pullbacks of \mathbb{C}^m are equivariantly diffeomorphic. Hence, $Y \cong Z_P$. \square

Corollary 6.3. *Suppose Y and Y' are smooth T^m -manifolds which are modeled on the standard representation and which have trivial bundles of principal orbits. Then Y is equivariantly diffeomorphic to Y' if and only if Y/T^m and Y'/T^m are diffeomorphic as manifolds with corners (via a strata-preserving diffeomorphism).*

Quasitoric manifolds. A T^n -action on a manifold M^{2n} is *locally standard* if it is locally modeled on the standard representation (\mathbb{C}^n, T^n) , up to automorphisms of T^n . (N.B. Because we are allowing ourselves to vary the local actions by automorphisms of T^n , a locally standard action is usually *not* modeled on the standard representation. For example, in the standard

representation only the coordinate circles T_1, \dots, T_n occur as isotropy subgroups, while in a locally standard action it is possible to have more isotropy subgroups isomorphic to S^1 .) The orbit space P of a locally standard action on M^{2n} is a smooth n -manifold with corners. In [21] Masuda and Panov impose the following additional requirement:

(*) P is an acyclic manifold with acyclic faces.

(Such a P is called a “homology polytope” in [21].) From now on we suppose that P satisfies (*). M^{2n} is a *quasitoric manifold* if P is a simple convex polytope (cf. [12]). Let F_1, \dots, F_m be the codimension one faces of P . The isotropy subgroup at an interior point of F_i is a subgroup Λ_i isomorphic to S^1 . If $F_{i_1} \cap \dots \cap F_{i_k} \neq \emptyset$, then $\Lambda_{i_1}, \dots, \Lambda_{i_k}$ span a k -dimensional subtorus of T^n . The subgroup Λ_i is determined by a vector $\lambda_i \in \text{Hom}(S^1, T^m) \cong \mathbb{Z}^m$, well defined up to sign. This defines an epimorphism $\lambda = (\lambda_1, \dots, \lambda_m) : T^m \rightarrow T^n$ called the *characteristic function*. We will sometimes also view λ as a homomorphism $\mathbb{Z}^m \rightarrow \mathbb{Z}^n$.) It is observed in [21] that, as in [12], M^{2n} is determined up to equivariant homeomorphism by P and the characteristic function λ . In fact, as we will see in Proposition 6.4 below, we can replace “equivariant homeomorphism” by “equivariant diffeomorphism.”

Put $T = T^n$ and let $M_T := ET \times_T M$ be the Borel construction. So, M_T is a bundle over BT with fiber M . Masuda and Panov prove in [21, Theorem 1] that when (*) holds, the cohomology of M vanishes in odd degrees and that its cohomology is generated by degree two classes. It follows that the Serre spectral sequence for $M_T \rightarrow BT$ degenerates at E_2 and that we have an isomorphism of $H^*(BT)$ -modules:

$$H^*(M_T) \cong H^*(BT) \otimes H^*(M).$$

In particular, there is a short exact sequence,

$$0 \rightarrow H^2(BT) \rightarrow H^2(M_T) \rightarrow H^2(M) \rightarrow 0. \quad (6.2)$$

Given P , an acyclic manifold with acyclic faces, and a characteristic function $\lambda : T^m \rightarrow T$, we construct a smooth T -manifold $M_{(P,\lambda)}$ as follows. Put $H = \text{Ker } \lambda$. Then $H \cong T^{m-n}$ and λ induces an isomorphism $\bar{\lambda} : T^m/H \rightarrow T$. The subgroup H acts freely on the moment angle manifold Z_P . So, the quotient

$$M_{(P,\lambda)} = Z_P/H \quad (6.3)$$

is a smooth manifold with a smooth action of T^m/H . After using $\bar{\lambda}^{-1}$ to identify T with T^m/H , we get a locally standard T -action on $M_{(P,\lambda)}$

with orbit space P . Moreover, the isotropy subgroup corresponding to the codimension one face F_i is the circle Λ_i .

I am grateful to Mikiya Masuda for suggesting the proof of the following proposition in some e-mail correspondence.

Proposition 6.4. *For $T = T^n$, suppose M^{2n} has a locally standard, smooth T -action with orbit space P , where P is an acyclic manifold with acyclic faces, with codimension one faces F_1, \dots, F_m . Let $\lambda : T^m \rightarrow T$ be its characteristic function and put $H = \text{Ker } \lambda$. There is a smooth principal H -bundle $\pi : Y^{n+m} \rightarrow M^{2n}$ with $H_1(Y) = 0$. Moreover, there is a lift of the T -action to Y giving a smooth $(T \times H)$ -action on Y and an isomorphism $L : T^m \rightarrow T \times H$ such that*

- (i) *If $p_1 : T \times H \rightarrow T$ denotes projection onto the first factor, then $p_1 \circ L = \lambda : T^m \rightarrow T$ and $L(H) = H$.*
- (ii) *After using L^{-1} to get a T^m -action on Y , the T^m -action is modeled on the standard representation (\mathbb{C}^m, T^m) .*
- (iii) *Y is T^m -equivariantly diffeomorphic to Z_P and M^{2n} is T -equivariantly diffeomorphic to $M_{(P, \lambda)}$ defined by (6.3).*

We begin with some notation that will be used in the proof. Principal H -bundles over M are classified by homotopy classes of maps from M to BH , i.e., by $[M, BH]$. According to a theorem of Hattori-Yoshida [15] (also cf. [26]), given a principal H -bundle $Y \rightarrow M$, the T -action on M lifts to a $(T \times H)$ -action on Y if and only if Y is the pullback of a bundle over M_T . Since $H \cong T^{m-n}$, $[M, BH]$ is the product of $(m - n)$ copies of $[M, BS^1]$, i.e., $[M, BH] \cong \bigoplus_{i=1}^{m-n} H^2(M)$. By (6.2), $H^2(M_T) \rightarrow H^2(M)$ is onto and hence, so is $[M_T, BH] \rightarrow [M, BH]$. Therefore, any principal H -bundle over M is the pullback of a principal H -bundle over M_T . So, the Hattori-Yoshida Theorem applies. Let $L_i < (T \times H)$ be the isotropy subgroup at a point in the relative interior of F_i . The group $T \times H$ is an m -dimensional torus. To get a T^m -action on Y which is modeled on the standard representation we need the homomorphism $T^m \rightarrow T \times H$ which takes the coordinate circle T_i to L_i to be an isomorphism. (Given a collection of L_1, \dots, L_m of circle subgroups in $T \times H$, we say that the L_i span $T \times H$ if the homomorphism $T^m \rightarrow T \times H$ which takes T_i to L_i is an isomorphism.)

Lemma 6.5. *Suppose M is as above and that Y is a principal H -bundle over M . Then the L_i span $T \times H$ if and only if $H_1(Y) = 0$.*

Proof. Let $q : M \rightarrow P$ be the orbit map and $\pi : Y \rightarrow M$ the bundle projection. Choose a point p in the interior of P . Let I_i be a line segment joining p to a point in the relative interior of F_i . Put $A = \bigcup I_i$, $U = q^{-1}(A)$ and $V = \pi^{-1}(U)$. Then $q^{-1}(p) = T$ and $q^{-1}(I_i)$ is a D^2 -bundle over T/Λ_i . We have $\pi_1(T) \cong \mathbb{Z}^n$ and $\pi_1(H) \cong \mathbb{Z}^{m-n}$. Hence, $\pi_1(q^{-1}(I_i)) = \mathbb{Z}^n / \langle \lambda_i \rangle$, where $\langle \lambda_i \rangle$ is the \mathbb{Z} -submodule determined by Λ_i , and $\pi_1(U) = \mathbb{Z}^n / \sum \langle \lambda_i \rangle$. Since the $\langle \lambda_i \rangle$ span \mathbb{Z}^n , $\pi_1(U)$ is trivial. Similarly, $\pi^{-1}(q^{-1}(p)) = T \times H$ and $\pi^{-1}(q^{-1}(I_i))$ is a D^2 -bundle over $(T \times H)/L_i$. Hence, $\pi_1(\pi^{-1}(q^{-1}(I_i))) = (\mathbb{Z}^n \oplus \mathbb{Z}^{m-n}) / \langle l_i \rangle$, where $\langle l_i \rangle$ is the \mathbb{Z} -submodule determined by L_i . So, $\pi_1(V) \cong (\mathbb{Z}^n \oplus \mathbb{Z}^{m-n}) / \sum \langle l_i \rangle$.

A spectral sequence argument can be used to prove the following.

Claim. The inclusion $V \hookrightarrow Y$ induces an isomorphism $H_1(V) \cong H_1(Y)$.

Assuming this claim, we have $H_1(Y) \cong H_1(V) \cong (\mathbb{Z}^n \oplus \mathbb{Z}^{m-n}) / \sum \langle l_i \rangle$ and since the quotient is trivial if and only if the L_i span $T \times H$, we get the lemma.

To prove the claim, note that M_T and U_T are the Davis-Januszkiewicz spaces for the simple polyhedral complexes P and A , respectively (cf. [12, Section 4]). By [12, Theorem 4.8], $H^*(M_T)$ and $H^*(U_T)$ are the face rings of the simplicial complexes $N(P)$ and $N(A)$ which are dual to P and A . Such face rings are generated by H^2 , and H^2 is free abelian on the vertex set of the simplicial complex. Since $N(P)$ and $N(A)$ have the same vertex set (namely, m points), the inclusion induces an isomorphism $H^2(M_T) \xrightarrow{\cong} H^2(U_T) \cong \mathbb{Z}^m$ and hence, also an isomorphism on homology, $H_2(U_T) \xrightarrow{\cong} H_2(M_T)$. Since P is acyclic, it follows from [21] that $H_1(M) = 0 = H_1(U)$. Comparing the Serre spectral sequences in homology for $U_T \rightarrow BT$ and $M_T \rightarrow BT$, we see that $H_2(U) \xrightarrow{\cong} H_2(M)$. Then comparing the spectral sequences for the principal H -bundles, $V \rightarrow U$ and $Y \rightarrow M$, we see that $H_2(V) \xrightarrow{\cong} H_2(Y)$. Finally, if $d_2 : E_{2,0}^2 \rightarrow E_{0,1}^2$ denotes the E^2 -differential,

$$\begin{aligned} H_1(V) &= \text{Coker}(d_2 : H_2(U) \rightarrow H_1(H) \cong \mathbb{Z}^{m-n}) \\ H_1(Y) &= \text{Coker}(d_2 : H_2(M) \rightarrow H_1(H) \cong \mathbb{Z}^{m-n}). \end{aligned} \quad (6.4)$$

So, $H_1(V) \cong H_1(Y)$ establishing the claim and consequently, the lemma. \square

Proof of Proposition 6.4. The characteristic class $c(Y) = (c_1, \dots, c_{m-n})$ of a principal H -bundle $Y \rightarrow M$ lies in $\bigoplus_{i=1}^{m-n} H^2(M)$, where $H^2(M) \cong \mathbb{Z}^{m-n}$. According to Lemma 6.5, we want $H_1(Y) = 0$. To achieve this, choose c_1, \dots, c_{m-n} to be a basis for $H^2(M)$. We will then have that $d_2 : H_2(M) \rightarrow H_1(H)$ is onto and by (6.4), that $H_1(Y) = 0$. By Lemma 6.5, the L_i span $T \times H$. Let $L : T^m \rightarrow T \times H$ be an isomorphism which sends T_i to L_i .

From the definitions of L_i and Λ_i , we see that $p_1 : T \times H \rightarrow T$ takes L_i to Λ_i ; so, $p_1 \circ L = \lambda : T_m \rightarrow T$. Hence, $L(H) = L(\text{Ker } \lambda) = \text{Ker } p_1 = H$. So, Properties (i) and (ii) hold. By Proposition 6.2, Y is T^m -equivariantly diffeomorphic to Z_P . Hence, $M = Y/H$ is T -equivariantly diffeomorphic to $M_{(P,\lambda)}$, establishing Property (iii). \square

By definition, the *equivariant cohomology* of M is the cohomology of M_T . It is an algebra over $H^*(BT)$. Masuda [19] shows that if quasitoric manifolds have isomorphic equivariant cohomology as algebras over $H^*(BT)$, then they are equivariantly homeomorphic ([19, Theorem 4]). (Also, compare Metaftsis-Prassidis [21].) Using Proposition 6.4 and Corollary 1.3 we can upgrade Masuda's rigidity result from equivariant homeomorphism to equivariant diffeomorphism, as in Theorem 6.6 below. (A different proof of this, using the theory of "normal systems," is given by Wiemeler [28].)

Theorem 6.6. (cf. Masuda [19], Wiemeler [28, Corollary 5.7]). *Quasitoric manifolds have isomorphic equivariant cohomology as algebras over $H^*(BT)$ if and only if they are equivariantly diffeomorphic.*

On the other hand, from Theorem 1.4 we get the following.

Theorem 6.7. *For each $n \geq 4$, there are locally standard T^n -manifolds over contractible manifolds with contractible faces (i.e., Coxeter orbifolds of type (I)) which are equivariantly homeomorphic but not equivariantly diffeomorphic.*

Toric varieties. A nonsingular, compact toric variety is determined by a nonsingular, complete simplicial fan in \mathbb{R}^n . The intersection of such a fan with the unit sphere S^{n-1} is a totally geodesic triangulation N of S^{n-1} . Such a geodesic triangulation of S^{n-1} need not be simplicially isomorphic to the boundary complex of a simplicial polytope, e.g., see [29, p. 194]. (We were not cognizant of this fact when we wrote [12].) The rays of the fan are rational and determine a characteristic function λ . So, the fan determines a locally standard T -action on M^{2n} . The orbit space $P^n = M^{2n}/T$ has the structure of a smooth manifold with faces. Since N is a PL triangulation of S^{n-1} , P is a Coxeter orbifold of type (II). When the toric variety is projective, one can use the moment map to identify P^n with a simple polytope. In particular, the faces of P are diffeomorphic to disks. As we will see below, this holds in general (it is not automatic since the smooth 4-dimensional Poincaré Conjecture is not known.)

Proposition 6.8. *Suppose M^{2n} is nonsingular, compact toric variety and $P = M^{2n}/T$. Then each face of P is diffeomorphic to a disk.*

Proof. Each top-dimensional simplex of N corresponds to a T -action on $(D^2)^n$ with orbit space the n -cube, $[0, 1]^n$. The manifold with faces P is obtained by gluing together these n -cubes via linear isomorphisms of faces. This defines a PL structure on P so that ∂P is N with its dual PL cell structure. Therefore, each face of P (and in particular, each 4-dimensional face) is PL homeomorphic to a disk. Theorem 1.2 completes the proof. \square

References

- [1] S. Akbulut, *A fake compact contractible 4-manifold*, J. Differential Geom. **33** (1991), 335–356.
- [2] J. Cerf, *Sur les difféomorphismes de la sphère de dimension trois ($\Gamma_4 = 0$)*, (French) Lecture Notes in Math. **53** Springer-Verlag, Berlin-New York, 1968.
- [3] J. Cerf, *La stratification naturelle des espaces de fonctions différentiables réelles et le théorème de la pseudo-isotopie*, (French) Inst. Hautes Études Sci. Publ. Math. No. **39** (1970), 5–173.
- [4] R. Charney and M.W. Davis, *When is a Coxeter system determined by its Coxeter group?*, J. London Math. Soc. **61** (2000), 441–461.
- [5] S. Choi and G-S. Lee, *Projective deformations of weakly orderable Coxeter orbifolds*, arXiv:1207.3527, preprint.
- [6] M. Cohen, *Homeomorphisms between homotopy manifolds and their resolutions*, Inv. Math. **10** (1970), 239–250.
- [7] M.W. Davis, *Multiaxial Actions on Manifolds*, Springer Lecture Notes in Math. **643**, Springer-Verlag, Berlin-New York, 1978.
- [8] M.W. Davis, *Universal G -manifolds*, Amer. J. Math. **103** (1981), 103–141.
- [9] M.W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. **117** (1983), 293–325.
- [10] M.W. Davis, *The Geometry and Topology of Coxeter Groups*, London Math. Soc. Monograph Series, vol. **32**, Princeton Univ. Press, 2008.

- [11] M.W. Davis, *Lectures on orbifolds and reflection groups*, in *Transformation Groups and Moduli Spaces of Curves* (eds. L. Ji and S-T. Yau) ALM **16**, Higher Education Press, Beijing 2010, pp. 63–93.
- [12] M.W. Davis and T. Januszkiewicz, *Convex polytopes, Coxeter orbifolds and torus actions*, Duke Math. J. **62** (1991), 417–451.
- [13] M. Freedman, *The topology of four-dimensional manifolds*, J. Diff. Geom. **17** (1982), 357–453.
- [14] A. Hatcher, *A proof of the Smale conjecture*, $\text{Diff}(S^3) \simeq O(4)$, Ann. of Math. (2) **117** (1983), 553–607.
- [15] A. Hattori and T. Yoshida, *Lifting compact group actions in fiber bundles*, Journal J. Math. **2**, No. 1 (1976), 13–25.
- [16] W.C. Hsiang and W.Y. Hsiang, *Differentiable actions of compact, connected classical groups: I*. Amer. J. Math. **89** (1967), 705–786.
- [17] M. Kervaire and J. Milnor, *Groups of homotopy spheres, I*, Ann. of Math. (2) **77** (1963), 504–537.
- [18] J. Levine, *Semi-free circle actions on spheres*, Inventiones math. **22** (1973), 161–186.
- [19] M. Masuda, *Equivariant cohomology distinguishes toric manifolds*, Adv. Math. **218** (2008), no. 6, 2005–2012.
- [20] M. Masuda and T. Panov, *On the cohomology of torus manifolds*, Osaka J. Math. **43** (2006), no. 3, 711–746.
- [21] V. Metaftsis and S. Prassidis, *Topological rigidity of quasitoric actions*, arXiv:1301.0469v1.
- [22] J. Morgan and G. Tian, *Ricci Flow and the Poincaré Conjecture* Clay Mathematics Monographs, **3**. American Mathematical Society, Providence, RI; Clay Mathematics Institute, Cambridge, MA, 2007.
- [23] G. Perelman, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*. math.DG/0307245, 2003.
- [24] S. Prassidis and B. Spieler, *Rigidity of Coxeter groups*, Trans. Amer. Math. Soc. **352** (2000), 2619–2642.

- [25] G.W. Schwarz, *Lifting smooth homotopies of orbit spaces*, Publ. Math. Inst. Hautes Étud. Sci. **51** (1980), 37–135.
- [26] T.E. Stewart, *Lifting group actions in fiber bundles*, Ann. of Math. **74** (1961), 192–198.
- [27] D. Sullivan, *Singularities in spaces*, in *Liverpool Symposium*, Springer Lecture Notes **209** (1971), 196-206.
- [28] M. Wiemeler, *Exotic torus manifolds and equivariant smooth structures on quasitoric manifolds*, Math. Z. **273** (2013), 1063–1084.
- [29] G.M. Ziegler, *Lectures on polytopes*, Graduate Texts in Mathematics, **152**, Springer-Verlag, New York, 1995.

Department of Mathematics
The Ohio State University
231 W. 18th Ave.
Columbus, Ohio 43210
davis.12@math.osu.edu