

SPECIAL RICCI-HESSIAN EQUATIONS ON KÄHLER MANIFOLDS

ANDRZEJ DERDZINSKI  and PAOLO PICCIONE 

(Received 3 August 2024; accepted 9 April 2025; first published online 7 July 2025)

Communicated by Graeme Wilkin

Abstract

Special Ricci–Hessian equations on Kähler manifolds (M, g) , as defined by Maschler [‘Special Kähler–Ricci potentials and Ricci solitons’, *Ann. Global Anal. Geom.* **34** (2008), 367–380], involve functions τ on M and state that, for some function α of the real variable τ , the sum of $\alpha \nabla d\tau$ and the Ricci tensor equals a functional multiple of the metric g , while $\alpha \nabla d\tau$ itself is assumed to be nonzero almost everywhere. Three well-known obvious ‘standard’ cases are provided by (non-Einstein) gradient Kähler–Ricci solitons, conformally-Einstein Kähler metrics, and special Kähler–Ricci potentials. We show that, outside of these three cases, such an equation can only occur in complex dimension two and, at generic points, it must then represent one of three types, for which, up to normalizations, $\alpha = 2 \cot \tau$, $\alpha = 2 \coth \tau$, or $\alpha = 2 \tanh \tau$. We also use the Cartan–Kähler theorem to prove that these three types are actually realized in a ‘nonstandard’ way.

2020 *Mathematics subject classification*: primary 53C55; secondary 53C25.

Keywords and phrases: Ricci–Hessian equation, Kähler manifold.

1. Introduction

Following Maschler [17, page 367], one says that functions τ, α, σ on a Riemannian manifold (M, g) with the Ricci tensor r satisfy a *Ricci–Hessian equation* if

$$\alpha \nabla d\tau + r = \sigma g \quad \text{for some function } \sigma : M \rightarrow \mathbb{R}, \quad (1-1)$$

∇ being the Levi–Civita connection of g . We call (1-1) *special* when

$$\alpha \nabla d\tau \neq 0 \text{ on a dense set, } \dim M = n > 2 \quad \text{and} \quad \alpha \text{ is a } C^\infty \text{ function of } \tau. \quad (1-2)$$

Conditions (1-1)–(1-2) are satisfied in several situations that have been studied (see below), raising a natural question: *Which functions $\tau \mapsto \alpha$ can be realized in this way?*

Both authors’ research was supported in part by a FAPESP OSU 2015 Regular Research Award (FAPESP grant: 2015/50265-6).

© The Author(s), 2025. Published by Cambridge University Press on behalf of Australian Mathematical Publishing Association Inc. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike licence (<https://creativecommons.org/licenses/by-nc-sa/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the same Creative Commons licence is included and the original work is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use.



The present paper provides an answer in the Kähler case, outside of the classes that are already well understood; see Theorems D and E.

There are three well-known classes of examples leading to (1-1)–(1-2).

- (I) Non-Einstein gradient Ricci almost-solitons [1, 20], including (non-Einstein) gradient Ricci solitons [15]. Here, α is a nonzero constant.
- (II) Conformally Einstein metrics g , with $\tau > 0$ and $\alpha = (n - 2)/\tau$, the Einstein metric being $\hat{g} = g/\tau^2$. Compare with [11, (6.2)].
- (III) Special Kähler–Ricci potentials τ on Kähler manifolds, at points where r is not a multiple of g . See [11, Remark 7.4].

A *special Kähler–Ricci potential* [11, Section 7] on a Kähler manifold (M, g) with the complex-structure tensor J is any nonconstant function τ on M having a real-holomorphic gradient $v = \nabla\tau$ for which, at points where $v \neq 0$, all nonzero vectors orthogonal to v and Jv are eigenvectors of both $\nabla d\tau$ and r . Such triples (M, g, τ) are completely understood, both locally [11] and in the compact case [12].

The classes (I)–(III) are far from disjoint: for instance [11, Corollary 9.3], in the Kähler category, if $n > 4$, class (II) is a special case of class (III).

We are interested in $M, g, \tau, \alpha, \sigma$ satisfying (1-1)–(1-2) and such that

$$2r(v, \cdot) = -dY \quad \text{for } v = \nabla\tau \quad \text{and} \quad Y = \Delta\tau. \quad (1-3)$$

Here, and throughout the paper, we use the notational conventions

$$v = \nabla\tau, \quad Q = g(v, v), \quad Y = \Delta\tau, \quad n = \dim M \quad (1-4)$$

whenever (M, g) is a Riemannian manifold and $\tau : M \rightarrow \mathbb{R}$. As we point out near the end of Section 2, with J denoting the complex-structure tensor,

for Kähler metrics g , conditions (1-1)–(1-2) imply (1-3),
and the gradient $v = \nabla\tau$ is a real-holomorphic vector field
or, equivalently, Jv is a real-holomorphic g -Killing field. (1-5)

Assuming (1-1)–(1-2), we may treat the derivatives $\alpha' = d\alpha/d\tau$ and α'' both as functions of the real variable τ and as functions $M \rightarrow \mathbb{R}$. In Sections 3 and 4, we prove the following two results, as well as Theorem D, stated below.

THEOREM A. *Under the hypotheses (1-1)–(1-3), at points where $\alpha'' + \alpha\alpha' \neq 0$ and $d\tau \neq 0$, both $Q = g(\nabla\tau, \nabla\tau)$ and $Y = \Delta\tau$ are, locally, functions of τ .*

THEOREM B. *Let functions τ, α, σ satisfy a special Ricci–Hessian equation (1-1), with (1-2), on a Kähler manifold (M, g) of real dimension $n \geq 4$. If $\alpha d\alpha$ and $d\tau$ are nonzero at all points of an open submanifold U of M , and:*

- (i) $n > 4$; or
- (ii) $n = 4$ and $d\sigma \wedge d\tau = 0$ identically in U ; or finally,
- (iii) $dQ \wedge d\tau = 0$ everywhere in U , where $Q = g(\nabla\tau, \nabla\tau)$,

then $\tau : U \rightarrow \mathbb{R}$ is a special Kähler–Ricci potential on the Kähler manifold (U, g) .

With v, Q, Y as in (1-4), a function τ on a Riemannian manifold (M, g) has $dQ \wedge d\tau = 0$ if and only if Q is locally, at points where $d\tau \neq 0$, a function of τ . This amounts to requiring the integral curves of v to be reparameterized geodesics (since, due to (2-2) below, the latter condition means that $\nabla_v v$ is a functional multiple of v). Such functions τ , called *transnormal*, have been studied extensively [3, 18, 21], and are referred to as *isoparametric* when, in addition, $dY \wedge d\tau = 0$.

Theorem B renders the transnormal case $dQ \wedge d\tau = 0$, as well as real dimensions $n > 4$, rather uninteresting in the context of special Ricci–Hessian equations (1-1)–(1-2) on Kähler manifolds, since at $d\alpha$ -generic points (see the end of Section 2), one then ends up with examples (I) or (III) above, compare with Remark 4.3, of which the former is the subject of a large existing literature, and the latter, as mentioned earlier, has been completely described. This is why our next two results focus exclusively on the real dimension four and functions τ with $dQ \wedge d\tau$ not identically zero.

REMARK C. Equation (1-1), with (1-2), remains satisfied after τ and the function $\tau \mapsto \alpha = \alpha(\tau)$ have been subjected to an *affine modification* in the sense of being replaced with $\hat{\tau}$ and $\hat{\tau} \mapsto \hat{\alpha}(\hat{\tau})$ given by $\hat{\tau} = p + \tau/c$ and $\hat{\alpha}(\hat{\tau}) = c\alpha(c\hat{\tau} - cp)$ for real constants $c \neq 0$ and p .

THEOREM D. *If the special Ricci–Hessian equation (1-1) and (1-2) both hold for functions τ, α, σ on a Kähler manifold (M, g) of real dimension four, while $dQ \wedge d\tau \neq 0$ everywhere in an open connected set $U \subseteq M$, then the function α of the variable τ and its derivative $\alpha' = d\alpha/d\tau$ satisfy, on U , the equation*

$$\alpha'' + \alpha\alpha' = 0, \text{ that is } 2\alpha' + \alpha^2 = 4\varepsilon \text{ with a constant } \varepsilon \in \mathbb{R}. \quad (1-6)$$

In addition, for Q and Y as in (1-4), and the scalar curvature s , the functions

$$2\theta = \alpha s + 4\varepsilon Y \quad \text{and} \quad \kappa = \theta\psi + \alpha^{-1}Y - Q \text{ are both constant,} \quad (1-7)$$

ψ being given by $4\varepsilon\psi = \tau - 2/\alpha$, if $\varepsilon \neq 0$, or $3\psi = 2/\alpha^3$, when $\varepsilon = 0$. Furthermore, σ in (1-1) and the function F of the variable τ characterized by

$$4\varepsilon F = \theta(2 - \tau\alpha) + 4\varepsilon\kappa\alpha \text{ for } \varepsilon \neq 0 \quad \text{and} \quad F = \kappa\alpha - 2\theta/(3\alpha^2) \text{ if } \varepsilon = 0, \quad (1-8)$$

and thus depending on the real constants θ, κ , satisfy the conditions

$$(a) \quad Y - Q\alpha = F, \quad (b) \quad 2\sigma = -(Q\alpha' + F'), \quad (c) \quad \Delta\alpha = Fa' = -F''. \quad (1-9)$$

Finally, up to affine modifications (see Remark C), the pair $(\alpha(\tau), \varepsilon)$ is one of the following five: $(2, 1)$, $(2/\tau, 0)$, $(2 \tanh \tau, 1)$, $(2 \coth \tau, 1)$, $(2 \cot \tau, -1)$.

THEOREM E. *Each of the five options listed in Theorem D, namely,*

$$(2, 1), \quad (2/\tau, 0), \quad (2 \tanh \tau, 1), \quad (2 \coth \tau, 1), \quad (2 \cot \tau, -1),$$

is realized by a special Ricci–Hessian equation (1-1)–(1-2) on a real-analytic Kähler manifold (M, g) of real dimension four such that, with $v = \nabla\tau$ and $Q = g(v, v)$, one has

$dQ \wedge d\tau \neq 0$ somewhere in M and Jv lies in a two-dimensional Abelian Lie algebra of Killing fields.

For $(2, 1)$ and $(2/\tau, 0)$, one can choose (M, g) to be compact and biholomorphic to the two-point blow-up of \mathbb{CP}^2 , where g is the Wang–Zhu toric Kähler–Ricci soliton [22, Theorem 1.1] or respectively the Chen–LeBrun–Weber conformally-Einstein Kähler metric [6, Theorem A].

In contrast with the final clause of Theorem E, we do not know whether the remaining three options, $(2 \tanh \tau, 1)$, $(2 \coth \tau, 1)$, and $(2 \cot \tau, -1)$, may be realized on a *compact* Kähler surface. An analytic-continuation phenomenon described below (Section 13) may hint at the plausibility of trying to obtain such compact examples via small deformations of the Wang–Zhu or Chen–LeBrun–Weber metric, combined with suitable affine modifications.

For the pairs $(2, 1)$ and $(2/\tau, 0)$ in Theorem D, the constancy conclusions of (1-7) are well known: [7, page 201], [9, page 417, Proposition 3(i) and page 419, (40)].

The paper is organized as follows. Section 2 contains the preliminaries. Consequences of special Ricci–Hessian equations, leading to proofs of Theorems A, B and D, are presented in the next two sections. Sections 5–12 are devoted to proving Theorem E: we rephrase it as solvability of the system (6-1) of quasi-linear first-order partial differential equations, subject to the additional conditions (6-2), which allows us to derive our claim from the Cartan–Kähler theorem for exterior differential systems.

2. Preliminaries

All manifolds and Riemannian metrics are assumed to be of class C^∞ . By definition, a manifold is connected. We use the symbol δ for divergence.

On a manifold with a torsion-free connection ∇ , the Ricci tensor r satisfies the Bochner identity $r(\cdot, v) = \delta \nabla v - d[\delta v]$, where v is any vector field. Its coordinate form $R_{jk}v^k = v^k_{,jk} - v^k_{,kj}$ arises via contraction from the Ricci identity $v^l_{,jk} - v^l_{,kj} = R_{jkq}^l v^q$. (We use the sign convention for R such that $R_{jk} = R_{jqq}^q$.) Applied to the gradient v of a function τ on a Riemannian manifold, this yields

$$\delta[\nabla d\tau] = r(v, \cdot) + dY \quad \text{with } v = \nabla\tau \quad \text{and} \quad Y = \Delta\tau. \quad (2-1)$$

Also, given a function τ on a Riemannian manifold,

$$2[\nabla d\tau](v, \cdot) = dQ, \quad \text{where } v = \nabla\tau \quad \text{and} \quad Q = g(v, v), \quad (2-2)$$

as one sees noting that, in local coordinates, $(\tau_{,k}\tau^k)_{,j} = 2\tau_{,kj}\tau^k$. We can rewrite the relations (2-1)–(2-2) using the interior product ι_v , and then they read

$$(a) \quad \delta[\nabla d\tau] = \iota_v r + dY, \quad (b) \quad 2\iota_v[\nabla d\tau] = dQ \quad \text{with (1-4).} \quad (2-3)$$

Finally, for the Ricci tensor r and scalar curvature s of any Riemannian metric,

$$2\delta r = ds, \quad (2-4)$$

which is known as the Bianchi identity for the Ricci tensor. Its coordinate form $2g^{kl}R_{jk,l} = s_j$ is immediate if one transvects with ('multiplies' by) g^{kl} the equality $R_{jkl}{}^q,_q = R_{jlk} - R_{klj}$ obtained by contracting the second Bianchi identity.

The *harmonic-flow condition* for a vector field v on a Riemannian manifold (M, g) , meaning that the flow of v consists of (local) harmonic diffeomorphisms, is known [19] to be equivalent to the equation

$$g(\Delta v, \cdot) = -r(v, \cdot), \quad (2-5)$$

the vector field Δv having the local components $[\Delta v]^j = v^{j,k}_k$. See also [14, Theorem 3.1]. When $v = \nabla\tau$ is the gradient of a function $\tau : M \rightarrow \mathbb{R}$,

$$\text{the harmonic-flow condition (2-5) amounts to (1-3).} \quad (2-6)$$

In fact, by (2-1), $2r(v, \cdot) + dY = \delta[\nabla d\tau] + r(v, \cdot) = g(\Delta v, \cdot) + r(v, \cdot)$, as $[\Delta v]_j = v_{j,k}{}^k = \tau_{jk}{}^k = \tau_{kj}{}^k = \tau^{kj}{}_k = (\delta[\nabla d\tau])_j$.

Furthermore (see, for example, [11, Lemma 5.2]), on a Kähler manifold (M, g) ,

conditions (1-1)–(1-2) imply real-holomorphicity of the gradient $v = \nabla\tau$, while Jv is then a holomorphic Killing field, due to the resulting Hermitian symmetry of $\nabla d\tau$. (2-7)

Since holomorphic mappings between Kähler manifolds are harmonic, every holomorphic vector field on a Kähler manifold satisfies (2-5); see also [14, Remark 3.2]. Now, (1-5) follows from (2-6). In other words, as observed by Calabi [5], on Kähler manifolds, one has

$$(1-3), \text{ with (1-4) for all real-holomorphic gradients } v = \nabla\tau. \quad (2-8)$$

Given a tensor field θ on a manifold M , we say that a point $x \in M$ is θ -generic if x has a neighborhood on which either $\theta = 0$ identically, or $\theta \neq 0$ everywhere. Such points clearly form a dense open subset of M .

3. Ricci–Hessian equations

As a consequence of (1-1), for the scalar curvature s , with (1-4),

$$n\sigma = Y\alpha + s, \quad \text{where } n = \dim M. \quad (3-1)$$

Applying $2\iota_v$ or 2δ to (1-1), we obtain, from (2-3)–(2-4) and (1-4),

$$\begin{aligned} \text{(i)} \quad & \alpha dQ + 2r(v, \cdot) = 2\sigma d\tau, \\ \text{(ii)} \quad & 2[\nabla d\tau](\nabla\alpha, \cdot) + 2\alpha[r(v, \cdot) + dY] + ds = 2d\sigma. \end{aligned} \quad (3-2)$$

In the case where (1-1)–(1-2) hold along with (1-3), one may rewrite (3-2) as

$$\begin{aligned} \text{(i)} \quad & \alpha dQ - dY = 2\sigma d\tau, \\ \text{(ii)} \quad & 2[\nabla d\tau](\nabla\alpha, \cdot) + \alpha dY + ds = 2d\sigma, \end{aligned} \quad (3-3)$$

which, in view of (2-2) and (3-1), amounts to nothing other than

$$\begin{aligned} \text{(i)} \quad d(Q\alpha - Y) &= (Q\alpha' + 2\sigma)d\tau, \\ \text{(ii)} \quad d[Q\alpha' + (n-2)\sigma] &= (Q\alpha'' + Y\alpha')d\tau, \end{aligned} \quad (3-4)$$

as the assumption, in (1-2), that α is a C^∞ function of τ allows us to write

$$d\alpha = \alpha'd\tau, \quad \nabla\alpha = \alpha'v, \quad 2[\nabla d\tau](\nabla\alpha, \cdot) = \alpha'dQ, \quad \text{where } \alpha' = d\alpha/d\tau, \quad (3-5)$$

since (2-2) gives $2[\nabla d\tau](\nabla\alpha, \cdot) = 2\alpha'[\nabla d\tau](v, \cdot) = \alpha'dQ$. Due to (3-4), conditions (1-1)–(1-3) imply that, locally, at points at which $d\tau \neq 0$,

$Q\alpha - Y$ and $Q\alpha' + (n-2)\sigma$ are functions of τ , with
the respective τ -derivatives $Q\alpha' + 2\sigma$ and $Q\alpha'' + Y\alpha'$,
which, consequently, must themselves be functions of τ .

PROOF OF THEOREM A. At the points in question, using (3-6) to equate both $Q\alpha - Y$ and $Q\alpha'' + Y\alpha'$ to some specific functions of τ , we obtain a system of two linear equations with the nonzero determinant $\alpha'' + \alpha\alpha'$, imposed on the unknowns Q, Y , and our assertion follows since $\alpha'' + \alpha\alpha'$ is also a function of τ . \square

Assuming only (1-1), for $n = \dim M$, with the aid of (3-1), we rewrite (3-2) as

$$\begin{aligned} n[\alpha dQ + 2r(v, \cdot)] - 2(Y\alpha + s)d\tau &= 0, \\ 2n\{[\nabla d\tau](\nabla\alpha, \cdot) + \alpha r(v, \cdot)\} + 2[(n-1)\alpha dY - Yd\alpha] + (n-2)ds &= 0. \end{aligned}$$

If (1-3) holds as well, replacing $2r(v, \cdot)$ here with $-dY$, we obtain $n(\alpha dQ - dY) - 2(Y\alpha + s)d\tau = 0$ and $2n\{[\nabla d\tau](\nabla\alpha, \cdot) + (n-2)(\alpha dY + ds)\} - 2Yd\alpha = 0$. Thus, when (1-1)–(1-3) are all satisfied, (3-5) gives

$$\begin{aligned} \text{(a)} \quad n(\alpha dQ - dY) - 2(Y\alpha + s)d\tau &= 0, \\ \text{(b)} \quad na'dQ + (n-2)(\alpha dY + ds) - 2Y\alpha'd\tau &= 0. \end{aligned} \quad (3-7)$$

4. Ricci–Hessian equations on Kähler manifolds

The goal of this section is to prove Theorems B and D.

In any complex manifold, $d\omega = 0$ and $\omega(J\cdot, \cdot)$ is symmetric if $\omega = i\partial\bar{\partial}\tau$, that is, if $2\omega = -d[J^*d\tau]$ for a real-valued function τ , with the 1-form $J^*d\tau = (d\tau)J$, which sends any tangent vector field v to $d_v\tau$. Our exterior-derivative and exterior-product conventions, for 1-forms ξ, ξ' and vector fields u, v , are

$$\begin{aligned} [d\xi](u, v) &= d_u[\xi(v)] - d_v[\xi(u)] - \xi([u, v]), \\ [\xi \wedge \xi'](u, v) &= \xi(u)\xi'(v) - \xi(v)\xi'(u). \end{aligned} \quad (4-1)$$

For a torsion-free connection ∇ , (4-1) gives $[d\xi](u, v) = [\nabla_u\xi](v) - [\nabla_v\xi](u)$, so that, if in addition $\nabla J = 0$, on an almost-complex manifold,

$$2i\partial\bar{\partial}\tau = [\nabla d\tau](J\cdot, \cdot) - [\nabla d\tau](\cdot, J\cdot). \quad (4-2)$$

In the case of a Kähler metric g on a complex manifold M , (1-1) implies that

$$i\alpha\partial\bar{\partial}\tau + \rho = \sigma\omega, \quad (4-3)$$

ω, ρ being the Kähler and Ricci forms, with both terms on the right-hand side of (4-2) equal, as the Hermitian symmetry of $\nabla d\tau$ follows from those of ρ and ω .

REMARK 4.1. As an obvious consequence of the last line in (2-7), if g is a Kähler metric, conditions (1-1)–(1-2) are *equivalent* to (4-3) along with (1-2) and real-holomorphicity of the gradient $v = \nabla\tau$.

REMARK 4.2. For the Kähler form ω of a Kähler manifold (M, g) of real dimension $n \geq 4$, the operator $\zeta \mapsto \zeta \wedge \omega$ acting on differential q -forms is injective if $q = 2$ and $n > 4$, or $q = 1$. Namely, the contraction of $\zeta \wedge \omega$ against ω yields a nonzero constant times $(n - 4)\zeta + 2\langle\omega, \zeta\rangle\omega$ (if $q = 2$), or times $(n - 2)\zeta$ (if $q = 1$). In the former case, ζ with $\zeta \wedge \omega = 0$ is thus a multiple of ω , and hence is 0.

REMARK 4.3. Whenever (4-3) with a *constant* α holds on a Kähler manifold of real dimension $n \geq 4$, constancy of σ follows (from Remark 4.2, as $d\sigma \wedge \omega = 0$).

We have the following result, due to Maschler [17, Proposition 3.3].

LEMMA 4.4. *Condition (1-1) on a Kähler manifold (M, g) of real dimension $n > 4$ implies that $d\sigma \wedge d\alpha = 0$. Equivalently, wherever $d\alpha$ is nonzero, σ must, locally, be a function of α .*

PROOF. By (4-3), $0 = d\rho = d\sigma \wedge \omega - d\alpha \wedge i\partial\bar{\partial}\tau$, so that $d\alpha \wedge d\sigma \wedge \omega = 0$, and our assertion is immediate from Remark 4.2. \square

PROOF OF THEOREM B. In all three cases (i)–(iii), $d\sigma \wedge d\tau = 0$. For case (i), this follows from Lemma 4.4 while, when $dQ \wedge d\tau = 0$ on U , we see that, in view of the equality $\alpha'dQ + \alpha dY = d(2\sigma - s)$ arising from (3-3ii) and (3-5), Y and $2\sigma - s$ are, locally, functions of τ , and hence so is σ , as a consequence of (3-1) with $n \geq 4$. Now, [11, Corollary 9.2] yields our claim. \square

PROOF OF THEOREM D. As $dQ \wedge d\tau \neq 0$ everywhere in U , Theorem A implies (1-6) and, consequently, also the final clause about the five possible pairs.

Next, in (1-6), $4d\theta = 2d[\alpha s + (2\alpha' + \alpha^2)Y] = 2[ads + sda + (2\alpha' + \alpha^2)dY]$, which, as $n = 4$, equals, in view of (3-5),

$$\alpha[n\alpha'dQ + (n - 2)(\alpha dY + ds) - 2Y\alpha'd\tau] - \alpha'[n(\alpha dQ - dY) - 2(Y\alpha + s)d\tau],$$

and hence vanishes due to (3-7). However, the function ψ of τ defined in the theorem is an antiderivative of $1/\alpha^2$, meaning that

$$\psi' = 1/\alpha^2. \quad (4-4)$$

Namely, by (1-6), $4\epsilon\psi' = 1 + 2\alpha'/\alpha^2 = (2\alpha' + \alpha^2)/\alpha^2 = 4\epsilon/\alpha^2$ if $\epsilon \neq 0$, and $3\psi' = -6\alpha'/\alpha^4 = 3/\alpha^2$ when $\epsilon = 0$, as $2\alpha' = -\alpha^2$.

Furthermore, $d(\theta\psi + \alpha^{-1}Y - Q) = 0$. In fact, $d\alpha = \alpha'd\tau$ in (3-5), and similarly for ψ , so that, from (4-4), $d(\theta\psi) = \theta d\psi = \theta\psi'd\tau = \theta\alpha^{-2}d\tau$, and $\alpha d(\alpha^{-1}Y) = dY - \alpha^{-1}Y\alpha'd\tau$. Also, $2(\theta - Ya') = (Y\alpha + s)\alpha$ from (1-6)–(1-7). These relations yield

$$\begin{aligned} -4\alpha d(\theta\psi + \alpha^{-1}Y - Q) &= 4[(\alpha dQ - dY) - (\theta - Ya')\alpha^{-1}d\tau] \\ &= n(\alpha dQ - dY) - 2(Y\alpha + s)d\tau, \end{aligned}$$

with $n = 4$, which equals 0 by (3-7a).

Finally, (1-8) and the second relation in (1-7) easily give (1-9a). Thus, by (1-4) and (3-5), $(Q\alpha' + F')Q = (Q\alpha' + F')d_v\tau = Qd_v\alpha + d_vF$, which, due to (1-9a), equals $d_vY - \alpha d_vQ$. At the same time, $-\iota_v$ applied to (3-3i) yields $d_vY - \alpha d_vQ = -2Q\sigma$. We thus get (1-9b). To obtain (1-9c), note that, from (1-4), $\Delta\alpha = Q\alpha'' + Ya'$, which, by (1-6) and (1-9a), equals $(Y - Q\alpha)\alpha' = Fa' = -F''$, where the last equality trivially follows from (1-8). \square

Theorem D has a partial converse: if a nonconstant function τ with real-holomorphic gradient $v = \nabla\tau$ on a Kähler surface (M, g) and a function α of the real variable τ satisfy (1-6) and (1-7), then they must also satisfy the Ricci–Hessian equation (1-1) with σ given by (3-1) for $n = 4$.

In fact, $b(v, \cdot) = 0$, where b denotes the traceless Hermitian symmetric 2-tensor field $\alpha\nabla d\tau + r - \sigma g$. Namely, (1-3)–(1-5) and (2-3b) yield $4b(v, \cdot) = 2\alpha dQ - 2dY - 4\sigma d\tau$ which, due to (3-1) and (3-5), equals $2\alpha dQ - 2dY - (Ya' + s)d\tau$, and so $-4\alpha b(v, \cdot) = 2\alpha^2 d(\theta\psi + \alpha^{-1}Y - Q) + (\alpha s + 4\epsilon Y - 2\theta)d\tau$. (Note that, by (1-6) and (4-4), $4\epsilon = 2\alpha' + \alpha^2$ and $2\alpha^2 d(\theta\psi) = 2\theta d\tau$.) Thus, $b = 0$, since b corresponds, via g , to a *complex-linear* bundle morphism $TM \rightarrow TM$.

5. The local Kähler potentials

This and the following seven sections are devoted to proving Theorem E.

In an open set $M \subseteq \mathbb{R}^4$ with the Cartesian coordinates x, x', u, u' arranged into the complex coordinates $(x + ix', u + iu')$ for the complex plane $\mathbb{C}^2 = \mathbb{R}^4$ carrying the standard complex structure J , one has $J^*dx = -dx'$ and $J^*du = -du'$, so that, if a C^∞ function f on M only depends on x and u , the relation $2i\partial\bar{\partial}f = -d[J^*df]$ yields, with subscripts denoting partial differentiations,

$$2i\partial\bar{\partial}f = f_{xx}dx \wedge dx' + f_{xu}(dx \wedge du' + du \wedge dx') + f_{uu}du \wedge du',$$

since $df = f_x dx + f_u du$. Furthermore, we set

$$v = \partial_x \quad \text{and} \quad w = \partial_u \quad (\text{the real coordinate vector fields}). \quad (5-1)$$

For the Kähler metric g on M having the Kähler form $\omega = 2i\partial\bar{\partial}\phi$, where the function $\phi : M \rightarrow \mathbb{R}$ is assumed to depend on x and u only, 2ϕ is a Kähler potential [2, page 85] of g , and the above formula for $2i\partial\bar{\partial}f$, with $f = \phi$, becomes

$$\omega = \phi_{xx}dx \wedge dx' + \phi_{xu}(dx \wedge du' + du \wedge dx') + \phi_{uu}du \wedge du'. \quad (5-2)$$

Generally, for a skew-Hermitian 2-form $\zeta = Qdx \wedge dx' + S(dx \wedge du' + du \wedge dx') + Bdu \wedge du'$ and the Hermitian symmetric 2-tensor field a with $\zeta = a(J \cdot, \cdot)$, one has

$$a = Q(dx \otimes dx + dx' \otimes dx') + S(dx \otimes du + du \otimes dx + dx' \otimes du' + du' \otimes dx') + B(du \otimes du + du' \otimes du'),$$

due to (4-1), and so the components of a relative to the coordinates (x, x', u, u') form the matrix

$$\begin{bmatrix} Q & 0 & S & 0 \\ 0 & Q & 0 & S \\ S & 0 & B & 0 \\ 0 & S & 0 & B \end{bmatrix} \quad \text{with the determinant } (QB - S^2)^2. \quad (5-3)$$

When $a = g$, (5-2) with $\zeta = \omega$ gives $(Q, S, B) = (\phi_{xx}, \phi_{xu}, \phi_{uu})$. Thus,

$$\phi_{xx} > 0 \quad \text{and} \quad \Pi > 0 \quad \text{for } \Pi = \phi_{xx}\phi_{uu} - \phi_{xu}^2, \quad (5-4)$$

which amounts to Sylvester's criterion for positive definiteness of g , namely, positivity of the upper left subdeterminants of (5-3). From now on, we set

$$\begin{aligned} (\tau, \lambda, Q, S, B) &= (\phi_x, \phi_u, \phi_{xx}, \phi_{xu}, \phi_{uu}), \text{ so that} \\ Q > 0 \text{ and } \Pi &= QB - S^2 > 0 \text{ due to (5-4).} \end{aligned} \quad (5-5)$$

With div , Δ denoting the g -divergence and g -Laplacian, for τ, λ, Q in (5-5):

- (a) the functions τ, λ have the holomorphic g -gradients $v = \partial_x$ and $w = \partial_u$;
- (b) the other coordinate fields Jv and Jw are holomorphic g -Killing fields;
- (c) $Q = g(v, v)$ and $\Delta\tau = \text{div } v = [\log \Pi]_x$, while $\Delta\lambda = \text{div } w = [\log \Pi]_u$.

Namely, (5-1) and (5-3) yield item (a). Also, item (b) follows since ϕ only depends on x and u . Finally, (5-3) has the determinant Π^2 , and so $\Pi dx \wedge dx' \wedge du \wedge du'$ is the volume form of g , on which $\mathfrak{L}_v, \mathfrak{L}_w$ act (see (5-1)) via partial differentiations ∂_x, ∂_u of the Π factor. Thus, $\text{div } v = [\log \Pi]_x$ and $\text{div } w = [\log \Pi]_u$, compare with [16, page 281]. Finally, by item (a), (5-1) and (5-5), $g(v, v) = d_v\tau = \partial_x\tau = \partial_x\phi_x = \phi_{xx} = Q$.

For our $(\tau, \lambda) = (\phi_x, \phi_u)$, the mapping $(x, u) \mapsto (\tau, \lambda)$ is locally diffeomorphic due to (5-4), which makes $(Q, S, B) = (\phi_{xx}, \phi_{xu}, \phi_{uu})$, locally, a triple of real-valued functions of the new variables τ, λ . Thus, from the chain rule, in matrix form,

$$\begin{aligned} \text{(i)} \quad \begin{bmatrix} \partial_x & \partial_u \end{bmatrix} &= \begin{bmatrix} \partial_\tau & \partial_\lambda \end{bmatrix} \begin{bmatrix} Q & S \\ S & B \end{bmatrix}, \quad \text{(ii)} \quad \begin{bmatrix} d\tau \\ d\lambda \end{bmatrix} = \begin{bmatrix} Q & S \\ S & B \end{bmatrix} \begin{bmatrix} dx \\ du \end{bmatrix}, \quad \text{and so} \\ \text{(iii)} \quad (QB - S^2) \begin{bmatrix} \partial_\tau & \partial_\lambda \end{bmatrix} &= \begin{bmatrix} \partial_x & \partial_u \end{bmatrix} \begin{bmatrix} B & -S \\ -S & Q \end{bmatrix}, \\ \text{(iv)} \quad (QB - S^2) \begin{bmatrix} dx \\ du \end{bmatrix} &= \begin{bmatrix} B & -S \\ -S & Q \end{bmatrix} \begin{bmatrix} d\tau \\ d\lambda \end{bmatrix}. \end{aligned} \quad (5-6)$$

With subscripts still denoting partial differentiations, the obvious integrability conditions $Q_u - S_x = S_u - B_x = 0$ and (5-5) give, due to (5-6i),

$$SQ_\tau + BQ_\lambda = QS_\tau + SS_\lambda, \quad SS_\tau + BS_\lambda = QB_\tau + SB_\lambda, \quad Q > 0, \quad QB > S^2. \quad (5-7)$$

Conversely, if functions Q, S, B of the variables τ, λ satisfy (5-7), then, locally:

(d) $(Q, S, B) = (\phi_{xx}, \phi_{xu}, \phi_{uu})$ for a function ϕ , with (5-4), of the variables x, u related to τ, λ via $(\tau, \lambda) = (\phi_x, \phi_u)$, and Q, S, B determine each of ϕ, x, u uniquely up to additive constants.

In fact, the equalities in (5-7) state precisely that the vector fields $Q\partial_\tau + S\partial_\lambda$ and $S\partial_\tau + B\partial_\lambda$ commute or, equivalently, the 1-forms

$$(QB - S^2)^{-1}(Bd\tau - Sd\lambda) \text{ and } (QB - S^2)^{-1}(Qd\lambda - Sd\tau),$$

dual to them, are closed, and we may declare these vector fields (or 1-forms) to be ∂_x, ∂_u or respectively dx, du . Now that, locally, x, u are defined, up to additive constants, we obtain ϕ by solving the system $(\phi_x, \phi_u) = (\tau, \lambda)$, where τ, λ are treated as functions of x, u via the resulting locally diffeomorphic coordinate change $(\tau, \lambda) \mapsto (x, u)$. Closedness of $\tau dx + \lambda du$ and the equality $(Q, S, B) = (\phi_{xx}, \phi_{xu}, \phi_{uu})$ are obvious: our choice of dx and du gives (5-6iv), and hence (5-6ii), so that $d\tau \wedge dx + d\lambda \wedge du = 0$.

The g -Laplacians of τ and λ can also be expressed as:

(e) $\Delta\tau = Q_\tau + S_\lambda$ and $\Delta\lambda = S_\tau + B_\lambda$; while
(f) $\Pi_x = (Q_\tau + S_\lambda)\Pi$ and $\Pi_u = (S_\tau + B_\lambda)\Pi$ for $\Pi = QB - S^2$.

To see this, first note that, by (5-6i), $(QB - S^2)_x = Q(QB - S^2)_\tau + S(QB - S^2)_\lambda = Q(QB_\tau + SB_\lambda - SS_\tau) - S(QS_\tau + SS_\lambda - BQ_\lambda) - S^2S_\lambda + QBQ_\tau$. Using (5-7) to replace the two three-term sums in parentheses by BS_λ and SQ_τ , we thus obtain the first part of item (f). For the second one, we similarly rewrite $(QB - S^2)_u = S(QB - S^2)_\tau + B(QB - S^2)_\lambda$ as $S(QB_\tau - BS_\lambda - SS_\tau) - S^2S_\tau + B(BQ_\lambda + SQ_\tau - SS_\lambda) + QBB_\lambda$, and use analogous three-term replacements based on (5-7). Now item (e) follows from item (c), (5-5), and item (f).

THEOREM 5.1. *In \mathbb{C}^2 with the complex coordinates $(x + ix', u + iu')$, given an open subset M and a function $\phi : M \rightarrow \mathbb{R}$ of the real variables x, u , having the property (5-4), let g be the Kähler metric on M with the Kähler potential 2ϕ . The following two conditions are equivalent.*

(i) *The special Ricci–Hessian equation (1-1)–(1-2) holds on M for $\tau = \phi_x$, and $dQ \wedge d\tau \neq 0$ everywhere in M , with $Q = g(\nabla\tau, \nabla\tau)$. Thus, by Theorem D, one has (1-6) and (1-9a), where $Y = \Delta\tau$ and F is the function of τ characterized by (1-8) for the constants θ, κ in (1-7).*

(ii) *The triple $(Q, S, B) = (\phi_{xx}, \phi_{xu}, \phi_{uu})$ of functions of the new variables $(\tau, \lambda) = (\phi_x, \phi_u)$ satisfies (5-7) with $Q_\lambda \neq 0$ everywhere, as well as the equations $S_\tau + B_\lambda = S\alpha + G$, $Q_\tau + S_\lambda = Q\alpha + F$, $G_\tau = -S\alpha'$, $G_\lambda = Q\alpha' + F'$ for some function G and $(\cdot)' = d/d\tau$.*

PROOF. By (1-9b), Equation (1-1) is, in case (i), equivalent to

$$2\alpha\nabla d\tau + 2r = -(Q\alpha' + F')g, \quad (5-8)$$

where all the terms are Hermitian symmetric 2-tensor fields, and hence correspond, via g , to *complex-linear* bundle morphisms $TM \rightarrow TM$. Thus, (5-8) amounts to:

(f) *equalities of the images of both sides in (5-8) under ι_v and ι_w .*

The equality of the ι_v -images is, by (2-3b) and (1-3), the result of applying d , via (3-5), to the relation (1-9a) in item (i): $Y - Q\alpha = F$. This last relation and item (e), with $Y = \Delta\tau$ due to (1-4), show that condition (i) implies the equality $Q_\tau + S_\lambda = Q\alpha + F$ in item (ii). Defining G to be $S_\tau + B_\lambda - S\alpha$, we get $S_\tau + B_\lambda = S\alpha + G$. Next, the equality of the ι_w -images in (5-8) reads

$$\alpha dS - d\Delta\lambda = -(Q\alpha' + F')d\lambda. \quad (5-9)$$

In fact, the first term equals αdS since, for the two commuting gradients $v = \nabla\tau$ and $w = \nabla\lambda$, one has $2\nabla_w d\tau = d[g(v, w)]$ or, in local coordinates, $2w^k v_{jk} = w^k v_{jk} + v^k w_{jk} = (v^k w_k)_j$ and $S = \phi_{xu} = g(v, w)$ by (5-3) and (5-1). The second term is $-d\Delta\lambda$ due to item (a) and (2-8). By item (e), $G = S_\tau + B_\lambda - S\alpha = \Delta\lambda - S\alpha$, and so (5-9) becomes $\alpha dS - d(G + S\alpha) = -(Q\alpha' + F')d\lambda$, that is, according to (3-5),

$$dG = (Q\alpha' + F')d\lambda - Sd\alpha = (Q\alpha' + F')d\lambda - S\alpha'd\tau$$

or, in other words, $G_\tau = -S\alpha'$ and $G_\lambda = Q\alpha' + F'$. Consequently, condition (i) implies condition (ii), since (5-7) arises as the integrability conditions $Q_u - S_x = S_u - B_x = 0$ combined with (5-4), and the equality $dQ = Q_\tau d\tau + Q_\lambda d\lambda$ yields $dQ \wedge d\tau = -Q_\lambda d\tau \wedge d\lambda$.

Conversely, assuming condition (ii), we get condition (i) from item (f). Namely, as we saw above, the equality of the ι_v -images in (5-8) arises by applying d to $Y - Q\alpha = F$, that is—compare with item (e)—to $Q_\tau + S_\lambda = Q\alpha + F$. Also, (5-9) follows from condition (ii) and item (e):

$$\begin{aligned} \alpha dS - d\Delta\lambda &= \alpha dS - d(S_\tau + B_\lambda) = \alpha dS - d(S\alpha + G) = -dG - Sd\alpha = -dG - S\alpha' d\tau \\ &= G_\tau d\tau - dG = -G_\lambda d\lambda = -(Q\alpha' + F')d\lambda. \end{aligned}$$

This completes the proof. □

Note that items (e), (f), Theorem 5.1(ii) and (5-1) give

$$\Delta\tau = Q\alpha + F, \quad \Delta\lambda = S\alpha + G, \quad d_v\Pi = \Pi\Delta\tau, \quad d_w\Pi = \Pi\Delta\lambda.$$

6. Some linear algebra

We now proceed to discuss the first-order system equivalent, as we saw in the last section (Theorem 5.1), to the Kähler-potential problem, the solution of which amounts to proving Theorem E. The main result established here, Theorem 6.3, will lead—in Section 9—to a unique-extension property of integral lines, which results in applicability of the Cartan–Kähler theorem to our situation.

Theorem 5.1 reduces constructing local examples of special Ricci–Hessian equations (1-1)–(1-2) with $dQ \wedge d\tau \neq 0$, on Kähler surfaces, which is a fourth-order problem in the Kähler potential 2ϕ , to solving the following system of quasi-linear first-order partial differential equations, with subscripts representing partial derivatives:

- (a) $Q_\tau + S_\lambda - Q\alpha - F = 0$,
- (b) $S_\tau + B_\lambda - S\alpha - G = 0$,
- (c) $QB_\tau + SB_\lambda - SS_\tau - BS_\lambda = 0$,
- (d) $SQ_\tau + BQ_\lambda - QS_\tau - SS_\lambda = 0$,
- (e) $G_\tau + S\alpha' = 0$,
- (f) $G_\lambda - Q\alpha' - F' = 0$,

(6-1)

on which one imposes the additional conditions

$$QB > S^2, \quad Q > 0, \quad Q_\lambda \neq 0 \quad \text{everywhere.} \quad (6-2)$$

Generally, if Q, S, B are real-valued functions of the real variables τ, λ and subscripts denote partial differentiations, writing $d[(QB - S^2)^{-1}(Bd\tau - Sd\lambda)] = \Phi d\tau \wedge d\lambda$ and $d[(QB - S^2)^{-1}(Sd\tau - Qd\lambda)] = \Psi d\tau \wedge d\lambda$ at points where $QB \neq S^2$, one easily verifies that

$$\begin{bmatrix} \Phi \\ \Psi \end{bmatrix} = \begin{bmatrix} S & B \\ Q & S \end{bmatrix} \begin{bmatrix} QB_\tau + SB_\lambda - SS_\tau - BS_\lambda \\ SQ_\tau + BQ_\lambda - QS_\tau - SS_\lambda \end{bmatrix}. \quad (6-3)$$

For the remainder of this section, we treat the letter symbols in (6-1)–(6-2) as real variables, so as to derive some consequences of conditions (6-1)–(6-2) just by using linear algebra.

LEMMA 6.1. *If $(Q, S, B, G, Q_\tau, S_\tau, B_\tau, G_\tau, Q_\lambda, S_\lambda, B_\lambda, G_\lambda) \in \mathbb{R}^{12}$ satisfies the conditions (6-1a)–(6-1d), with some $(\alpha, F) \in \mathbb{R}^2$, then*

- (i) $QB_\tau + BQ_\tau - 2SS_\tau - (QB - S^2)\alpha - BF + SG = 0$,
- (ii) $QB_\lambda + BQ_\lambda - 2SS_\lambda - QG + SF = 0$.

(6-4)

PROOF. The left-hand side of (6-4i) is, obviously,

$$(QB_\tau + SB_\lambda - SS_\tau - BS_\lambda) + (Q_\tau + S_\lambda - Q\alpha - F)B - (S_\tau + B_\lambda - S\alpha - G)S,$$

and each sum in the parentheses vanishes due to (6-1). Similarly, (6-4ii) has the left-hand side

$$(SQ_\tau + BQ_\lambda - QS_\tau - SS_\lambda) + (S_\tau + B_\lambda - S\alpha - G)Q - (Q_\tau + S_\lambda - Q\alpha - F)S = 0 + 0 + 0.$$

□

With subscripts denoting partial differentiations, (6-1e–f) and (6-4) read

- (i) $dG = -S\alpha'd\tau + (Q\alpha' + F')d\lambda$,
- (ii) $d\Pi = (\Pi\alpha + BF - SG)d\tau + (QG - SF)d\lambda$ for $\Pi = QB - S^2$. (6-5)

Since $d\Pi = \Pi[(Q_\tau + S_\lambda)dx + (S_\tau + B_\lambda)du]$ due to item (f), one can also obtain (6-5ii) from (5-6iv), with $\Pi = QB - S^2$, and (6-1a–b).

LEMMA 6.2. *Let $(\dot{\tau}, \dot{\lambda}, Q, S, B, G) \in \mathbb{R}^6$ have $(\dot{\tau}, \dot{\lambda}) \neq (0, 0)$ and $QB > S^2$.*

- (a) *$QB > 0$ and $\Psi \neq 0$, where $\Psi = B\dot{\tau}^2 - 2S\dot{\tau}\dot{\lambda} + Q\dot{\lambda}^2$.*
- (b) *$(0, 0, 0, 0, 0, 0)$ is the only vector $(Q_\tau, S_\tau, B_\tau, Q_\lambda, S_\lambda, B_\lambda) \in \mathbb{R}^6$ with*

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 1 \\ 0 & -S & Q & 0 & -B & S \\ S & -Q & 0 & B & -S & 0 \\ \dot{\tau} & 0 & 0 & \dot{\lambda} & 0 & 0 \\ 0 & \dot{\tau} & 0 & 0 & \dot{\lambda} & 0 \\ 0 & 0 & \dot{\tau} & 0 & 0 & \dot{\lambda} \end{bmatrix} \begin{bmatrix} Q_\tau \\ S_\tau \\ B_\tau \\ Q_\lambda \\ S_\lambda \\ B_\lambda \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}.$$

- (c) *The first four rows of the above 7×6 matrix are linearly independent.*

PROOF. Assertion (a) follows as Ψ is positive or negative definite in $(\dot{\tau}, \dot{\lambda})$. Next, one easily verifies that, for Ψ as in assertion (a), and the rows \mathbf{r}_i of the above matrix,

$$(B\dot{\tau} - S\dot{\lambda})\dot{\lambda}\mathbf{r}_1 + \Psi\mathbf{r}_2 + \dot{\lambda}^2\mathbf{r}_4 - B\dot{\lambda}\mathbf{r}_5 + (2S\dot{\lambda} - B\dot{\tau})\mathbf{r}_6 = (0, 0, 0, 0, 0, \Psi).$$

Depending on whether $\dot{\lambda} \neq 0$ (or $\dot{\lambda} = 0$ and hence $\dot{\tau} \neq 0$), the 6×6 matrix with the rows $\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3 + S\mathbf{r}_2, \mathbf{r}_4 - S\mathbf{r}_1 + Q\mathbf{r}_2$ followed by $\mathbf{r}_6 - \dot{\tau}\mathbf{r}_2$ (or respectively by $\mathbf{r}_5 - \dot{\tau}\mathbf{r}_1$), and then by $(0, 0, 0, 0, 0, \Psi)$ displayed above, is upper triangular, with the diagonal entries $1, 1, Q, B, \dot{\lambda}, \Psi$ or respectively $1, 1, Q, B, -\dot{\tau}, \Psi$, all nonzero by assertion (a). These six new rows thus form a basis of \mathbb{R}^6 (so that the matrix has rank 6), while the first four are linear combinations of the first four original rows, which proves both assertions (b) and (c). \square

THEOREM 6.3. *Given vectors $(\dot{\tau}, \dot{\lambda}, Q, S, B, G) \in \mathbb{R}^6$ and $(\alpha, F, \alpha', F') \in \mathbb{R}^4$ with $(\dot{\tau}, \dot{\lambda}) \neq (0, 0)$ and $QB > S^2$, the set of all*

$$(Q_\tau, S_\tau, B_\tau, G_\tau, Q_\lambda, S_\lambda, B_\lambda, G_\lambda) \in \mathbb{R}^8 \quad (6-6)$$

satisfying (6-1) forms a two-dimensional affine subspace \mathcal{A} of \mathbb{R}^8 . Furthermore, the restriction to \mathcal{A} of the linear operator $\Phi : \mathbb{R}^8 \rightarrow \mathbb{R}^4$ sending (6-6) to

$$(\dot{Q}, \dot{S}, \dot{B}, \dot{G}) = (Q_\tau\dot{\tau} + Q_\lambda\dot{\lambda}, S_\tau\dot{\tau} + S_\lambda\dot{\lambda}, B_\tau\dot{\tau} + B_\lambda\dot{\lambda}, G_\tau\dot{\tau} + G_\lambda\dot{\lambda}) \quad (6-7)$$

is an affine isomorphism $\Phi : \mathcal{A} \rightarrow \mathcal{L}$ onto the two-dimensional affine subspace \mathcal{L} of \mathbb{R}^4 consisting of all $(\dot{Q}, \dot{S}, \dot{B}, \dot{G})$ such that

$$\begin{aligned}\dot{G} &= -Sa'\dot{\tau} + (Q\alpha' + F')\dot{\lambda}, \\ Q\dot{B} + B\dot{Q} - 2S\dot{S} &= [(QB - S^2)\alpha + BF - SG]\dot{\tau} + (QG - SF)\dot{\lambda}.\end{aligned}\quad (6-8)$$

PROOF. That $\mathcal{A} \subseteq \mathbb{R}^8$, or $\mathcal{L} \subseteq \mathbb{R}^4$, if nonempty, is an affine subspace, clearly follows since (6-1), or (6-8), is a system of nonhomogeneous linear equations imposed on (6-6) or respectively $(\dot{Q}, \dot{S}, \dot{B}, \dot{G})$. Also, \mathcal{A} is nonempty, and two-dimensional, being the pre-image of $(Q\alpha + F, Sa + G, 0, 0, -Sa', Q\alpha' + F')$ under the obvious linear operator $\mathbb{R}^8 \rightarrow \mathbb{R}^6$. Namely, this operator is surjective: it equals the direct sum of the identity operator $\mathbb{R}^2 \rightarrow \mathbb{R}^2$, acting on (G_τ, G_λ) , and an operator $\mathbb{R}^6 \rightarrow \mathbb{R}^4$, which has rank 4 due to Lemma 6.2(c). Next, Φ maps \mathcal{A} into \mathcal{L} , which one sees adding (6-1e), or (6-4i), times $\dot{\tau}$ to (6-1f), or (6-4ii), times $\dot{\lambda}$, and then using (6-7). Thus, \mathcal{L} is nonempty, and $\dim \mathcal{L} = 2$, the matrix of the homogeneous system associated with (6-8) having rank 2 since Lemma 6.2(a) gives $Q \neq 0$. Let $\mathcal{A}' \subseteq \mathbb{R}^8$ and $\mathcal{L}' \subseteq \mathbb{R}^4$ now be the vector subspaces parallel to \mathcal{A} and \mathcal{L} . Our assertion will thus follow once we establish injectivity of $\Phi : \mathcal{A} \rightarrow \mathcal{L}$, that is, injectivity of its linear part $\Phi : \mathcal{A}' \rightarrow \mathcal{L}'$. Equivalently, we need to show that zero is the only vector (6-6) lying in \mathcal{A}' and having the Φ -image $(0, 0, 0, 0)$ or, in other words, the only solution to the matrix equation in Lemma 6.2, with $(\dot{G}, G_\tau, G_\lambda) = (0, 0, 0)$. This, however, is precisely what Lemma 6.2(b) states. \square

7. The existence of solutions

After the preceding foray into linear algebra, we now return to treating (6-1) as a system of quasi-linear first-order partial differential equations with four unknown real-valued functions Q, S, B, G of the real variables τ, λ , subject to the additional conditions (6-2).

Subscripts again denote partial differentiations, while α and F are functions of the variable τ , also depending on three fixed real constants $\varepsilon, \theta, \kappa$, so that

$$\begin{aligned}2\alpha' + \alpha^2 &= 4\varepsilon, \text{ where } ()' = d/d\tau, \\ 4\varepsilon F &= \theta(2 - \tau\alpha) + 4\varepsilon\kappa\alpha \text{ if } \varepsilon \neq 0, \\ F &= \kappa\alpha - 2\theta/(3\alpha^2) \text{ when } \varepsilon = 0.\end{aligned}\quad (7-1)$$

In addition, it is natural to assume here that

$$\tau \text{ ranges over the domain of } \alpha \text{ (which is also the domain of } F). \quad (7-2)$$

Consequently, α and F satisfy the ordinary differential equations

$$\alpha'' + \alpha\alpha' = 0, \quad F'' = -F\alpha'. \quad (7-3)$$

THEOREM 7.1. *For any fixed α, F as in (7-1)–(7-2), real-analytic solutions $\mathbf{Z} = (Q, S, B, G)$ to (6-1)–(6-2) exist, locally, on a neighborhood of any $(\tau, \lambda) \in \mathbb{R}^2$ with the property (7-2).*

More precisely, one obtains a locally unique such solution \mathbf{Z} by prescribing \mathbf{Z} and the partial derivatives $\mathbf{Z}_\tau, \mathbf{Z}_\lambda$ real-analytically along an arbitrary real-analytic embedded curve $t \mapsto (\tau, \lambda) \in \mathbb{R}^2$, so as to satisfy (6-1), (6-2), (7-2), and the condition $\dot{\mathbf{Z}} = \dot{\tau}\mathbf{Z}_\tau + \dot{\lambda}\mathbf{Z}_\lambda$, where $(\cdot)' = d/dt$.

We prove Theorem 7.1 at the end of Section 10. As we point out in Remark 10.2, there is an infinite-dimensional freedom of choosing the data described in the second paragraph of Theorem 7.1.

8. The associated exterior differential system

By an *exterior differential system* on a manifold M , one means an ideal \mathcal{I} in the graded algebra Ω^*M , closed under exterior differentiation; its *integral manifolds* (or *integral elements*) are those submanifolds of M (or subspaces of its tangent spaces) on which every form in \mathcal{I} vanishes [4, pages 16 and 65]. When such objects have dimension 1 or 2, we call them *integral curves/surfaces* or *lines/planes*.

If $E \subseteq T_zM$ is a p -dimensional integral element of \mathcal{I} , one sets [4, pages 67–68]:

$$H(E) = \{v \in T_zM : \zeta(v, e_1, \dots, e_p) = 0 \text{ for all } \zeta \in \mathcal{I} \cap \Omega^{p+1}M\} \quad (8-1)$$

and $r(E) = \dim H(E) - (p + 1)$, for any basis e_1, \dots, e_p of E ,

so that $H(E)$ is a vector subspace of T_zM , not depending on e_1, \dots, e_p since

$$H(E) = \{v \in T_zM : \text{span}(v, E) \text{ is an integral element of } \mathcal{I}\}. \quad (8-2)$$

For fixed real constants $\varepsilon, \theta, \kappa$, let the open subset \mathcal{Y} of \mathbb{R}^6 consist of all points $(\tau, \lambda, Q, S, B, G) \in \mathbb{R}^6$ such that Q and $QB - S^2$ are both positive, while τ lies in the domains of α and F , chosen so as to satisfy (7-1). Consider now

the exterior differential system \mathcal{I} on this \mathcal{Y} generated
by the two 1-forms $dG + Sa'd\tau - (Qa' + F')d\lambda$ and
 $d(QB - S^2) - [(QB - S^2)\alpha + BF - SG]d\tau - (QG - SF)d\lambda$,
the 2-form $dQ \wedge d\lambda + d\tau \wedge dS - (Qa + F)d\tau \wedge d\lambda$,
the 2-form $dS \wedge d\lambda + d\tau \wedge dB - (Sa + G)d\tau \wedge d\lambda$,
their exterior derivatives, and the exterior derivatives of
 $(QB - S^2)^{-1}(Bd\tau - Sd\lambda)$ and $(QB - S^2)^{-1}(Sd\tau - Qd\lambda)$. (8-3)

The choice of \mathcal{I} is justified as follows. We want \mathcal{I} to consist of differential forms that are expected to vanish on all graphs of solutions to (6-1) with (6-2). Since such a graph is horizontal (in the sense that $d\tau \wedge d\lambda \neq 0$ on it), forms generating \mathcal{I} will result from multiplying by $d\tau \wedge d\lambda$ the left-hand sides in (6-1), along with those in its consequence (6-4), as including the latter changes nothing except for enriching the differential system. Each of the terms $\mathcal{E}Z_\tau$, or $\mathcal{E}Z_\lambda$, or Θ , where $Z \in \{Q, S, B, G\}$ (with various \mathcal{E} depending on Q, S, B, G , and Θ which may further depend also on a, F, a', F') then becomes $\mathcal{E}dZ \wedge d\lambda$, or $\mathcal{E}d\tau \wedge dZ$, or simply $\Theta d\tau \wedge d\lambda$.

The 2-forms in the fourth and fifth lines of (8-3) arise as above from (6-1a) and (6-1b), the exterior derivatives of the 1-forms in the last line—from (6-1c)

and (6-1d) via (6-3). The rest of (8-3) is based on an additional principle: if our exterior differential system ends up containing $\xi \wedge d\tau$ and $\xi \wedge d\lambda$, for some 1-form ξ , we are free to include ξ among the system's generators, as the horizontal integral surfaces/planes then obviously remain unaffected. The 1-form ξ in the second, or third, line of (8-3) corresponds in this way to (6-1e-f) or respectively (6-4).

Instead of invoking the general 'additional principle', one can also justify the second and third lines in (8-3) directly from the fact that (6-5) is a consequence of (6-1), which causes the two 1-forms to vanish on all graphs of solutions to (6-1).

9. The unique-extension theorem

In \mathbb{R}^6 with the coordinates $\tau, \lambda, Q, S, B, G$, given a subspace $E \subseteq \mathbb{R}^6$,

we call E horizontal when $(d\tau, d\lambda) : E \rightarrow \mathbb{R}^2$ is injective. (9-1)

REMARK 9.1. If E_1 is an integral line of \mathcal{I} in (8-3) and $E_1 \subseteq E_2$ for a unique horizontal integral plane E_2 , then E_2 is the only integral plane containing E_1 . Namely, another such plane E'_2 , being nonhorizontal, would intersect the kernel of $(d\tau, d\lambda)$ along a line. As $E_3 \subseteq H(E_1)$ for the vector subspace $H(E)$ in (8-2) and the three-dimensional span E_3 of E_2 and E'_2 , all planes in E_3 containing the line E_1 , other than E'_2 , would be horizontal integral planes, making E_2 nonunique.

REMARK 9.2. The only 1-forms in \mathcal{I} are, obviously, the functional combinations of those in the second and third lines of (8-3). Due to their linear independence at every point, the simultaneous kernel of these two 1-forms is a codimension-two distribution \mathcal{D} on \mathcal{Y} , that is, a vector subbundle of $T\mathcal{Y}$, and its fiber at any $(\tau, \lambda, Q, S, B, G) \in \mathcal{Y}$ consists of all $(\dot{\tau}, \dot{\lambda}, \dot{Q}, \dot{S}, \dot{B}, \dot{G}) \in \mathbb{R}^6$ with (6-8). Hence,

vectors $(\dot{\tau}, \dot{\lambda}, \dot{Q}, \dot{S}, \dot{B}, \dot{G})$ at $(\tau, \lambda, Q, S, B, G)$ spanning horizontal integral lines of \mathcal{I} are characterized by (6-8) and $(\dot{\tau}, \dot{\lambda}) \neq (0, 0)$, (9-2)

where α, F, α', F' satisfy (7-1)–(7-2).

THEOREM 9.3. *Every horizontal integral line of the system \mathcal{I} defined by (8-3) is contained in a unique integral plane of \mathcal{I} , and this unique plane is also horizontal.*

PROOF. Any horizontal plane in \mathbb{R}^6 has, by (9-1), a unique basis of the form

$$(1, 0, Q_\tau, S_\tau, B_\tau, G_\tau), \quad (0, 1, Q_\lambda, S_\lambda, B_\lambda, G_\lambda). \quad (9-3)$$

The span of (9-3) is an *integral plane of \mathcal{I}* if and only if all the 1-forms (and 2-forms) listed in (8-3) yield the value 0 when evaluated on both vectors in (9-3) or respectively on the pair (9-3). Due to the two final paragraphs of Section 8, and (6-3), this is equivalent to (6-1) and (6-4), and hence (Lemma 6.1) just to (6-1).

Every vector in (9-2) is a linear combination of a unique pair (9-3) satisfying (6-1): as the coefficients of the combination must be $\dot{\tau}$ and $\dot{\lambda}$, this is immediate from Theorem 6.3. In other words, every horizontal integral line of \mathcal{I} lies within a unique

horizontal integral plane. Remark 9.1 now allows us to drop the last occurrence of the word ‘horizontal’, completing the proof. \square

10. Existence of integral surfaces

The next fact—used below to derive our Theorem 7.1—is a special case of the celebrated Cartan–Kähler theorem [4, pages 81–82]. Since our phrasing differs from that of [4], we devote the next section to clarifying how our version amounts to adapting the one in [4] to our particular case.

The symbols \mathcal{Y} , \mathcal{I} and \mathcal{D} stand here for more general objects than those in Sections 8–9. The definition (9-1) of horizontality, for integral elements, is used more generally, as well as extended, in an obvious fashion, to integral manifolds.

THEOREM 10.1. *Let real-analytic functions τ, λ and 1-forms ξ_1, \dots, ξ_q on a manifold \mathcal{Y} , where $0 < q < \dim \mathcal{Y}$, have the property that $d\tau, d\lambda, \xi_1, \dots, \xi_q$ are linearly independent at every point. Denoting by \mathcal{D} and \mathcal{I} the distribution on \mathcal{Y} arising as the simultaneous kernel of the 1-forms ξ_1, \dots, ξ_q and respectively the exterior differential system on \mathcal{Y} generated by ξ_1, \dots, ξ_q and, possibly, some higher-degree forms, along with their exterior derivatives, let us suppose that*

$$\text{every horizontal integral line of } \mathcal{I}, \text{ at any point of } \mathcal{Y}, \text{ is contained in a unique integral plane of } \mathcal{I}. \quad (10-1)$$

Then, every horizontal real-analytic integral curve of \mathcal{I} is contained, locally, in a locally-unique horizontal real-analytic integral surface. Examples of such curves are provided by unparameterized integral curves of any real-analytic vector field without zeros forming a horizontal local section of the vector bundle \mathcal{D} over \mathcal{Y} . Also,

$$\text{integral lines of } \mathcal{I} \text{ are the same as lines tangent to } \mathcal{D}. \quad (10-2)$$

REMARK 10.2. Due to Theorem 9.3, our \mathcal{Y} and \mathcal{I} , introduced in Section 8, satisfy the hypotheses of Theorem 10.1, with $q = 2$, the coordinate functions τ, λ , and the two 1-forms in the second and third lines of (8-3). Therefore, our \mathcal{D} (see Remark 9.2) then corresponds to \mathcal{D} in Theorem 10.1, and hence satisfies (10-2). Horizontal integral curves of our \mathcal{I} thus are, by (9-2), precisely those curves that have parameterizations $t \mapsto (\tau, \lambda, \mathbf{Z}) = (\tau, \lambda, Q, S, B, G) \in \mathcal{Y}$ with (6-8), where $(\cdot)^\cdot = d/dt$, and $(\dot{\tau}, \dot{\lambda}) \neq (0, 0)$ for all t . Choosing such a curve as in the sentence preceding (10-2), we obtain the additional data $\mathbf{Z}_\tau, \mathbf{Z}_\lambda$ required in Theorem 7.1 by applying to $\dot{\mathbf{Z}}$ the inverse of the affine isomorphism $\Phi : \mathcal{A} \rightarrow \mathcal{L}$ of Theorem 6.3. This clearly results in the infinite-dimensional freedom of choices mentioned at the end of Section 6.

PROOF OF THEOREM 7.1. The image of the mapping $t \mapsto (\tau, \lambda, \mathbf{Z}) \in \mathcal{Y}$ in the second paragraph of Theorem 7.1 is a horizontal real-analytic integral curve of \mathcal{I} . In fact, horizontality follows since $t \mapsto (\tau, \lambda)$ is an embedding, while, as $\dot{\mathbf{Z}} = \dot{\tau}\mathbf{Z}_\tau + \dot{\lambda}\mathbf{Z}_\lambda$, the resulting tangent directions are integral lines of \mathcal{I} as an immediate consequence of Theorem 9.3 combined with (9-2).

The integral surface of \mathcal{I} arising in Theorem 10.1, being horizontal (Theorem 9.3), forms, locally, the graph of a function $(\tau, \lambda) \mapsto \mathbf{Z} = (Q, S, B, G)$, which is a solution to (6-1) according to the description of \mathcal{I} in (8-3) and the paragraph following (8-3). To realize the condition $Q_\lambda \neq 0$ required in (6-2), we solve (6-1), at a given point $(\tau, \lambda, Q, S, B, G) \in \mathcal{Y}$, by setting $(Q_\tau, Q_\lambda) = (0, 1)$, which uniquely determines $S_\tau, B_\tau, G_\tau, S_\lambda, B_\lambda, G_\lambda$, and then choosing the quadruple (6-7), with fixed $(\tau, \lambda) \neq (0, 0)$, associated with the resulting octuple (6-6). \square

Under the assumptions of Theorem 10.1, let $k = \dim \mathcal{Y}$. For all p -dimensional horizontal integral elements $E = E_p$ of \mathcal{I} , with $p \in \{0, 1\}$, and for $r(E) = \dim H(E) - (p + 1)$ in (8-1):

- (a) the integer $r(E)$ has a fixed nonnegative value, namely,
- (b) $\dim H(E_0) = k - q$ and $r(E_0) = k - q - 1$ when $p = 0$,
- (c) $\dim H(E_1) = 2$ and $r(E_1) = 0$ in the case where $p = 1$.

This is obvious from (10-2) or respectively (10-1).

11. Where Theorem 10.1 comes from

Here is the Cartan–Kähler theorem, cited *verbatim* from [4, pages 81–82]:

Let $\mathcal{I} \subset \mathcal{Q}^(M)$ be a real analytic differential ideal. Let $P \subset M$ be a connected, p -dimensional, real analytic, Kähler-regular integral manifold of \mathcal{I} .*

Suppose that $r = r(P)$ is a nonnegative integer. Let $R \subset M$ be a real analytic submanifold of M which is of codimension r , which contains P , and which satisfies the condition that $T_x R$ and $H(T_x R)$ are transverse in $T_x M$ for all $x \in P$.

Then, there exists a real analytic integral manifold of \mathcal{I} , X , which is connected and $(p + 1)$ -dimensional and which satisfies $P \subset X \subset R$. This manifold is unique in the sense that any other real analytic integral manifold of \mathcal{I} with these properties agrees with X on an open neighborhood of P .

As we verify in the following paragraphs, the hypotheses of our Theorem 10.1 imply those listed above, for $(p, r) = (1, 0)$, the manifolds M, R above which are both replaced by our \mathcal{Y} , and the same ideal \mathcal{I} as ours. By our \mathcal{Y} and \mathcal{I} , we mean the ‘general’ ones (see the three lines preceding Theorem 10.1), rather than the very special choices of \mathcal{Y} and \mathcal{I} made in Section 8.

Furthermore, P mentioned above is our (arbitrary) horizontal real-analytic integral curve of \mathcal{I} . The resulting manifold X corresponds to the horizontal real-analytic integral surface of \mathcal{I} claimed to exist in Theorem 10.1.

We now proceed to explain why our horizontal integral curve must automatically be Kähler-regular [4, page 81], meaning that its tangent lines are all Kähler-regular in the sense of [4, page 68, Definition 1.7]. To verify this last claim, we first apply Cartan’s test [4, page 74, Theorem 1.11]. Namely, in the notation of [4, page 74, Theorem 1.11], $n = 1$ (as we are dealing with tangent *lines*). Due to the relation $\dim H(E_0) = k - q$ in (10-3b), and (10-2), $H(E_0)$ is of codimension q in the tangent space of \mathcal{Y} containing

it, which is the same as the codimension, in the $(2k - 1)$ -dimensional Grassmann manifold $\text{Gr}_1 \mathcal{Y}$ of lines tangent to \mathcal{Y} , of the $(2k - q - 1)$ -dimensional submanifold $V_1(\mathcal{I})$ formed by all integral lines of \mathcal{I} . Cartan's test thus shows that every line E_1 tangent to our horizontal integral curve is *ordinary* [4, page 73, Definition 1.9]. The Kähler-regularity of E_1 now trivially follows, as r in [4, pages 67–68] has the constant value 0 according to (10-3c). This is also the value $r = r(P)$ in the italicized statement cited above from [4] (compare with [4, pages 81–82, the lines preceding Theorem 2.2]).

12. Proof of Theorem E

Let ∇ (or g) be a connection (or a pseudo-Riemannian metric) on a C^∞ manifold M . We call ∇ or g *real-analytic* if, in a suitable coordinate system around every point of M , its components Γ_{jk}^l (or g_{jk}) are real-analytic functions of the coordinates. The C^∞ structure of M then contains a unique real-analytic structure (maximal atlas) making ∇ or g real-analytic. (The atlas consists of all coordinate systems just mentioned; their mutual transition mappings are real-analytic due to real-analyticity of affine mappings, or isometries, between manifolds with real-analytic connections/metrics, which follows since such mappings appear linear in geodesic coordinates.) Real-analyticity of a metric g obviously implies that of its Levi–Civita connection ∇ (and *vice versa*, since $\nabla g = 0$).

For a real-analytic (Riemannian) Kähler metric g on a complex manifold M , the unique real-analytic structure described above coincides with that induced by the complex structure of M . In fact, local holomorphic coordinate functions, being g -harmonic, must be real-analytic relative to the former structure, as a consequence of the standard regularity theory of elliptic partial differential equations applied to the g -Laplacian Δ .

PROOF OF THEOREM E. Combining Theorems 7.1 and 5.1, we obtain the first assertion of Theorem E.

For the second one, we invoke the existence results of [6, 22]. In both cases, $dQ \wedge d\tau \neq 0$ somewhere, and the metric is real-analytic. The former claim follows, for instance, since a compact Kähler surface with a nontrivial holomorphic gradient $\nabla\tau$ having $dQ \wedge d\tau = 0$ identically for $Q = g(\nabla\tau, \nabla\tau)$ must necessarily [10, Section 1] be biholomorphic to \mathbb{CP}^2 or a \mathbb{CP}^1 bundle over \mathbb{CP}^1 (rather than the two-point blow-up of \mathbb{CP}^2). The latter, in the case of [22], is due to a general reason: all Ricci solitons are real-analytic [8, Lemma 3.2]. So are, however, all Riemannian Einstein metrics [13, Theorem 5.2], and the Chen–LeBrun–Weber metric of [6] is conformal to an Einstein metric \hat{g} , while again, for a general reason [9, page 417, Proposition 3(ii)], the conformal change leading from \hat{g} to g has a canonical form (up to a constant factor, it is the multiplication by the cubic root of the norm-squared of the self-dual Weyl tensor). This causes g to be real-analytic as well. \square

13. The analytic-continuation phenomenon

We elaborate here on the plausibility of small deformations mentioned in the lines following Theorem E, beginning with the coth-cot analytic continuation. The real-analytic function $\mathbb{R} \ni y \mapsto y^{-1}\tanh y$, with the value 1 at $y = 0$, being even, has the form $\Sigma(y^2)$ for some real-analytic function Σ . Now, $(\varepsilon, \tau) \mapsto \beta_\varepsilon(\tau) = \tau\Sigma(\varepsilon\tau^2)$ is a real-analytic function on an open subset of \mathbb{R}^2 and $\beta_\varepsilon(\tau)$ equals $\varepsilon^{-1/2} \tanh(\varepsilon^{1/2}\tau)$, or τ , or $|\varepsilon|^{-1/2} \tan(|\varepsilon|^{1/2}\tau)$, depending on whether $\varepsilon > 0$, or $\varepsilon = 0$, or $\varepsilon < 0$. For $\alpha_\varepsilon(\tau) = 2/\beta_\varepsilon(\tau)$, the analogous expressions are

$$2\varepsilon^{1/2} \coth(\varepsilon^{1/2}\tau) \text{ (if } \varepsilon > 0\text{), } \quad 2/\tau \text{ (if } \varepsilon = 0\text{), } \quad 2|\varepsilon|^{1/2} \cot(|\varepsilon|^{1/2}\tau) \text{ (if } \varepsilon < 0\text{).}$$

All α_ε with $\varepsilon > 0$, as well as those with $\varepsilon < 0$, are thus affine (in fact, linear) modifications (see Remark C) of α_1 or respectively α_{-1} , and $\alpha_0(\tau) = 2/\tau$.

For a tanh-coth analytic-continuation argument, we define $(t, \tau) \mapsto \alpha_t(\tau)$ by $\alpha_t(\tau) = 2(e^\tau - te^{-\tau})/(e^\tau + te^{-\tau})$. Thus, with q such that $2q = \log|t|$, if $t > 0$ (or $t < 0$), $\alpha_t(\tau) = 2\tanh(\tau - q)$ or respectively $\alpha_t(\tau) = 2\coth(\tau - q)$. Again, all α_t for $t > 0$, or those with $t < 0$, are affine (this time, translational) modifications of α_1 , or of α_{-1} , while $\alpha_0(\tau) = 2$.

References

- [1] A. Barros and E. Ribeiro, ‘Some characterizations for compact almost Ricci solitons’, *Proc. Amer. Math. Soc.* **140** (2012), 1033–1040.
- [2] A. L. Besse, *Einstein Manifolds*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge / A Series of Modern Surveys in Mathematics, 10 (Springer-Verlag, Berlin–Heidelberg–New York, 1987).
- [3] J. Bolton, ‘Transnormal systems’, *Quart. J. Math. Oxford Ser. (2)* **24** (1973), 385–395.
- [4] R. L. Bryant, S. S. Chern, R. B. Gardner, H. L. Goldschmidt and P. A. Griffiths, *Exterior Differential Systems*, Mathematical Sciences Research Institute Publications, 18 (Springer-Verlag, New York, 1991).
- [5] E. Calabi, ‘Extremal Kähler metrics’, in: *Seminar on Differential Geometry*, Annals of Mathematics Studies, 102 (ed. S.-T. Yau) (Princeton University Press, Princeton, NJ, 1982), 259–290.
- [6] X. Chen, C. LeBrun and B. Weber, ‘On conformally Kähler, Einstein manifolds’, *J. Amer. Math. Soc.* **21** (2008), 1137–1168.
- [7] B. Chow, ‘Ricci flow and Einstein metrics in low dimensions’, in: *Surveys in Differential Geometry, Volume VI* (eds. C. LeBrun and M. Wang) (International Press, Boston, MA, 1999), 187–220.
- [8] A. S. Dancer and M. Y. Wang, ‘On Ricci solitons of cohomogeneity one’, *Ann. Global Anal. Geom.* **39** (2011), 259–292.
- [9] A. Derdzinski, ‘Self-dual Kähler manifolds and Einstein manifolds of dimension four’, *Compos. Math.* **49** (1983), 405–433.
- [10] A. Derdzinski, ‘Killing potentials with geodesic gradients on Kähler surfaces’, *Indiana Univ. Math. J.* **61** (2012), 1643–1666.
- [11] A. Derdzinski and G. Maschler, ‘Local classification of conformally-Einstein Kähler metrics in higher dimensions’, *Proc. Lond. Math. Soc. (3)* **87** (2003), 779–819.
- [12] A. Derdzinski and G. Maschler, ‘Special Kähler–Ricci potentials on compact Kähler manifolds’, *J. reine angew. Math.* **593** (2006), 73–116.

- [13] D. M. DeTurck and J. L. Kazdan, ‘Some regularity theorems in Riemannian geometry’, *Ann. Sci. Éc. Norm. Supér. (4)* **14** (1981), 249–260.
- [14] C. T. J. Dodson, M. Trinidad Pérez and M. E. Vázquez-Abal, ‘Harmonic–Killing vector fields’, *Bull. Belg. Math. Soc. Simon Stevin* **9** (2002), 481–490.
- [15] R. S. Hamilton, ‘The Ricci flow on surfaces’, in *Mathematics and General Relativity, Santa Cruz, CA, 1986*, Contemporary Mathematics, 71 (ed. J. A. Isenberg) (American Mathematical Society, Providence, RI, 1988), 237–262.
- [16] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry, Volume I* (Interscience, New York, 1963).
- [17] G. Maschler, ‘Special Kähler–Ricci potentials and Ricci solitons’, *Ann. Global Anal. Geom.* **34** (2008), 367–380.
- [18] R. Miyaoka, ‘Transnormal functions on a Riemannian manifold’, *Differential Geom. Appl.* **31** (2013), 130–139.
- [19] O. Nouhaud, ‘Transformations infinitésimales harmoniques’, *C. R. Acad. Sci. Paris Sér. A-B* **274** (1972), A573–A576.
- [20] S. Pigola, M. Rigoli, M. Rimoldi and A. G. Setti, ‘Ricci almost solitons’, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (4)* **10** (2011), 757–799.
- [21] Q. M. Wang, ‘Isoparametric functions on Riemannian manifolds, I’, *Math. Ann.* **277** (1987), 639–646.
- [22] X. J. Wang and X. Zhu, ‘Kähler–Ricci solitons on toric manifolds with positive first Chern class’, *Adv. Math.* **188** (2004), 87–103.

ANDRZEJ DERDZINSKI, Department of Mathematics,
The Ohio State University, 231 W. 18th Avenue, Columbus, OH 43210, USA
e-mail: andrzej@math.ohio-state.edu

PAOLO PICCIONE, Department of Mathematics,
School of Sciences, Great Bay University, Dongguan, Guangdong 523000, PR China

Permanent address: Departamento de Matemática,
Instituto de Matemática e Estatística, Universidade de São Paulo, Rua do Matão 1010,
CEP 05508-900, São Paulo, SP, Brazil
e-mail: paolo.piccione@usp.br