



# Affine vector fields on compact pseudo-Kähler manifolds

Andrzej Derdzinski<sup>1</sup>

Received: 22 December 2024 / Accepted: 12 December 2025  
© The Author(s) 2025

## Abstract

It is known that a Killing field on a compact pseudo-Kähler manifold is necessarily (real) holomorphic, as long as the manifold satisfies some relatively mild additional conditions. We provide two further proofs of this fact and discuss the natural open question whether the same conclusion holds for affine—rather than Killing—vector fields. The question cannot be settled by invoking the Killing case: Boubel and Mounoud [Trans. Amer. Math. Soc. 368, 2016, 2223–2262] constructed examples of non-Killing affine vector fields on compact pseudo-Riemannian manifolds. We show that an affine vector field  $v$  is necessarily symplectic, and establish some algebraic and differential properties of the Lie derivative of the metric along  $v$ , such as its being parallel, antilinear and nilpotent as an endomorphism of the tangent bundle. As a consequence, the answer to the above question turns out to be ‘yes’ whenever the underlying manifold admits no nontrivial holomorphic quadratic differentials, which includes the case of compact almost homogeneous complex manifolds with nonzero Euler characteristic.

**Keywords** Compact pseudo-Kähler manifold · Affine vector field · Real-holomorphic vector field

**Mathematics Subject Classification** Primary 53C50 · Secondary 53C56

## 1 Introduction

A *pseudo-Kähler manifold* is a pseudo-Riemannian manifold endowed with a parallel almost-complex structure  $J$ , making the metric Hermitian. This is well known to imply integrability of  $J$ , via the Newlander-Niren-

---

Dedicated to Paolo Piccione on the occasion of his 60th birthday.

---

✉ Andrzej Derdzinski  
andrzej@math.ohio-state.edu

<sup>1</sup> Department of Mathematics, The Ohio State University, Columbus 43210, OH, USA

berg theorem: the Nijenhuis tensor  $N$  sends vector fields  $v, w$  to the vector field  $N(v, w) = [v, w] + J[Jv, w] + J[v, Jw] - [Jv, Jw]$ , so that  $N(v, w) = [J\nabla_v J - \nabla_{Jv} J]w + [\nabla_{Jw} J - J\nabla_w J]v$  for any torsion-free connection  $\nabla$ . Thus,  $N = 0$  whenever  $\nabla J = 0$ .

It is also well known [1, pp. 60–61] that Killing fields on compact Riemannian Kähler manifolds are necessarily (real) holomorphic, compactness being essential (as illustrated by flat manifolds). This remains valid, under some additional assumptions, in the pseudo-Kähler case [3, 8].

A vector field  $v$  on a manifold endowed with a connection  $\nabla$  is said to be *affine* if its local flow preserves  $\nabla$ . When  $\nabla$  is the Levi-Civita connection of a Riemannian metric, such  $v$  are usually Killing fields, with very few—always noncompact—exceptions [5, Ch. IV]. However, Boubel and Mounoud [2] provided examples of compact pseudo-Riemannian manifolds admitting non-Killing affine vector fields. Their examples are not of the pseudo-Kähler type, which raises a question: are the affine-to-Killing and affine-to-holomorphic implications true for compact pseudo-Kähler manifolds?

This question remains open. The present paper establishes some properties of the Lie derivative  $\mathcal{L}_v g$  for an affine vector field  $v$  on a compact pseudo-Kähler manifold  $(M, g)$  with the  $\partial\bar{\partial}$  property (Theorem 6.1): in addition to being obviously parallel,  $\mathcal{L}_v g$  is also antilinear and nilpotent as an endomorphism of  $TM$ , while  $v$  is a Killing field if and only if it is real holomorphic. Also,  $\mathcal{L}_v \omega = 0$  for the Kähler form  $\omega$  (Corollary 4.2).

Finally, as a partial answer to the above question, we show that  $v$  must be real holomorphic when  $M$  admits no nontrivial holomorphic quadratic differentials (Theorem 7.2) and hence, in particular, when  $M$  is almost homogeneous and has nonzero Euler characteristic (Corollary 7.3). We also provide, in Theorem 7.1, a pseudo-Riemannian analog of a Hodge decomposition for the 1-form  $g(v, \cdot)$ , and observe that it coincides with the Riemannian Hodge decomposition of  $g(v, \cdot)$  relative to any Riemannian Kähler metric  $h$  on  $M$ , if such  $h$  exists.

## 2 Preliminaries

Manifolds and mappings are assumed smooth, the former also connected.

Let  $(M, g)$  be a pseudo-Riemannian manifold. We write  $\beta \sim B$  when tensor fields  $\beta$  and  $B$  of types  $(0, 2)$  and  $(1, 1)$  are related by  $\beta = g(B\cdot, \cdot)$ . Thus,  $g \sim \text{Id}$ . On a pseudo-Kähler manifold

$$\omega \sim J \text{ for the Kähler form } \omega \text{ and the complex-structure tensor } J. \quad (2.1)$$

If  $\beta \sim B$ , as defined above, and we set

$$A = \nabla v \quad (2.2)$$

for a fixed vector field  $v$ , one easily sees that

$$\mathcal{L}_v\beta \sim \nabla_v B + BA + A^*B, \tag{2.3}$$

$A^*$  being the pointwise  $g$ -adjoint of  $A$ . Two obvious special cases are

$$\text{a) } \mathcal{L}_v g \sim A + A^*, \quad \text{b) } \mathcal{L}_v \omega \sim JA + A^*J, \tag{2.4}$$

the latter in the pseudo-Kähler case. Assuming (2.2), we obviously get

$$d[g(v, \cdot)] \sim A - A^*. \tag{2.5}$$

If  $\beta \sim B$  for a real differential 2-form on a complex manifold,

$$\beta \text{ is a } (1, 1)\text{-form if and only if } [J, B] = 0, \text{ as both conditions are clearly equivalent to } \beta(J\cdot, J\cdot) = \beta. \tag{2.6}$$

Given a pseudo-Riemannian manifold  $(M, g)$ , using  $J$ ,

$$\text{we treat } TM \text{ as a complex vector bundle.} \tag{2.7}$$

Cartan’s homotopy formula  $\mathcal{L}_v = v_d + dv_v$  for  $\mathcal{L}_v$  acting on differential forms [6, Thm. 14.35, p. 372] and the Leibniz rule  $\mathcal{L}_v[\nabla\Theta] = [\mathcal{L}_v\nabla]\Theta + \nabla[\mathcal{L}_v\Theta]$  imply that, for any vector field  $v$  on a manifold,

$$\mathcal{L}_v\omega = d[\omega(v, \cdot)] \text{ if } \omega \text{ is a closed differential form,} \tag{2.8}$$

while, whenever  $v$  happens to be affine relative to a connection  $\nabla$ ,

$$\nabla[\mathcal{L}_v\Theta] = 0 \text{ if } \Theta \text{ is a tensor field with } \nabla\Theta = 0. \tag{2.9}$$

**Remark 2.1** Any constant-rank twice-covariant symmetric tensor field  $\beta$  on a manifold has the same algebraic type at all points: its positive and negative indices, being lower semicontinuous, with a constant sum, must be locally constant.

**Remark 2.2** Due to the Leibniz rule, for any vector field  $v$  on a pseudo-Kähler manifold,  $\mathcal{L}_v J = [J, A]$ , where  $A = \nabla v$ , so that  $v$  is real holomorphic if and only if  $\nabla v$  commutes with  $J$ . On the other hand, holomorphic complex-valued functions  $\phi$  on a complex manifold  $M$  are characterized by the Cauchy-Riemann condition  $(d\phi)J = id\phi$ , where  $(d\phi)J$  denotes the composite bundle morphism  $TM \rightarrow TM \rightarrow M \times \mathbb{C}$ .

### 3 The $\partial\bar{\partial}$ property

Every compact complex manifold admitting a Riemannian Kähler metric has the following  $\partial\bar{\partial}$  property, also referred to as *the  $\partial\bar{\partial}$  lemma* [9, Prop. 6.17 on p. 144]: given integers  $p, q \geq 0$ , any closed  $\partial$ -exact or  $\bar{\partial}$ -exact  $(p, q)$ -form equals  $\partial\bar{\partial}\lambda$  for some  $(p - 1, q - 1)$ -form  $\lambda$ . Then, since the exactness of a  $(p, 0)$ -form amounts to its  $\bar{\partial}$ -exactness, and implies its closedness,

$$M \text{ admits no nonzero exact } (p, 0)\text{- or } (0, p)\text{-forms.} \tag{3.1}$$

As a special case, on a compact complex  $\partial\bar{\partial}$  manifold  $M$ ,

$$\text{every exact real } (1, 1)\text{-form } \alpha \text{ equals } i\partial\bar{\partial}\phi \text{ for some } \phi : M \rightarrow \mathbb{R}. \tag{3.2}$$

In fact, writing  $\alpha = d\xi = \partial\xi + \bar{\partial}\xi$ , with a real 1-form  $\xi$ , we see that  $\partial\xi$  and  $\bar{\partial}\xi$  are both closed:  $d\partial\xi = \bar{\partial}\partial\xi = \bar{\partial}d\xi = \bar{\partial}\alpha = 0 = \partial\alpha = \partial\bar{\partial}\xi = d\bar{\partial}\xi$ , since  $d = \partial + \bar{\partial}$ , while  $\partial^2 = \bar{\partial}^2 = 0$  and  $\alpha$ , being closed, has  $\partial\alpha = \bar{\partial}\alpha = 0$ . Also, as  $\alpha$  is a  $(1, 1)$ -form, so must be  $\partial\xi$  and  $\bar{\partial}\xi$ . In view of the  $\partial\bar{\partial}$  property, the  $(1, 1)$ -forms  $\partial\xi$  and  $\bar{\partial}\xi$ , being closed and  $\partial$ -exact or  $\bar{\partial}$ -exact, equal  $\partial\bar{\partial}\lambda$  and  $\partial\bar{\partial}\mu$  for some functions  $\lambda, \mu$ , so that  $\alpha = \text{Re } \alpha = i\partial\bar{\partial}\phi$ , where  $\phi = \text{Im}(\lambda + \mu)$ .

On the other hand, whenever  $\phi : M \rightarrow \mathbb{R}$ ,

$$2i\partial\bar{\partial}\phi = -d[(d\phi)J], \tag{3.3}$$

$(d\phi)J$  being the composite bundle morphism  $TM \rightarrow TM \rightarrow M \times \mathbb{R}$ . Consequently,

$$-2\alpha(J\cdot, \cdot) = \theta(J\cdot, J\cdot) + \theta(\cdot, \cdot) \text{ if } \alpha = i\partial\bar{\partial}\phi \text{ and } \theta = \nabla d\phi, \tag{3.4}$$

where  $\nabla$  is any torsion-free connection on  $M$  with  $\nabla J = 0$ . Whether or not such  $\nabla$  exists, at any critical point  $z$  of a function  $\phi : M \rightarrow \mathbb{R}$ , by (3.3),

$$-2\alpha(J\cdot, \cdot) = \theta(J\cdot, J\cdot) + \theta(\cdot, \cdot) \text{ if } \alpha = i\partial\bar{\partial}\phi \text{ and } \theta = \text{Hess}_z\phi. \tag{3.5}$$

**Lemma 3.1** *A compact complex manifold  $M$  satisfying the special case (3.2) of the  $\partial\bar{\partial}$  condition admits no nonzero constant-rank real-valued exact  $(1, 1)$ -forms.*

**Proof** Applying (3.2) and (3.5) to a constant-rank real-valued exact  $(1, 1)$ -form  $\alpha$ , at a point  $z \in M$  where  $\phi$  assumes its maximum (or, minimum) value, we see that the symmetric 2-tensor field  $\alpha(J\cdot, \cdot)$  is positive (or, negative) semidefinite at  $z$ . Remark 2.1 applied to  $\beta = \alpha(J\cdot, \cdot)$ , which has constant rank (equal to the rank of  $\alpha$ ), implies both positive and negative semidefiniteness of  $\beta$  at all points, so that  $\beta = 0$  everywhere.  $\square$

### 4 Consequences of the Hodge decomposition

For each cohomology space  $H^p(M, \mathbb{C})$  of a compact complex manifold  $M$  with the  $\partial\bar{\partial}$  property, denoting by  $H^{r,s}M$  the space of cohomology classes of closed complex-valued  $(r, s)$ -forms, one has the Hodge decomposition

$$H^p(M, \mathbb{C}) = H^{p,0}M \oplus H^{p-1,1}M \oplus \dots \oplus H^{1,p-1}M \oplus H^{0,p}M. \tag{4.1}$$

See, e.g., [4, p. 296, subsect. (5.21)].

**Lemma 4.1** *Let  $\zeta$  be an exact  $\nabla$ -parallel complex-valued  $p$ -form on a compact complex  $\partial\bar{\partial}$  manifold  $M$  with a torsion-free connection  $\nabla$  such that  $\nabla J = 0$ .*

- (i)  $\zeta$  has zero  $(p, 0)$  and  $(0, p)$  components.
- (ii) The  $(r, s)$  components of  $\zeta$ ,  $r + s = p$ , are all exact and  $\nabla$ -parallel.
- (iii)  $\zeta = 0$  when  $p = 2$ .

**Proof** The decomposition of the bundle of complex-valued exterior  $p$ -forms on  $M$  into its  $(r, s)$  summands, with  $r + s = p$ , is invariant under parallel transports, since  $J$  uniquely determines the decomposition and  $\nabla J = 0$ . The components  $\zeta^{r,s}$  of the decomposition of  $\zeta$  are thus all  $\nabla$ -parallel, and hence closed. The resulting cohomology relation  $\sum_{r,s} [\zeta^{r,s}] = [\zeta] = 0$  gives, by (4.1),  $[\zeta^{r,s}] = 0$  whenever  $r + s = p$ , proving (ii), while (i) follows from (ii) and (3.1). Lemma 3.1 and (i)–(ii) now easily yield (iii). □

We have an obvious consequence of (2.8)–(2.9) and Lemma 4.1(iii).

**Corollary 4.2** *Let  $v$  be an affine vector field on a compact pseudo-Kähler  $\partial\bar{\partial}$  manifold  $(M, g)$  with the Kähler form  $\omega = g(J\cdot, \cdot)$ . Then  $\mathcal{L}_v\omega = 0$ .*

### 5 The case of Killing fields

The paper [3] provides two different proofs of the fact that, on a compact pseudo-Kähler  $\partial\bar{\partial}$  manifold, all Killing fields are real holomorphic.

Our preceding discussion gives rise to two more simple proofs of this fact. As

$$\mathcal{L}_v [g(J\cdot, \cdot)] = g(\mathcal{L}_v J\cdot, \cdot) \text{ when } \mathcal{L}_v g = 0, \tag{5.1}$$

and  $\omega = g(J\cdot, \cdot)$ , one new proof comes directly from Corollary 4.2.

For the other new proof, note that  $\zeta = \mathcal{L}_v\omega$ , being exact and parallel by (2.8) and (2.9), must—due to Lemma 4.1(i)—be a  $(1, 1)$ -form. Since (5.1) gives  $\mathcal{L}_v\omega \sim \mathcal{L}_v J$ , (2.6) implies that  $J$  and  $C = \mathcal{L}_v J$  commute. However, they also anticommute:  $0 = -\mathcal{L}_v \text{Id} = \mathcal{L}_v J^2 = CJ + JC$ . Consequently,  $\mathcal{L}_v J = C = 0$ .

The first proof in [3], at the end of Sect. 3, proceeds as follows. By (2.9) above,  $\mathcal{L}_v J$  is parallel, and hence so is the  $(2, 0)$ -form  $\zeta = g(\mathcal{L}_v J \cdot, \cdot) - ig((\mathcal{L}_v J)J \cdot, \cdot)$ , which makes  $\zeta$  closed as well. However,  $\zeta$  is also  $\partial$ -exact, namely—see [3, Lemma 3.4]—equal to  $\partial[g(Jv, \cdot) - ig(v, \cdot)]$ . Combined with its closedness and the  $\partial\bar{\partial}$  property, this, according to [3, Lemma 3.1], gives  $\zeta = 0$ .

The second proof in [3], in Sect. 4, uses the parallel  $(2, 0)$ -form  $\zeta$  of the last paragraph. By (2.8),  $\text{Re } \zeta = \mathcal{L}_v \omega$  is exact. This makes  $[i\zeta] \in H^{2,0}M$  a real cohomology class, equal to its conjugate lying in  $H^{0,2}M$ , and so  $\zeta = 0$ , as  $H^{2,0}M$ ,  $H^{0,2}M$  and  $H^{1,1}M$  are direct summands of  $H^2(M, \mathbb{C})$ ,

### 6 The four components of the covariant derivative

On a pseudo-Kähler manifold  $(M, g)$ , the operation  $A \mapsto JAJ$  applied to bundle morphisms  $A : TM \rightarrow TM$  obviously commutes with  $A \mapsto A^*$  and, as both are involutions, every  $A$  is decomposed into four components (complex-linear self-adjoint, complex-linear skew-adjoint, antilinear self-adjoint, antilinear skew-adjoint). In the case where  $A = \nabla v$  for an affine vector field  $v$  on a compact pseudo-Kähler  $\partial\bar{\partial}$  manifold  $(M, g)$ , two of the four components—the first and last ones—are absent, according to the assertions (6.1-b) and (6.1-c) below, while the third one has rather special algebraic and differential properties, cf. (6.1-d).

**Theorem 6.1** *Let  $v$  be an affine vector field on a compact pseudo-Kähler  $\partial\bar{\partial}$  manifold  $(M, g)$ . Then, for  $A = \nabla v$ ,*

- a)  $A^* = JAJ$ ,
  - b)  $A - A^*$  commutes with  $J$ ,
  - c)  $A + A^*$  anticommutes with  $J$ ,
  - d)  $A + A^*$  is parallel, and nilpotent at every point,
  - e)  $\text{div } v = 0$ ,
  - f)  $v$  is a Killing field if and only if it is real-holomorphic.
- (6.1)

**Proof** Corollary 4.2 and (2.4.b) yield (6.1-a), while (6.1-a) trivially implies (6.1-b)–(6.1-c). Next, (2.9) and (2.4.b) prove the first part of (6.1-d). Thus,  $2 \text{div } v = 2 \text{tr } A = \text{tr } (A + A^*)$  is constant, and has zero integral, which gives (6.1.e) and the equality  $\text{tr } (A + A^*)^k = 0$  for  $k = 1$ . The same equality for  $k \geq 2$  now follows: setting  $C = (A + A^*)^{k-1}$ , we get  $\text{tr } (A + A^*)^k = \text{tr } (CA + CA^*) = 2 \text{tr } CA$ . (Note that  $\text{tr } CA^* = \text{tr } (CA^*)^* = \text{tr } AC = \text{tr } CA$ .) Due to the first part of (6.1-d),  $C$  is parallel and  $\text{tr } CA$  constant. As the constant  $\text{tr } CA$  equals  $C_k^i v^k_{,i} = (C_k^i v^k)_{,i} = \text{div } Cv$ , it has zero integral. The zero-integral constant  $\text{tr } (A + A^*)^k$  thus equals 0, which yields the induction step, proving the remainder of (6.1-d). Finally, (6.1-f) follows from Corollary 4.2: since  $\mathcal{L}_v \omega = 0$  and  $\omega = g(J \cdot, \cdot)$ , the condition  $\mathcal{L}_v g = 0$  is equivalent to  $\mathcal{L}_v J = 0$ . □

## 7 Two holomorphic covariant tensors

Equation (7.2) in Theorem 7.1 constitutes both a pseudo-Riemannian analog of a harmonic-coexact Hodge decomposition for the 1-form  $g(v, \cdot)$ , and the actual Riemannian Hodge decomposition of  $g(v, \cdot)$  relative to any Riemannian Kähler metric  $h$ , as long as one exists on  $M$ . This follows since  $(d\phi)J$  and  $\xi$  appearing in (7.2) are, respectively,  $h$ -coexact and  $h$ -harmonic, with the exact part absent, for every pseudo-Riemannian Kähler metric  $h$  which either equals our  $g$ , or is positive definite (and, in the case of  $g$ , by  $g$ -harmonic one means closed and  $g$ -coclosed). In fact,  $(d\phi)J$  is the  $h$ -divergence of  $-\phi h(J, \cdot)$ , for the  $h$ -Kähler form  $h(J, \cdot)$ . Furthermore, we prove below that both  $\xi$  and  $\xi J$  are closed, while, given any real 1-form  $\xi$  on a complex manifold,

$$\xi - i\xi J \text{ is holomorphic whenever } d\xi = d(\xi J) = 0, \tag{7.1}$$

since, in general, for a  $(p, 0)$ -form  $\zeta$ , closedness ( $d\zeta = 0$ ) obviously yields holomorphicity ( $\bar{\partial}\zeta = 0$ ). Being holomorphic,  $\xi - i\xi J$  must be  $h$ -harmonic [1, Ch. 5] when  $h$  is Riemannian. Finally, (7.2) and (6.1-e) imply  $g$ -coclosedness of  $\xi$ .

**Theorem 7.1** *Given an affine vector field  $v$  on a compact pseudo-Kähler  $\partial\bar{\partial}$  manifold  $(M, g)$ , one has*

$$g(v, \cdot) = (d\phi)J + \xi \tag{7.2}$$

for some  $\phi : M \rightarrow \mathbb{R}$  and the real part  $\xi$  of a holomorphic 1-form  $\xi - i\xi J$ .

**Proof** The assertions (2.5), (6.1.b), and (2.6) applied to  $B = A - A^*$ , show that  $d[g(v, \cdot)]$  is an exact  $(1, 1)$ -form. Choosing  $\phi$  as in (3.2) for  $\alpha = -d[g(v, \cdot)]/2$ , we now see that, by (3.3), the 1-form  $\xi = g(v, \cdot) - (d\phi)J$  is closed. However,  $\xi J = d\phi - g(Jv, \cdot) = d\phi - \omega(v, \cdot)$  is also closed, due to (2.8) and Corollary 4.2. The holomorphicity of  $\xi - i\xi J$  now follows from (7.1). □

Theorem 7.1 does not seem to be relevant to the question stated in the Introduction: for instance, vanishing of the holomorphic 1-form  $\xi - i\xi J$  (which follows if  $M$  is simply connected) does not lead to any immediate answer. This stands in marked contrast with the next result.

**Theorem 7.2** *Suppose that  $v$  is an affine vector field on a compact pseudo-Kähler  $\partial\bar{\partial}$  manifold  $(M, g)$ . Then the  $(0, 2)$  tensor field  $\mathcal{L}_v g$  is the real part of a holomorphic section  $\theta$  of the second complex symmetric power of the complex dual of  $TM$ , with the convention (2.7).*

Consequently,  $\mathcal{L}_v g = 0$  if no such nonzero holomorphic section  $\theta$  exists.

**Proof** By (2.4.a) and (6.1-c),  $[\mathcal{L}_v g](J \cdot, J \cdot) = -\mathcal{L}_v g$ , so that  $\theta = \mathcal{L}_v g - i[\mathcal{L}_v g](J \cdot, \cdot)$  is complex-bilinear at every point. As it is parallel—by (2.9)—its holomorphicity follows: Remark 2.2 easily implies that, for any (local) real holomorphic vector fields  $w, w'$ , the function  $\theta(w, w')$  is holomorphic.  $\square$

A complex manifold  $M$  is called *almost homogeneous* [7] if, for some  $x \in M$ , every vector in  $T_x M$  is the value at  $x$  of some real holomorphic vector field.

**Corollary 7.3** *If  $g$  is a pseudo-Kähler metric on a compact almost homogeneous complex  $\partial\bar{\partial}$  manifold  $M$  with nonzero Euler characteristic  $\chi(M)$ , then every  $g$ -affine vector field  $v$  on  $M$  is real holomorphic.*

In fact, let  $\theta$  be the holomorphic quadratic differential mentioned in Theorem 7.2. Thus,  $\theta(w, w')$  is constant for any real holomorphic vector fields  $w, w'$ . If  $\theta$  were nonzero—everywhere, due to its being parallel by (2.9)—choosing  $x$  as above and a real holomorphic vector field  $w$  with  $\theta(w, w) \neq 0$  at  $x$  we would get  $w \neq 0$  everywhere, and hence, by the Poincaré-Hopf theorem,  $\chi(M) = 0$ .

**Acknowledgements** The author wishes to thank Ivo Terek for very helpful discussions, and the anonymous referee for numerous comments and suggestions that greatly improved the exposition. Research supported in part by a FAPESP grant: 2015/50265-6.

## Declarations

**Conflict of interest** The author states that there is no conflict of interest.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Ballmann, W.: Lectures on Kähler Manifolds, ESI Lectures in Mathematics and Physics. European Mathematical Society, Zürich (2006)
2. Boubel, C., Mounoud, P.: Affine transformations and parallel lightlike vector fields on compact Lorentzian 3-manifolds. *Trans. Amer. Math. Soc.* **368**, 2223–2262 (2016)
3. Derdzinski, A., Terek, I.: Killing fields on compact pseudo-Kähler manifolds. *J. Geom. Anal.* **34**, 7, 144 (2024)
4. Deligne, P., Griffiths, P., Morgan, J., Sullivan, D.: Real homotopy theory of Kähler manifolds. *Invent. Math.* **29**, 245–274 (1975)

5. Kobayashi, S.: Transformation Groups in Differential Geometry. Springer-Verlag, Berlin-Heidelberg-New York (1972)
6. Lee, J. M.: Introduction to Smooth Manifolds. 2nd ed., Graduate Texts in Mathematics, vol. 218, Springer-Verlag, New York (2013)
7. Oeljeklaus, E.: Almost homogeneous manifolds. Inst. Élie Cartan, Université de Nancy, Institut Élie Cartan, Nancy **8**, 172–193 (1983)
8. Schouten, J. A., Yano, K.: On pseudo-Kählerian spaces admitting a continuous group of motions, Nederl. Akad. Wetensch. Proc. Ser. A **58**, Indag. Math. **17**, 565–570 (1955)
9. Voisin, C.: Hodge theory and complex algebraic geometry, I. In: Cambridge Studies in Advanced Mathematics, vol. 76. Cambridge University Press, Cambridge (2002)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.