

LECTURE 13

Operations with Linear Transformations

I. Sum of two linear transformations

II. composition of linear transformation

III. Inverse of an invertible

lin. transformation.

Operation with Linear Transformations

(3.1)

I. The Sum of Two Linear Transformations

For a moving viscous fluid passing

through a vectorial element of area

[A_i], the measured force vector [F_j]

acting on this area is expressed

quantitatively in terms of the stress

tensor (pressures, shear stresses)

[T_{ji}] which characterizes this fluid:

$$F_j = \sum_{i=1}^3 T_{ji} A_i.$$


If there are two kinds of fluid,

say type a and type b with stress

tensors [T_{ji}^a] and [T_{ji}^b], then the

force on this element of area is

$$F_j = \sum_{i=1}^3 (T_{ji}^a + T_{ji}^b) A_i$$



$$\vec{A}: A_1 = \frac{a_2 a_3}{2}$$

$$A_2 = \frac{a_3 a_1}{2}$$

$$A_3 = \frac{a_1 a_2}{2}$$

$$\left\{ \begin{array}{l} \vec{A} = \frac{1}{2} \vec{B} \times \vec{C} \\ = \frac{1}{2} \begin{vmatrix} i & j & k \\ a_1 & a_2 & 0 \\ a_1 & 0 & a_3 \end{vmatrix} \\ = i \underbrace{\frac{a_2 a_3}{2}}_{A_1} + j \underbrace{\frac{a_3 a_1}{2}}_{A_2} + k \underbrace{\frac{a_1 a_2}{2}}_{A_3} \end{array} \right.$$

Figure 13.1:
Components of an element of area \vec{A} as

projections of this area onto the respective coordinate planes.

$$(\vec{F})_j = \sum_i T_{ji} (\vec{A})_i$$

The force - area relation

The observation of such a relation between the concept "area" and the concept "force" is mathematized by linear transformations

$$T^a : U \rightarrow V$$

$$u_1 + u_2 \text{ and } T^a(u_1 + u_2) = T^a(u_1) + T^a(u_2)$$

$$T^b : U \rightarrow V$$

$$u_1 + u_2 \text{ and } T^b(u_1 + u_2) = T^b(u_1) + T^b(u_2)$$

and the definition of their sum:

Definition 13.1 (sum of two linear transformations)

The linear transformation

$$T^{a+b} = T^a + T^b$$

defined by the function

$$T^{a+b} : U \rightarrow V$$

$$u \text{ and } T^{a+b}(u) = T^a(u) + T^b(u)$$

is called the sum of T^a and T^b .

It is an easy exercise to validate
the following:

Theorem 13.1

If $T^a: U \rightarrow V$ and $T^b: U \rightarrow V$
are linear transformations, then so

is is their sum:

$$T^{a+b} = T^a + T^b : U \rightarrow V.$$

and their scalar multiple defined by

$$cT^a: U \rightarrow V$$

$$\text{such that } (cT^a)(u) = c(T(u)) \quad \forall u \in U$$

Comment:

At the risk of stating the obvious, note
that the c scalar multiple of T^a
depends on the validity of the c scalar
multiple of a vector being another vector.

II Composition of Two Transformations

1. Consider two rotation in \mathbb{R}^2

$$T^{\theta_1} = \begin{bmatrix} \cos\theta_1 & -\sin\theta_1 \\ \sin\theta_1 & \cos\theta_1 \end{bmatrix} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

$$T^{\theta_2} = \begin{bmatrix} \cos\theta_2 & -\sin\theta_2 \\ \sin\theta_2 & \cos\theta_2 \end{bmatrix} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

Then

$$T^{\theta_2} T^{\theta_1} = \begin{bmatrix} \cos\theta_2 \cos\theta_1 - \sin\theta_2 \sin\theta_1 & -(\cos\theta_2 \sin\theta_1 + \sin\theta_2 \cos\theta_1) \\ \sin\theta_2 \cos\theta_1 + \cos\theta_2 \sin\theta_1 & -\sin\theta_2 \sin\theta_1 + \cos\theta_2 \cos\theta_1 \end{bmatrix}$$

$$= \begin{bmatrix} \cos(\theta_2 + \theta_1) & -\sin(\theta_2 + \theta_1) \\ \sin(\theta_2 + \theta_1) & \cos(\theta_2 + \theta_1) \end{bmatrix}$$

$$= T^{\theta_2 + \theta_1} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

Conclusion: The composite of two rotations is another rotation.



2. Next consider the linear transformations.

$$T: \mathbb{R}^2 \rightarrow \mathbb{P}, \quad \text{and } S = P_1 \rightarrow P_2$$

$$\begin{bmatrix} a \\ b \end{bmatrix} \mapsto T\left(\begin{bmatrix} a \\ b \end{bmatrix}\right) = a + (a+b)x \quad \text{and } p(x) \mapsto S(p(x)) = xp(x)$$

This example and others like it lead to the concept of the composition of two linear transformations as follows

Definition 13.2 (The composition map)

If $T: U \rightarrow V$ and $S: V \rightarrow W$

are linear transformations, then the composition of S with T is the mapping defined by

$$S \circ T(u) = S(T(u))$$

where u is in U .

Observe that $S \circ T$ maps U to W :




Figure 13.2 $\xrightarrow{S \circ T} S \circ T(u) = S(T(u))$

Composite of two linear transformations

Returning to the example at the top of

page 13.6 with $u = \begin{bmatrix} a \\ b \end{bmatrix}$ one finds,

FIND $S \circ T\left(\begin{bmatrix} 5 \\ -4 \end{bmatrix}\right)$ and $S \circ T\left(\begin{bmatrix} a \\ b \end{bmatrix}\right)$.

SOLUTION

computing, one obtains

$$\begin{aligned} S \circ T\left(\begin{bmatrix} 5 \\ -4 \end{bmatrix}\right) &= S\left(T\begin{bmatrix} 5 \\ -4 \end{bmatrix}\right) = S(5 + (5-4)x) = S(5+x) \\ &= x(5+x) = 5x + x^2 \end{aligned}$$

and

$$\begin{aligned} S \circ T\left(\begin{bmatrix} a \\ b \end{bmatrix}\right) &= S\left(T\begin{bmatrix} a \\ b \end{bmatrix}\right) = S(a + (a+b)x) = ax + (a+b)x^2 \end{aligned}$$

As expected, one has the following

Theorem 13.2 (Composition preserves Linearity)

Given:

$T: U \rightarrow V$ is linear

$S: V \rightarrow W$ is linear.

Conclusion:

$S \circ T$ is linear.

Proof:

The validation of this conclusion hinges on the validity of the equation

$$\begin{aligned} S \circ T(c_1 u_1 + c_2 u_2) &= S(T(u_1) + c_2 T(u_2)) = \\ &= c_1 S(T(u_1)) + c_2 S(T(u_2)) = c_1 S \circ T(u_1) + c_2 S \circ T(u_2) \end{aligned}$$

$\forall u_1, u_2 \in U$ and \forall scalars c_1 and c_2 .

It is an easy exercise to verify this using the Definitions 13.2 and P13.1

on pages 13.6 and 11.3 respectively.

III. Invertible Linear Transformations.^{13.9}

There is a third way of obtaining a new transformation from a given linear transformation. This method consists of inverting a given linear transformation. The result of this process is the inverse of that transformation.

However, it is only invertible transformations which can be inverted. Such transformations are identified by means of the following

Definition 13.3 (Inverse of a transformation)

A linear transformation

$$T: U \rightarrow V$$

is said to be invertible if there exists
a transformation

$$T': V \rightarrow U$$

with the property that

$$\boxed{T' \circ T = I_U} \text{ and } \boxed{T \circ T' = I_V}$$

For example: $T' \circ T_{\text{PC}}(\text{poly}) = \text{poly}$ $T \circ T'((n+1)\text{-tuple}) = (n+1)\text{-tuple}$

where I_U and I_V are the identity

$$T: \text{poly's} \rightarrow n+1 \text{ tuples} \quad T'^{(n+1) \text{ tuples}} \rightarrow \text{poly's}$$

transformations on U and V respectively.

If there exists such a T' , one says that

T' is an inverse for T .

Comment

i. The domain space U and the target space V

do not have to be the same, as they do

in the case of invertible matrix
transformations.

2. If T' is an inverse for T , then
the definition implies that T is
an inverse for T' .

3. We shall see that T is invertible
whenever T is onto and one-to-one.

A. There are two ways of determining
whether T is onto (i.e. $R(T) = V$):

(i) Determine whether $\forall v \in V$ the
equation

$T(u) = v$
can be solved for $u \in U$.

(ii) Use the dimension argument to determine whether $R(T) = V$
 (i.e., is onto)

In, by the following 4-step argument:

1. Suppose V is finite dimensional,
 say

$$\dim V = p. \quad (*)$$

and

$$\dim R(T) = \dim V. \quad (**)$$

2. The fact that

$$R(T) \subseteq V \quad (*)$$

implies that $R(T)$ is a subspace, which

according to Eqs. $(**)$ and $(*)$ is

p -dimensional and hence has a basis

$$\{T(u_1), \dots, T(u_p)\} \equiv Q,$$

which is a linearly independent set.