

LECTURE 33

Diagonal Form of a Matrix.

Hamilton's Theorem

The Internal Input-Output Processing
Structure of a Matrix

Diagonalizable Matrices.

I. DIAGONAL FORM OF A MATRIX

Read Strang 5.2

The essential properties of a $n \times n$ matrix with a set of n linearly independent eigenvectors are expressed in terms of its eigenvalues and the transition matrix formed from these eigenvectors. This is expressed by the following

Proposition

Let $A: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be an $n \times n$ matrix

Let A have a set of n lin. indep. eigenvectors $\{\vec{e}_i\}$

Let S be the transition matrix formed from these eigenvalues.

Then

$$S^{-1}AS = \begin{bmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{bmatrix} \equiv \Lambda; \quad A = S\Lambda S^{-1}$$

One says that A can be diagonalized.

proof:

consider the equations

$$A\vec{e}_i = \lambda_i \vec{e}_i \quad i=1, \dots, n$$

satisfied by the given eigenvectors.

One has

$$A \begin{bmatrix} \vec{e}_1 \\ \vdots \\ \vec{e}_m \end{bmatrix} = \begin{bmatrix} \vec{e}_1 \lambda_1 & \dots & \vec{e}_n \lambda_n \end{bmatrix}$$

$$= \underbrace{\begin{bmatrix} \vec{e}_1 & \dots & \vec{e}_n \\ \vdots & & \vdots \end{bmatrix}}_S \underbrace{\begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix}}_\Lambda \leftarrow \Lambda !!$$

or

$$A S = S \Lambda \quad \text{where } S \text{ is invertible (why?)}$$

Thus $\boxed{S^{-1} A S = \Lambda}$ or $\boxed{A = S \Lambda S^{-1}}$ where $S^{-1} = \begin{bmatrix} \langle \omega^1 | \dots \end{bmatrix}$
 where $\omega^i(\vec{e}_j) = \langle \omega^i | \vec{e}_j \rangle = \delta_{ij}$

A is diagonalizable by means of a similarity transformation.

Comments:

1. It is easy to compute powers of a diagonalizable matrix:

$$A^m = (S \Lambda S^{-1}) \dots (S \Lambda S^{-1})$$

$$= S \Lambda \dots \Lambda S^{-1}$$

$$\boxed{A^m = S \Lambda^m S^{-1}}$$

2. One can compute matrix power series:
 consider a function defined by a convergent power series

$$g(x) = a_0 + a_1 x + a_2 x^2 + \dots \quad ; \text{ e.g. } e^x = 1 + x + \frac{x^2}{2} + \dots$$

Then $g(A)$ is defined and convergent in an appropriate sense:

$$\begin{aligned}
 e^{At} &= I + At + A^2 \frac{t^2}{2} + \dots \\
 &= S I S^{-1} + S A S^{-1} t + S A^2 S^{-1} \frac{t^2}{2!} + \dots \\
 &= S \left(I + \lambda t + \frac{\lambda^2 t^2}{2!} + \dots \right) S^{-1} \\
 &= S \begin{bmatrix} e^{\lambda_1 t} & & & \\ & e^{\lambda_2 t} & & \\ & & \ddots & \\ & & & e^{\lambda_n t} \end{bmatrix} S^{-1} \\
 &= S e^{\Lambda t} S^{-1}
 \end{aligned}$$

Thus $g(A) = S g(\Lambda) S^{-1}$

3. Consider the secular (a.k.a. characteristic) polynomial

$$\det |A - \lambda I| = a_0 + a_1(-\lambda) + \dots + a_n(-\lambda)^n$$

$$= (\lambda_1 - \lambda)(\lambda_2 - \lambda) \dots (\lambda_n - \lambda) \equiv f(\lambda)$$

$$= \lambda_1 \dots \lambda_n + (\dots)(-\lambda) + (\dots)(-\lambda)^2 + \dots + (\lambda_1 + \dots + \lambda_n)(-\lambda)^{n-1} + (-\lambda)^n = f(\lambda)$$

The secular pol' f in the powers of A vanishes!

Recalling that $\Lambda = \begin{bmatrix} \lambda_1 & & 0 \\ & \lambda_2 & \\ 0 & & \lambda_n \end{bmatrix}$, we have $f(\Lambda) = (I\Lambda_1 - \Lambda)(I\Lambda_2 - \Lambda) \dots (I\Lambda_n - \Lambda)$

$$f(A) = S^{-1} f(\Lambda) S =$$

$$= S^{-1} \begin{bmatrix} (\lambda_1 - \lambda_1)(\lambda_2 - \lambda_1) \dots (\lambda_n - \lambda_1) & & \\ & 0 & (\lambda_1 - \lambda_2)(\lambda_2 - \lambda_2) \dots (\lambda_n - \lambda_2) \\ & & \ddots & \\ & & & (\lambda_1 - \lambda_n)(\lambda_2 - \lambda_n) \dots (\lambda_n - \lambda_n) \end{bmatrix}$$

$$= \begin{bmatrix} \lambda_1 - \lambda_1 & & \\ \lambda_1 - \lambda_2 & & \\ & \ddots & \\ & & \lambda_n - \lambda_n \end{bmatrix} \begin{bmatrix} \lambda_1 - \lambda_1 & & \\ & \lambda_2 - \lambda_2 & \\ & & \lambda_n - \lambda_n \end{bmatrix} \\ = S \begin{bmatrix} 0 & & \\ & 0 & \\ & & 0 \end{bmatrix} S \\ = [0] \text{ matrix}$$

SKIP
obsolete

3. Let $A: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be diagonalizable

(i) $f(\lambda) = \det(A - \lambda I)$ is the characteristic poly of A .

$$= a_0 + a_1(-\lambda) + \dots + a_n(-\lambda)^n$$

$$= (\lambda_1 - \lambda)(\lambda_2 - \lambda) \dots (\lambda_n - \lambda)$$

(ii) $A = S^{-1} \Lambda S$; $\Lambda = \begin{bmatrix} \lambda_1 & & 0 \\ & \lambda_2 & \\ 0 & & \lambda_n \end{bmatrix}$

(iii) $f(\Lambda) = (\lambda_1 I - \Lambda)(\lambda_2 I - \Lambda) \dots (\lambda_n I - \Lambda)$

$$= \begin{bmatrix} \lambda_1 - \lambda_1 & & \\ & \lambda_1 - \lambda_2 & \\ & & \ddots \\ & & & \lambda_1 - \lambda_n \end{bmatrix} \begin{bmatrix} \lambda_2 - \lambda_1 & & \\ & \lambda_2 - \lambda_2 & \\ & & \ddots \\ & & & \lambda_2 - \lambda_n \end{bmatrix} \dots \begin{bmatrix} \lambda_n - \lambda_1 & & \\ & \lambda_n - \lambda_2 & \\ & & \ddots \\ & & & \lambda_n - \lambda_n \end{bmatrix}$$

$$= \begin{bmatrix} 0 & & \\ & 0 & \\ & & \ddots \\ & & & 0 \end{bmatrix}$$

(iv) $f(A) = S^{-1} f(\Lambda) S = \begin{bmatrix} 0 & & \\ & 0 & \\ & & \ddots \\ & & & 0 \end{bmatrix}$

Cayley-Hamilton's Theorem

A satisfies its characteristic eq'n, $f(A) = 0$.

$$f(A) = 0$$

We have

$$f(\lambda) = \lambda_1 \cdots \lambda_n + (\cdots)(-\lambda) + (\cdots)(-\lambda)^2 + \cdots + (\lambda_1 + \lambda_2 + \cdots + \lambda_n)(-\lambda)^{n-1} + (-\lambda)^n$$

$$f(A) = a_0 + a_1(A) + a_2(A)^2 + \cdots + a_{n-1}(A)^{n-1} + (-A)^n = \text{zero matrix}$$

This illustrates the Cayley-Hamilton theorem:

A matrix satisfies its characteristic equation.

(v) Here the coefficients

$$a_0 = f(0) = \lambda_1 \lambda_2 \cdots \lambda_n = \det A$$

$$a_1 =$$

$$a_{n-1} = \lambda_1 + \cdots + \lambda_n = \sum_{i=1}^n A_{ii} = \text{trace } A \quad (*)$$

⊙ \uparrow see below

depend only on the eigenvalues of A .

(v₂) In general $AB \neq BA$.

Question: True or False; $\text{tr } AB = \text{tr } BA$?

Answer: True, indeed one has

$$\text{tr } AB = A_{ij} B_{ji} = B_{ji} A_{ij} = \text{tr } BA$$

Application: For $S^{-1}AS = \Lambda$ one has

$$\begin{aligned} \lambda_1 + \cdots + \lambda_n &= \text{tr } \Lambda \\ &= \text{tr } S^{-1}AS \\ &= \text{tr } ASS^{-1} \\ &= \text{tr } A \end{aligned}$$


$$\text{tr } \Lambda = \text{tr } A$$

This validates Eq. (*).

4. A diagonalizable matrix A is a linear system whose internal structure is characterized by its eigenvalues

$$\lambda_1, \lambda_2, \dots, \lambda_n$$

Contact with its exterior is provided by its eigenvector basis and the corresponding dual basis. Diagrammatically one has



eigen vectors $\{\vec{e}_i\}$ $A = \sum_{i=1}^n \vec{e}_i \lambda_i \omega_i^i$ eigen covectors $\{\omega_i^i\}$

$\langle \omega_i^i | e_j \rangle = \delta_{ij}$
 $\langle \omega_i^i | \vec{v} \rangle = v^i$
 $A \vec{v} = \sum e_i \lambda_i v^i$

(a) This diagram conceptualizes the following physical process:

A matrix, say A , is a linear input-output device. Its output exhibits a causal relationship to the input. As depicted in the diagram, this relationship is the result of a three-stage process

1. The input is decomposed (by the eigen-covector basis $\{\omega_i^z\}$; see page 33.8) into its eigenvector components,
2. Each eigenvector component gets multiplied ("amplified", "phase-shifted", "filtered") by its respective eigenvalue λ_i .
3. The results of this modification get recombined by the eigenvector basis $\{e_i\}$. This recombination is the output from A .

(SKIP this in class, but hold student responsible for it) GOTO P33.9

(b) The mathematical (here algebraic) formulation of the diagram on page 33.6 is achieved by the following

4-step process:

33.7

STEP I.

Let $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ be the eigenvector basis corresponding to $\{\lambda_1, \lambda_2, \dots, \lambda_n\}$

Let $\{\omega^1, \omega^2, \dots, \omega^m\}$ be the corresponding cobasis for the dual space V^* (See Lecture 8).

The two bases are related by the duality relation

$$\langle \omega^i | \vec{e}_j \rangle = \delta^i_j.$$

Explicitly one has

$$\vec{e}_j = \begin{bmatrix} e_{j1} \\ e_{j2} \\ \vdots \\ e_{jn} \end{bmatrix}, \text{ the } j^{\text{th}} \text{ basis vector;}$$
$$A \vec{e}_j = \lambda_j \vec{e}_j$$

STEP II

Construct the cobasis elements

$$\omega^i = [\omega_1^i, \omega_2^i, \dots, \omega_n^i]$$

from the duality relation

$$\delta^i_j = \langle \omega^i | \vec{e}_j \rangle = [\omega_1^i, \dots, \omega_n^i] \begin{bmatrix} e_{j1} \\ \vdots \\ e_{jn} \end{bmatrix}$$

one solves for ω_i^z , the components of each ω^z :

$$\sum_{\ell=1}^n a_{\ell}^z e_{\ell}^z = \delta^z_j$$

Note that if the \vec{e}_j 's are the columns of

$$S = \begin{bmatrix} e_{11}^1 & e_{21}^1 & \dots & e_{n1}^1 \\ e_{12}^2 & e_{22}^2 & \dots & e_{n2}^2 \\ \vdots & \vdots & \ddots & \vdots \\ e_{1n}^n & e_{2n}^n & \dots & e_{nn}^n \end{bmatrix}$$

then the ω^z are the rows of

$$S^{-1} = \begin{bmatrix} \omega_1^1 & \omega_2^1 & \dots & \omega_n^1 \\ \vdots & \vdots & \ddots & \vdots \\ \omega_1^n & \omega_2^n & \dots & \omega_n^n \end{bmatrix} \begin{matrix} \leftarrow \omega^1 \\ \\ \leftarrow \omega^n \end{matrix}$$

STEP III

Exhibit the matrix

$$A = \sum_{z=1}^n \vec{e}_z \lambda_z \omega^z$$

STEP IV

Validate the correctness of this representation by applying A to the vector

$$\vec{x} = \sum_{j=1}^n e_j \langle \omega^j | \vec{x} \rangle = \sum_{j=1}^n \vec{e}_j x^j$$

we have

$$\begin{aligned}
 A(\vec{x}) &= \sum_i \vec{e}_i \lambda_i \omega_m^i(\vec{x}) \\
 &= \sum_i \vec{e}_i \lambda_i \langle \omega_m^i | \vec{e}_j x^j \rangle \\
 &= \sum_i \vec{e}_i \lambda_i \underbrace{\langle \omega_m^i | \vec{e}_j \rangle}_{\delta_j^i} x^j \\
 &= \sum_{i=1}^n \vec{e}_i \lambda_i x^i
 \end{aligned}$$

If \vec{x} is one of the eigenvectors, say

$$\vec{x} = \vec{e}_k,$$

then

$$A \vec{e}_k = \sum_i \vec{e}_i \lambda_i \underbrace{\langle \omega_m^i | \vec{e}_k \rangle}_{\delta_k^i}$$

$$\boxed{A \vec{e}_k = \vec{e}_k \lambda_k} \quad (\text{no sum over } k)$$

Thus

$$\boxed{A = \sum_{i=1}^n \vec{e}_i \lambda_i \omega_m^i}$$

mathematizes the diagram on page 33.5 indeed.

5.11) The eigenvectors $\{\vec{e}_i\}$,

$$A \vec{e}_i = \lambda_i \vec{e}_i \quad i=1, \dots, n$$

are right eigenvectors. By contrast

$\{\omega^k\}$ are left eigenvectors. Indeed,

$$\omega^k A = \langle \omega^k | \sum_i e_i \lambda_i \omega^i$$

$$= \sum_i \langle \omega^k | e_i \rangle \lambda_i \omega^i$$

$$\therefore \boxed{\omega^k A = \lambda_k \omega^k} \quad \text{Left eigenvector eq'n.}$$

Comment

Notice that the decomposition of A ,

$$A = \sum_{i=1}^n \vec{e}_i \lambda_i \omega_i'$$

and hence the eigenvalue eq'n

$$A \vec{x} = \lambda \vec{x}$$

did not in any way depend on the existence of an inner product.

The only necessary ingredient was

the construction of the dual eigen

basis $\{\omega_i'\}$ from the already known

eigenbasis $\{\vec{e}_i\}$ of A .

II. DIAGONALIZABLE MATRICES

33.12

It is not obvious that all matrices are diagonalizable. In fact it is not even true as exemplified by the ("defective") matrix $\begin{bmatrix} 1 & \\ 0 & 1 \end{bmatrix}$, whose characteristic polynomial $(1-\lambda)^2=0$ has degenerate roots.

However if the roots are distinct, then there exists a corresponding set of eigenvectors. This fact is made precise by the following

Theorem 33.1

A matrix with n distinct eigenvalues has a set of n eigenvectors which is linearly independent.

Because of this, one has the following

33, 13
35, 12

Corollary

A matrix with distinct eigenvalues can be diagonalized. (Because the transition matrix is non-singular)

Comment

1. The converse is not true: is false;

∃ many many diagonalizable matrices with degenerate eigenvalues.

For example, $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$.

2. The Theorem does not apply to "defective" matrices