

LECTURE 19

The Einstein Field Equations (E.F.E.)

- I) Einstein's tensorial line of reasoning
- II) Cartan's, Misner's and Wheeler geometrization of the E.F.E.

consigned
to lecture 20

- III) "Rotation" as a tensor.
- IV) Curvature as rotation.

I Einstein's line of reasoning that led to his gravitational field equations

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \frac{8\pi G}{c^2} T_{\mu\nu}$$

or equivalently

$$R_{\mu\nu} = \frac{8\pi G}{c^2} \left(T_{\mu\nu} - \frac{1}{2} g_{\mu\nu} T \right).$$

(2) Geometrize Newton's 1st Law relative to non-inertial reference frame.

$$\frac{d^2 x^\mu}{d\tau^2} = -\Gamma^\mu_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}$$

(i') Special Relativity:

Uniformly accelerated frame as a sequence of inertial frames.

(iii) Mathematize the dynamical laws governing particles and fields into coordinate frame independent form.

(iv) Recognize and incorporate the Equivalence

Principle as the metaphysical corner-

(Here "metaphysical" means: that which pertains to reality, to the nature of things, to existence.)

stone in conceptualizing gravitation:

a) "uniformly acc'd frame \equiv static, homogeneous gravitational field"

or

b) "inertial force = grav'l force"

(v) Apply the Equivalence Principle (E.P.) to the motion of bodies:

$$\frac{d^2 x^\mu}{d\tau^2} = -\Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau} \rightarrow \frac{d^2 x^{z'}}{d\tau^2} = -\Gamma_{00}^{z'}$$

$$= \frac{1}{2} g_{00,z'}$$

$$= -\phi_{,z'}$$

(E.P.)

(vi) Mathematize the momenergy properties and the dynamics of matter particles, and fields in geometrical form based on the momenergy tensor

$$\{T^{\mu\nu}\}:$$

$$T^{\mu\nu}_{;\nu} = 0.$$

(vii) Generalize the Newtonian gravitational field equation

$$\nabla^2 \phi = 4\pi G \rho$$

taking advantage of

a) the special relativistic mass-energy relation and

b) the fact that the Riemann curvature tensor

$$\{R^{\alpha}_{\beta\gamma\delta}\} = \{\Gamma^{\alpha}_{\beta\delta,\gamma} - \Gamma^{\alpha}_{\beta\gamma,\delta} + \Gamma^{\alpha}_{\sigma\gamma}\Gamma^{\sigma}_{\beta\delta} - \Gamma^{\alpha}_{\sigma\delta}\Gamma^{\sigma}_{\beta\gamma}\}$$

is the only tensor containing

2nd derivatives of $g_{\mu\nu}$, including

$$g_{00},{}^{;i}{}_{;i} = \left(-1 - \frac{2\phi}{c^2}\right),{}^{;i}{}_{;i} = -2 \nabla^2 \frac{\phi}{c^2},$$

imply that the tensorial generalization of the Newtonian gravitational field equation is

$$R_{\mu\nu} \equiv R^{\alpha}{}_{\mu\alpha\nu} = \text{expression in } T_{\mu\nu} \text{ and } g_{\mu\nu} \cdot T^{\alpha}{}_{\alpha}$$

(200) By demanding that momentum conservation

$$T^{\mu\nu}{}_{;\nu} = 0$$

be contained in a tensorial way of the tensorially generalized Newtonian equations

$$\underbrace{-\nabla^2(g_{00})}_{\left(-1 - 2\frac{\phi}{c^2}\right)} = \frac{8\pi G}{c^2} \rho,$$

i.e.,

$$\nabla^2 \frac{\phi}{c^2} = 4\pi G \frac{\rho}{c^2},$$

Einstein arrived at

$$R_{\mu\nu} = \frac{8\pi G}{c^2} \left(T_{\mu\nu} - \frac{1}{2} g_{\mu\nu} T^\alpha{}_\alpha \right)$$

which is equivalent to

$$\underbrace{R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R}_{\equiv G_{\mu\nu}} = \frac{8\pi G}{c^2} T_{\mu\nu}.$$

This equation incorporates momentum conservation

$$G_{\mu}{}^{\nu}{}_{;\nu} = 0$$

identically, and has the Newtonian grav'l equations as an asymptotic limit.

(v) Such a construction and line of reasoning is necessary, but not enough.

In physics and mathematics both sides, the l.h.s. and the r.h.s. of an equation (e.g. a stress-strain relation, $\vec{F} = m\vec{a}$, etc) must have a well-defined identity.

The r.h.s. of Einstein's equation, $T_{\mu\nu}$, is well-defined geometrically and physically.

However, this is not the case for the l.h.s.

(v2) In 1928 Cartan, and in 1964, 1972, 1990

Misner and Wheeler filled that cognitive

gap by restating Einstein's field equation ⁱⁿ geometrical form, both for the l.h.s. and the r.h.s.

$$L, h, s = \underbrace{\text{sum of moments of}}_{\text{curvature induced rotation for the 6 faces of a small 3-cube}} = r, h, s = \frac{8\pi G}{c^2} \underbrace{\text{amount of momenergy inside this 3-cube}}$$

A prerequisite for understanding and using the Einstein field equations is that one grasp the meaning and the geometrical formulation of the concepts

(i) "rotation" and (ii) "moment".

III)

ROTATION AS A TENSOR

19-8

(25.1)

The Physical Origin of Rotation,

In three dimensions consider a vector \vec{v} rotating with a given angular velocity around a given axis. The vectorial change $\Delta\vec{v}$ in this vector during time Δt is



$$\Delta\vec{v} = \Delta t \vec{\omega} \times \vec{v}$$

$$= \Delta t \begin{vmatrix} \vec{e}_1 & \vec{e}_2 & \vec{e}_3 \\ \omega^1 & \omega^2 & \omega^3 \\ v^1 & v^2 & v^3 \end{vmatrix}$$

Such a vectorial determinant can be generalized to higher dimensions. But, as far as I know, it will not represent a rotation in that case.

This is because the essential (= most consequential) property of the rotation process a plane in which the rotation

19-9,
(2.5.2)

takes place, not some normal to it. This

plane is spanned by a bivector as follows:

$$\begin{aligned}\Delta \vec{v} &= \Delta t \left[e_1 (\omega^2 \overset{(2b)}{v^3} - \omega^3 \overset{(2a)}{v^2}) - e_2 (\omega^1 \overset{(2b)}{v^3} - \omega^3 \overset{(2a)}{v^1}) + e_3 (\omega^1 \overset{(2b)}{v^2} - \omega^2 \overset{(2a)}{v^1}) \right] \\ &= -\Delta t \left[\omega^1 (e_2 \otimes e_3 - e_3 \otimes e_2) + \omega^2 (e_3 \otimes e_1 - e_1 \otimes e_3) + \omega^3 (e_1 \otimes e_2 - e_2 \otimes e_1) \right] \cdot \vec{v} \\ &= -\Delta t \left[\omega^1 e_2 \wedge e_3 + \omega^2 e_3 \wedge e_1 + \omega^3 e_1 \wedge e_2 \right] \cdot \vec{v}\end{aligned}$$

This change expresses an infinitesimal rotation.

It has components which express rotation

in each of three planes spanned by the

pairs of basis vectors in the ambient Euclidean inner product space.

In terms of the antisymmetric matrix

$$[R^{em}] = \begin{bmatrix} 0 & -\omega^3 & \omega^2 \\ \omega^3 & 0 & -\omega^1 \\ -\omega^2 & \omega^1 & 0 \end{bmatrix}$$

19-10
(25.3)

one has

$$\Delta \vec{v} = \frac{\Delta t}{2!} R^{lm} \vec{e}_l \wedge \vec{e}_m \cdot \vec{v} \quad (\text{Einstein Summation Convention})$$

$$\Delta \vec{v} = \Delta t R^{[lm]} \vec{e}_l \wedge \vec{e}_m \cdot \vec{v} \quad (\text{Summation restricted to } l < m)$$

By omitting reference to any particular vector \vec{v} one arrives at the concept of rotation as a tensor of rank $\begin{pmatrix} 2 \\ 0 \end{pmatrix}$.

Thus one has the following definition


Definition

1) A rotation is a second rank antisymmetric tensor

$$\frac{\Delta t}{2!} R^{lm} \vec{e}_l \wedge \vec{e}_m = \Delta t \vec{e}_l \wedge \vec{e}_m R^{[lm]}$$

IV) Curvature as Rotation

The concept of rotation defined this way generalizes to four (and higher) dimensions of spaces with an inner product (i.e. metric structure). Indeed, applying it to the curvature-induced rotational change associated with the u - v spanned face of a cube,



one has

$$\begin{aligned}
 \Delta W &= e_k w^k R^{\alpha\beta}{}_{k|} \omega^\alpha \omega^\beta(u, v) \\
 &= e_k w^k g_{km} R^{lm}{}_{|\alpha\beta|} \omega^\alpha \omega^\beta(u, v) \\
 &= e_k w^k \underbrace{e_i \cdot e_m}_{\delta_{im}} R^{lm}{}_{|\alpha\beta|}(u, v) \\
 &= e_k \otimes e_m \cdot \underbrace{W}_{\omega^\alpha \omega^\beta} R^{lm}{}_{|\alpha\beta|}(u, v)
 \end{aligned}$$

Taking advantage of the curvature's metric-induced antisymmetry,

$$R^{lm}{}_{\alpha\beta} = -R^{ml}{}_{\alpha\beta}, \text{ one has}$$

$$\begin{aligned} \Delta W &= \frac{1}{2} (e_l \otimes e_m - e_m \otimes e_l) \cdot W R^{lm}{}_{\alpha\beta} (u, v) \\ &= e_l \wedge e_m \cdot W R^{lm}{}_{\alpha\beta} (u, v) \end{aligned}$$

Comparing this with the rotation

defined on page 3, one arrives at

$$e_l \wedge e_m R^{lm}{}_{\alpha\beta} (u, v) = \text{"rotation"}$$

which is induced by the curvature in the area subtended by the vectors u and v .

This "rotation" is a $\begin{pmatrix} 2 \\ 0 \end{pmatrix}$ tensor. For infinitesimal vectors u and v its components $R^{lm}{}_{\alpha\beta} (u, v)$ are the angles by which vector such as w get

get rotated in the plane spanned by e_2 and e_m .

Notabene: In the context of spacetime the rotation can refer to Euclidean rotation, Lorentzian rotation or any of their combinations.