

Axioms for \mathbb{R}

The handout contains a list of axioms for the real numbers.

Observations:

- Most of these were also axioms for \mathbb{Z}

- There is no Well-Ordering axiom

- There are 2 new axioms.

⑦ Multiplicative Inverses: For each $a \in \mathbb{R}$ such that $a \neq 0$, there exists $a^{-1} \in \mathbb{R}$ such that

$$a \cdot a^{-1} = 1.$$

Write $\frac{b}{a}$ to mean $b \cdot a^{-1}$.

⑧ Least Upper Bound Property:
Every non-empty subset of \mathbb{R} which has an upper bound has a least upper bound in \mathbb{R} .

So • everything we proved about \mathbb{Z} without using Well-Ordering will also be true for \mathbb{R} .

• these new axioms will give \mathbb{R} new properties that we did not have in \mathbb{Z} .

Division and Rational Numbers

Lemma: For all $a, b \in \mathbb{R}$ with $a \neq 0$ and $b \neq 0$,

(a) If $a \cdot b = 1$, then $b = a^{-1}$. [Uniqueness of Mult. Inverses]

(b) $(a^{-1})^{-1} = a$.

(c) $(a \cdot b)^{-1} = a^{-1} \cdot b^{-1}$

(d) $(-a)^{-1} = -a^{-1}$

(e) $a > 0$ if and only if $a^{-1} > 0$.

In fraction notation: • $a \cdot b = 1 \Rightarrow b = \frac{1}{a}$

$$\bullet \frac{1}{\frac{1}{a}} = a$$

$$\bullet \frac{1}{ab} = \frac{1}{a} \cdot \frac{1}{b}$$

$$\bullet \frac{1}{(-a)} = -\frac{1}{a}$$

$$\bullet a > 0 \Leftrightarrow \frac{1}{a} > 0.$$

Proof: See handout.

Thm: Every integer is a real number.

Proof: The integers consist of positive numbers (\mathbb{N}), 0, and negative numbers ($-n$ for $n \in \mathbb{N}$).

$0 \in \mathbb{R}$ by Identity axiom. ✓

To show each $n \in \mathbb{N}$ is in \mathbb{R} , we use induction.

Base Case: $1 \in \mathbb{R}$ by Identity axiom.

Inductive Step: Let $n \in \mathbb{N}$ and suppose $n \in \mathbb{R}$. Then, since $1 \in \mathbb{R}$, we have $n+1 \in \mathbb{R}$. ✓

Lastly, since each $n \in \mathbb{N}$ is in \mathbb{R} , $-n$ will also be in \mathbb{R} by the Additive Inverses axiom. ■

Def: A real number $x \in \mathbb{R}$ is a rational number if there exist integers $a, b \in \mathbb{Z}$ such that $b \neq 0$ and $x = a \cdot b^{-1}$.

Write $x = \frac{a}{b}$, and say $\frac{a}{b}$ is a fraction representing x .

The set of all rational numbers is \mathbb{Q} .

Ex: $\frac{2}{3}$ and $\frac{8}{12}$ and $\frac{10}{15}$ are all different functions representing the same rational number.

Rule: $\frac{a}{b} = \frac{c}{d} \Leftrightarrow a \cdot b^{-1} = c \cdot d^{-1} \Leftrightarrow ad = bc$

"cross-multiply"

Lemma: For all $x, y \in \mathbb{Q}$,

a) $x + y \in \mathbb{Q}$

b) $x - y \in \mathbb{Q}$

c) $x \cdot y \in \mathbb{Q}$

d) if $y \neq 0$, then $x \cdot y^{-1} \in \mathbb{Q}$.

Proof: (a) Since x and y are rational, there exist integers $a, b, c, d \in \mathbb{Z}$ such that $b \neq 0$, $d \neq 0$, and

$$x = \frac{a}{b}, \quad y = \frac{c}{d}.$$

Then

$$x+y = \frac{a}{b} + \frac{c}{d} = a \cdot b^{-1} + c \cdot d^{-1}$$

So

$$\begin{aligned} (bd) \cdot (x+y) &= (bd)(ab^{-1} + cd^{-1}) \\ &= ad + bc. \end{aligned}$$

Thus,

$$\begin{aligned} x+y &= (ad+bc) \cdot (bd)^{-1} \\ &= \frac{ad+bc}{bd}. \end{aligned}$$

Now

- $ad+bc, bd \in \mathbb{Z}$

- $bd \neq 0$ because $b \neq 0$ and $d \neq 0$.

So $x+y = \frac{ad+bc}{bd} \in \mathbb{Q}$.

(b)-(d): HW 13



Lemma: Let $x \in \mathbb{Q}$. Then there is $m \in \mathbb{Z}$ and $n \in \mathbb{N}$ such that

$$x = \frac{m}{n}.$$

Proof: Since x is rational, there exist $a, b \in \mathbb{Z}$ such that

$$x = \frac{a}{b}.$$

- If $b > 0$, take $m = a$ and $n = b$.
- If $b < 0$, take $m = -a$ and $n = -b$, since

$$x = \frac{a}{b} = \frac{-a}{-b}.$$

Def: A fraction $\frac{a}{b}$ is in lowest terms if for every $d \in \mathbb{N}$, if $d|a$ and $d|b$, then $d=1$.

That is, 1 is the only positive divisor a and b have in common.

Ex: $\frac{2}{3}$ is in lowest terms. $\frac{8}{12}$ is not, because $4|8$ and $4|12$.

Def: Let $x \in \mathbb{Q}$. A possible positive denominator for x is a positive integer $n \in \mathbb{N}$ such that there exists $m \in \mathbb{Z}$ with $x = \frac{m}{n}$.

Ex: $\frac{2}{3} = \frac{4}{6} = \frac{8}{12} = \frac{20}{30} = \dots$

so 3, 6, 12, 30 are some of the possible denominators for this rational number.

Thm: Let $x \in \mathbb{Q}$. There exist $m \in \mathbb{Z}$ and $n \in \mathbb{N}$ such that $x = \frac{m}{n}$ and $\frac{m}{n}$ is in lowest terms.

Proof: Let S be the set of possible positive denominators for x .

By the lemma, x has a possible positive denominator, so S is a non-empty subset of \mathbb{N} .

By the Well-Ordering Principle, S has a smallest element. Call it n .

So $x = \frac{m}{n}$ for some $m \in \mathbb{Z}$.

Claim: $\frac{m}{n}$ is in lowest terms.

To prove this, assume it is not. Then there exists $d \in \mathbb{N}$ such that $d|m$ and $d|n$, and $d \neq 1$. So there exist $k, l \in \mathbb{Z}$ such that

$$m = dk \quad \text{and} \quad n = dl$$

Thus,

$$x = \frac{m}{n} = \frac{dk}{dl} = \frac{k}{l}.$$

Now, $\bullet l \in \mathbb{N}$ [because $n, d \in \mathbb{N}$]

$\bullet l < n$ [because $d > 1$]

Thus, l is a possible positive denominator for x which is smaller than n , a contradiction. \bullet