

Warm-Up: What is the difference between the following three sets?

- $\{0, 1\}$
 - $\{\{0, 1\}\}$
 - $\{\{0\}, \{1\}\}$
-

Other ways to specify sets

In words: • Let B be the set whose elements are the first five prime numbers

$$\text{so } B = \{2, 3, 5, 7, 11\}$$

• Let $\mathbb{R}_{>0}$ be the set of all positive real numbers.

By patterns: • $E = \{2, 4, 6, 8, \dots\}$

(E is the set of positive even numbers)

• $P = \{2, 3, 5, 7, 11, \dots\}$

(P is the set of all prime numbers)

These first two methods are somewhat limited.

Set-Builder Notation: If $P(x)$ is a sentence,
then

$$\{x \mid P(x)\} \quad \text{or} \quad \{x : P(x)\}$$

\uparrow "such that" \uparrow "such that"

is the set of all x such that $P(x)$
is true.

If A is a set, then

$$\{x \in A \mid P(x)\}$$

is the set of all x such that $x \in A$
and $P(x)$ is true. (x is a bound variable)

- $E = \{n \mid n \in \mathbb{N} \text{ and } n \text{ is even}\}.$

Also, $E = \{n \in \mathbb{N} \mid n \text{ is even}\}.$

Also, $E = \{n \in \mathbb{N} : 2 \mid n\}.$

- $\mathbb{R}_{>0} = \{x \in \mathbb{R} \mid x > 0\}.$

Also, $\mathbb{R}_{>0} = \{y \in \mathbb{R} \mid y > 0\}$

Transformation notation: If A is a set and f is some function defined on A , then

$$\{f(x) \mid x \in A\}$$

is the set of all objects $f(x)$ obtained from all $x \in A$.

$$\bullet E = \{2n \mid n \in \mathbb{N}\}$$

$$\bullet S = \{n^2 \mid n \in \mathbb{N}\} \leftarrow \text{Transformation}$$
$$= \{m \mid \text{there exists } n \in \mathbb{N} \text{ such that } n^2 = m\}$$

\leftarrow Set-Builder

Ex: Let $S = \{n^2 \mid n \in \mathbb{N}\}$ be the set of (positive) squares and $C = \{n^3 \mid n \in \mathbb{N}\}$ the set of (positive) cubes.

Define $A = \{x + y \mid x \in S \text{ and } y \in C\}$.

Then

In words: A is the set of integers which can be written as the sum of a positive square and a positive cube.

Pattern: $A = \{2, 5, 9, 10, 12, 17, 24, \dots\}$

Set-Builder:

$$A = \{n \in \mathbb{N} \mid \text{there exist } a, b \in \mathbb{N} \text{ such that } n = a^2 + b^3\}$$

Subsets

Def: Let A and B be sets. We say A is a subset of B , written $A \subseteq B$, if $x \in A$ implies $x \in B$.

That is, $A \subseteq B$ means every element of A is also an element of B .

Can write as $(\forall x \in A)(x \in B)$
or $(\forall x)[x \in A \Rightarrow x \in B]$

- Ex:
- $\{2, 3, 5\} \subseteq \{1, 2, 3, 4, 5\}$
 - $\mathbb{N} \subseteq \mathbb{Z}$
 - $\mathbb{Z} \subseteq \mathbb{Q}$
 - $\mathbb{Q} \subseteq \mathbb{R}$

Ex: Any time we use set-builder notation to write

$$A = \{x \in B \mid P(x)\},$$

we have $A \subseteq B$.

Thm: For every set A , $\emptyset \in A$.

Proof: The sentence $x \in \emptyset$ is always false.

Thus,

$$x \in \emptyset \Rightarrow x \in A$$

is always true, so $\emptyset \in A$. \blacksquare

Thm: For every set A , $A \subseteq A$.

Proof: The sentence

$$x \in A \Rightarrow x \in A$$

is true for all x , so $A \subseteq A$. \blacksquare

Def: Let A and B be sets. We say $A = B$ if $A \subseteq B$ and $B \subseteq A$.

So to prove $A = B$, we usually have to prove 2 things:

• $A \subseteq B$ ($x \in A \Rightarrow x \in B$)

• $B \subseteq A$ ($x \in B \Rightarrow x \in A$)

Thm: Let A and B be sets. Then $A = B$ if and only if $(x \in A \Leftrightarrow x \in B)$ for all x .

Proof: $A = B$ is logically equivalent to

$$(A \subseteq B) \wedge (B \subseteq A)$$

$$\equiv (\forall x)(x \in A \Rightarrow x \in B) \wedge (\forall x)(x \in B \Rightarrow x \in A)$$

$$\equiv (\forall x) [(x \in A \Rightarrow x \in B) \wedge (x \in B \Rightarrow x \in A)]$$

$$\equiv (\forall x) [x \in A \Leftrightarrow x \in B].$$

since $(\forall x) P(x) \wedge (\forall x) Q(x) \equiv (\forall x) [P(x) \wedge Q(x)]$

Ex: Let's prove

$$\underbrace{\{x \in \mathbb{Z} \mid x^2 = 1\}}_{=A} = \underbrace{\{1, -1\}}_{=B}$$

We must prove $A \subseteq B$ ($x \in A \Rightarrow x \in B$)
and $B \subseteq A$ ($x \in B \Rightarrow x \in A$).

Proof: (\subseteq) Let $x \in A$. Then $x \in \mathbb{Z}$ and $x^2 = 1$. So

$$x^2 - 1 = 0$$

$$(x-1)(x+1) = 0.$$

Thus, $x = 1$ or $x = -1$, so $x \in B$.

(\supseteq) Let $x \in B$. Then $x = 1$ or $x = -1$.
Either way, $x \in \mathbb{Z}$ and $x^2 = 1$,
so $x \in A$. \square