

HOMEWORK 16
MATH 3345 – SPRING 2023 – KUTLER

Exercises

Please complete the following problems on your own paper. Solutions should be written clearly, legibly, and with appropriate style.

1. Let $a, b \in \mathbb{N}$ and set $d = \gcd(a, b)$.
 - (a) Explain why $\frac{a}{d}$ and $\frac{b}{d}$ are integers.
 - (b) Prove that $\gcd\left(\frac{a}{d}, \frac{b}{d}\right) = 1$.
[HINT: Consider the prime factorizations of a and b . What does the prime factorization of d look like?]

2. **[Falkner Section 4 Exercise 8]** Let x and y be rational numbers. Prove the following statements.
 - (a) $-y$ is a rational number.
 - (b) $x - y$ is a rational number.
 - (c) xy is a rational number.
 - (d) If $y \neq 0$, then $1/y$ is a rational number.
 - (e) If $y \neq 0$, then x/y is a rational number.

3. **[Falkner Section 4 Exercise 10 – modified]** Let x be a rational number and let y be an irrational number. Prove the following statements.
 - (a) $-y$ is irrational.
 - (b) $x - y$ is irrational.
 - (c) $y - x$ is irrational.
 - (d) If $x \neq 0$, then xy is irrational. [Be sure to explain where you use the condition that $x \neq 0$ in your proof.]
 - (e) Explain why the condition that $x \neq 0$ was necessary for part (d). That is explain why xy is rational when $x = 0$.
 - (f) $1/y$ is irrational. [You should explain why $y \neq 0$ **must** be true.]
 - (g) If $x \neq 0$, then x/y is irrational.
 - (h) If $x \neq 0$, then y/x is irrational.

Practice Problems

It is strongly recommended that you complete the following problems. There is no need to write up polished, final versions of your solutions (although you may find this a useful exercise). Please do not submit any work for these problems.

1. **[Falkner Section 4 Exercise 20 (The rational roots theorem)]** Let $r \in \mathbb{Q}$ be a rational number such that

$$c_n r^n + c_{n-1} r^{n-1} + \cdots + c_1 r + c_0 = 0,$$

where $n \in \mathbb{N}$ and $c_0, c_1, \dots, c_n, c_{n-1} \in \mathbb{Z}$, and $c_n \neq 0$. In other words, r is a rational root of the polynomial $f(x) = c_n x^n + c_{n-1} x^{n-1} + \cdots + c_1 x + c_0$, and every coefficient of this polynomial is an integer.

Prove that r can be written in the form $r = \frac{a}{b}$, where a is an integer such that $a|c_0$ and b is a nonzero integer such that $b|c_n$.

[HINT: Write $r = \frac{a}{b}$ as a fraction in lowest terms. Then $\gcd(a, b) = 1$. (Why?) What does this say about the prime factorizations of a and b ?]

2. **[Falkner Section 4 Exercise 21]** Let $f(x) = 3x^3 - 40x^2 + 97x + 10$.

- (a) Find a rational number $r \in \mathbb{Q}$ such that $f(r) = 0$. [HINT: Use the rational roots theorem to narrow down the possibilities for r .]
- (b) Find two other real numbers s and t such that $f(s) = 0$ and $f(t) = 0$. [HINT: Use part (a) and polynomial long division to write $f(x) = (x - r)g(x)$, where $g(x)$ is a quadratic polynomial.]
- (c) Explain why s and t must be irrational. [HINT: There are several ways to do this. One elegant way is to notice that $g(x) = 3h(x)$, where $h(x)$ is a quadratic polynomial to which it is particularly easy to apply the rational roots theorem.]