

Recall: A set A is finite if

- $A = \emptyset$

or

- For some $n \in \mathbb{N}$, there is a bijection

$$f: \{1, 2, \dots, n\} \rightarrow A.$$

Think: f lists all elements of A on n lines with no repeats.

In this case, write $|A| = n$.

Warm-Up: You probably intuit that \mathbb{N} is infinite (i.e., not finite).

Try to prove this.

Hint: Use contradiction.

Here's one solution: Suppose $|N| = n$, so

there is a bijection $f: \{1, 2, \dots, n\} \rightarrow N$.

Let

$M = \text{maximum of } f(1), f(2), \dots, f(n).$

Then $f(k) \leq M < M+1$ for all $k \in \{1, \dots, n\}$, so
 $M+1 \in N \setminus \text{Rng}(f).$

Thus, f is not surjective, so cannot be a bijection. 

Ex: Similarly, \mathbb{Q} and \mathbb{R} are infinite.

WARNING: It may be tempting to write

$$|N| = \infty$$

$$|\mathbb{Q}| = \infty$$

$$|\mathbb{R}| = \infty$$

We will not do this.

As we will see, $|N| = |\mathbb{Q}|$, but $|N| \neq |\mathbb{R}|$.

First, more on finite sets.

Thm: Let S be a finite set and $T \subseteq S$.
Then

- T is finite
- $|T| \leq |S|$
- $|T| = |S|$ if and only if $T = S$.

Proof: Book Thms 13.30, 13.33.

(By induction on $|S|$. Not hard - just tedious.)

Cor: Let A and B be finite sets, and let $f: A \rightarrow B$ be a function. Then

- ① If f is an injection, then $|A| \leq |B|$
- ② If f is a surjection, then $|A| \geq |B|$

Proof: ① Suppose $f: A \rightarrow B$ is injective. Then

$$f: A \rightarrow \text{Rng}(f)$$

is a bijection. Hence, $|A| = |\text{Rng}(f)|$.

But $\text{Rng}(f) \subseteq B$, so $|\text{Rng}(f)| \leq |B|$ by the Theorem. Together, we get $|A| \leq |B|$.

② Suppose $f: A \rightarrow B$ is surjective. Since B is finite, $|B| = n$ for some $n \in \mathbb{N}$, so we can write

$$B = \{b_1, b_2, \dots, b_n\}.$$

For each $i \in \{1, \dots, n\}$, let $a_i \in A$ be such that $f(a_i) = b_i$.

If $i \neq j$, then $f(a_i) = b_i \neq b_j = f(a_j)$, so $a_i \neq a_j$.

Thus, $|\{a_1, \dots, a_n\}| = n$. But $\{a_1, \dots, a_n\} \subseteq A$, so $n \leq |A|$. Since $|B| = n$, we have $|A| \geq |B|$.



The contrapositive of ① is the

Pigeonhole Principle: Let A and B be finite sets and $f: A \rightarrow B$ a function.

If $|A| > |B|$, then f is not injective.

A - set of pigeons

B - set of pigeonholes

$f: A \rightarrow B$ puts each pigeon in a pigeonhole

Then there is a pigeonhole containing more than one pigeon.

Ex: If $a_1, a_2, a_3, a_4 \in \mathbb{Z}$, then the difference $a_i - a_j$ will be divisible by 3 for some $i \neq j$.

Ex: Suppose n people are at a party. Then there are two people who have the same number of friends at the party.

↳ Cannot be someone with 0 friends and someone with $n-1$ friends. So possibilities are $0, \dots, n-2$ or $1, \dots, n-1$.