

FINITENESS OF MAXIMAL GEODESIC SUBMANIFOLDS IN HYPERBOLIC HYBRIDS

DAVID FISHER, JEAN-FRANÇOIS LAFONT, NICHOLAS MILLER,
AND MATTHEW STOVER

ABSTRACT. We show that large classes of non-arithmetic hyperbolic n -manifolds, including the hybrids introduced by Gromov and Piatetski-Shapiro and many of their generalizations, have only finitely many finite-volume immersed totally geodesic hypersurfaces. In higher codimension, we prove finiteness for geodesic submanifolds of dimension at least 2 that are *maximal*, i.e., not properly contained in a proper geodesic submanifold of the ambient n -manifold. The proof is a mix of structure theory for arithmetic groups, dynamics, and geometry in negative curvature.

1. INTRODUCTION

To simplify the discussion, throughout this introduction a hyperbolic manifold will mean a connected, oriented, complete, finite-volume hyperbolic n -manifold, and a geodesic submanifold will be a complete, immersed, finite-volume, totally geodesic subspace (note that a totally geodesic subspace need not be orientable). The main motivation of this paper is the following question which, as far as we know, has been asked independently by Reid and in dimension 3 by McMullen [13, Qn. 7.6]:

Question 1.1. *Let M be a non-arithmetic hyperbolic manifold of dimension at least 3. Does M have at most finitely many geodesic submanifolds of codimension one?*

We will prove the answer is yes for a large class of non-arithmetic hyperbolic manifolds that includes the famous non-arithmetic manifolds constructed by Gromov and Piatetski-Shapiro [18]. We also prove more general results valid in arbitrary codimension $k < n - 1$, answering a more general question of Reid. Making a precise statement requires some more care and notation. Before embarking on that, we discuss some background and motivation for the conjecture.

Totally geodesic submanifolds of hyperbolic manifolds, when they exist, have proven fundamental in solving a number of important problems. Perhaps most famously, Gromov and Piatetski-Shapiro used cut-and-paste of arithmetic hyperbolic manifolds along codimension one geodesic submanifolds to build non-arithmetic hyperbolic manifolds in all dimensions [18]. More recently, variants on their construction first introduced in [1] were used by Geland and Levit to prove that “most” hyperbolic manifolds in dimension at least 4 are non-arithmetic [16].

In another famous application, Millson used geodesic submanifolds to construct hyperbolic manifolds in all dimensions with positive 1st betti number [28]. Millson’s examples are arithmetic, and a key idea in the proof is that if an arithmetic hyperbolic manifold contains a geodesic submanifold, then it contains many of them. One can see this as an easy application of the fact that the commensurator of an

arithmetic group is dense in the isometry group of the associated symmetric space [7, Th. 2]. In particular, density of the commensurator allows one to prove the following:

Arithmetic geodesic submanifold dichotomy: *For any $1 \leq k \leq n - 1$, an arithmetic hyperbolic n -manifold either contains no codimension k geodesic submanifolds, or it contains infinitely many and they are everywhere dense.*

This was perhaps first made precise in dimension 3 by Maclachlan–Reid and Reid [20, 34], who exhibited the first hyperbolic 3-manifolds with no totally geodesic surfaces. (For non-arithmetic examples, see [8] for fibered knots with no totally geodesic surfaces.)

In the non-arithmetic setting, this commensurator argument is not available. A theorem of Margulis shows that an irreducible lattice Γ in an adjoint semisimple Lie group G is arithmetic if its commensurator is topologically dense in G . From this one deduces that if Γ is non-arithmetic, then its commensurator is itself a lattice that contains Γ with finite index [24, p. 2], and so Question 1.1 is reasonable. The simplest form of our main result is the following.

Theorem 1.2. *For every $n \geq 3$, there exist infinitely many commensurability classes of finite-volume non-arithmetic hyperbolic n -manifolds for which the collection of all codimension one finite-volume totally geodesic submanifolds is finite but nonempty.*

As we will describe below, Theorem 1.2 applies in particular to the class of manifolds constructed by Gromov and Piatetski-Shapiro, along with many variants of their construction. In dimension $n \geq 4$, we do not know an example of a non-arithmetic hyperbolic n -manifold containing no codimension one geodesic submanifolds. Interestingly, finiteness results of this kind were only known previously for compact immersed totally geodesic surfaces in *infinite volume* hyperbolic 3-manifolds with compact convex core, by McMullen–Mohammadi–Oh [25, Thm. 1.4]. See very recent work of Benoist–Oh for the geometrically finite case [6, Thm. 1.5].

Even for finite-volume hyperbolic 3-manifolds, Theorem 1.2 is new and the first result of its kind on the question of McMullen and Reid. All previous arguments in that setting could only show finiteness of geodesic subsurfaces by showing there were none at all. In fact, we will show in §6.1 that explicit examples of hyperbolic links with a finite but nonempty set of totally geodesic surfaces are relatively easy to construct.

Theorem 1.3. *There are infinitely many distinct hyperbolic links L in the 3-sphere so that the collection of all finite-area totally geodesic surfaces in $S^3 \setminus L$ is finite but nonempty. One can choose L such that all its totally geodesic surfaces are noncompact.*

In short, our examples are *belted sums* of arithmetic links. See §6 for more on various examples and nonexamples, along with connections between our work and the Menasco–Reid conjecture [26].

As mentioned briefly above, Theorem 1.2 is a special case of a more general theorem on geodesic submanifolds of arbitrary codimension. To state this result we require some more notation. A *building block* of a hyperbolic n -manifold M

is a connected n -dimensional submanifold N with nonempty (but possibly disconnected) totally geodesic boundary such that $\pi_1(N) < \pi_1(M)$ is Zariski dense in $\text{Isom}(\mathbb{H}^n)$. In the finite-volume noncompact case, we require all cusps of N to be finite-volume.

We say a hyperbolic manifold M is *built from building blocks* when it is the union of finitely many building blocks with disjoint interiors, hence they intersect only along their boundaries. That is, M is obtained by gluing finitely many building blocks together isometrically along their boundaries. We call two such blocks *adjacent* if they meet in M along a common boundary component. We will frequently refer to any connected component of this common boundary as a *cutting hypersurface*.

We say that a building block $N \subset M$ is *arithmetic* when there exists an arithmetic hyperbolic n -manifold M' such that N is isometric to a building block for M' . We will see that, up to commensurability, M' is uniquely determined by N . In particular, it makes sense to say that two arithmetic building blocks are *similar* when they are building blocks for commensurable arithmetic manifolds and are *dissimilar* otherwise. We will also call a geodesic submanifold *maximal* if it is not contained in a proper geodesic submanifold of smaller codimension.

Our main result is:

Theorem 1.4. *Let M be a non-arithmetic hyperbolic n -manifold built from building blocks for which there are two adjacent building blocks N_1 and N_2 that are arithmetic and dissimilar. For each $1 \leq k \leq n - 2$, the collection of all maximal codimension k finite-volume totally geodesic submanifolds of M is finite.*

Note that the word maximal in the theorem is necessary, since maximal codimension k finite volume totally geodesic submanifolds may be arithmetic and thus contain infinitely many finite volume totally geodesic submanifolds of all greater codimensions. This occurs quite frequently in explicit constructions, particularly those built by Gromov and Piatetski-Shapiro.

Also note that the manifolds constructed by Gromov and Piatetski-Shapiro are built from two dissimilar building blocks (see §2.2), so Theorem 1.4 applies. The theorem also applies to the manifolds used to study invariant random subgroups in [1] and to those used by Raimbault [30] and Gelander–Levit [16] in studying growth of the number of maximal lattices in $\text{SO}(n, 1)$. These lattices are all built from *subarithmetic pieces* in the language of Gromov and Piatetski-Shapiro [18, Qn. 0.4]. We state one immediate corollary of our result and the results of Gelander–Levit more carefully:

Theorem 1.5. *Fix $n > 3$. Then there exists a number $b_n > 0$ such that the number of commensurability classes of hyperbolic manifolds satisfying the conclusions of Theorem 1.4 and having a representative of volume less than V is proportional to $V^{b_n V}$.*

As in Gelander–Levit, it is possible to see from this that there are “more” manifolds satisfying the conclusions of Theorem 1.4 than there are arithmetic manifolds. In fact, the set of such manifolds has positive logarithmic density if one counts by minimal volume in the commensurability class.

We note here that our proofs do not apply to the class of hyperbolic manifolds built by inbreeding that were introduced by Agol [3] and developed further by Belolipetsky and Thomson [5]. Our approach depends at a key point on the adjacent pieces being associated with different arithmetic groups, and in this family of examples there is only one arithmetic group involved. In other words, in our terminology the building blocks used to construct the examples in [3] and [5] are similar, hence Theorem 1.4 does not apply.

The proof of Theorem 1.4 is a mix of structure theory for arithmetic groups, dynamics, and geometry in negative curvature.

Our proof involves three key ingredients. The first two are a pair of ‘rigidity’ theorems for geodesic submanifolds of manifolds M satisfying the properties of Theorem 1.4. Each shows that geodesic submanifolds of M that meet both N_1 and N_2 must be of a very particular kind. The third ingredient consists of equidistribution results from homogeneous dynamics.

The first ingredient is ‘Closure Rigidity’, which we prove in §3. Given a geodesic submanifold Σ that crosses both N_1 and N_2 , we show that $\Sigma \cap N_i$ is actually the intersection with N_i of a geodesic submanifold in the finite volume arithmetic manifold from which N_i is cut. This is not true for geodesics, and in fact a generic geodesic in the examples we consider is built from pieces that do not close in the corresponding arithmetic manifold.

The second ingredient and rigidity result is ‘Angle Rigidity’, proven in §4. This says that, once we know the closure rigidity results from §3, geodesic submanifolds must meet $N_1 \cap N_2$ orthogonally.

The last key ingredient in the proof is a generalization of Shah’s work on orbit closures in hyperbolic manifolds [37], which we prove in §5. This allows us to conclude that either the collection of codimension k geodesic submanifolds of M satisfies the conclusions of Theorem 1.4, or they determine a dense subset of the oriented orthonormal frame bundle of M . The combination of closure and angle rigidity forbids the second conclusion, hence the theorem follows.

A special case of our angle rigidity result was recently and independently proven by Benoist–Oh and used to study related questions in the case of geodesic surfaces in geometrically finite hyperbolic 3 manifolds [6, Section 12.5]. After we showed our paper to Oh, she pointed out that in dimension 3 one can use their case of that result and their Theorem 11.8 in place of our Theorem 3.1 to prove our Theorem 1.2.

We close with some discussion of the following famous question asked by Gromov and Piatetski-Shapiro.

Question 1.6 (Qn. 0.4 [18]). *Call a discrete subgroup $\Gamma_0 < \mathrm{PO}(n, 1)$ subarithmetic if Γ_0 is Zariski dense and if there exists an arithmetic subgroup $\Gamma_1 < \mathrm{PO}(n, 1)$ such that $\Gamma_0 \cap \Gamma_1$ has finite index in Γ_0 . Does every lattice Γ in $\mathrm{PO}(n, 1)$ (maybe for large n) contain a subarithmetic subgroup? Is Γ generated by (finitely many) such subgroups? If so, does $V = \mathbb{H}^n/\Gamma$ admit a “nice” partition into “subarithmetic pieces”.*

Manifolds built from arithmetic building blocks certainly admit a nice partition into subarithmetic pieces. In a different direction, in the 1960s Vinberg introduced the notion of a *quasi-arithmetic* lattice [39] (see §6.2). For example, many lattices acting on \mathbb{H}^n generated by reflections are quasi-arithmetic. However, the typical

variants of Gromov and Piatetski-Shapiro’s construction of non-arithmetic lattices are not quasi-arithmetic, so Question 1.6 is well-known to be more subtle than “are all non-arithmetic lattices iterations of the construction in [18]”.

A key step in the proof that the examples in [18] are non-arithmetic is the assumption that one interbreeds noncommensurable arithmetic manifolds. More recently, Agol [3] (for $n = 4$) and Belolipetsky–Thomson [5] (for $n \geq 4$) constructed non-arithmetic but quasi-arithmetic lattices via inbreeding commensurable arithmetic lattices. In particular, the existence of quasi-arithmetic lattices that are non-arithmetic does not rule out the possibility that all non-arithmetic lattices for $n \geq 4$ are constructed by inbreeding arithmetic lattices (there are hyperbolic 3-manifolds with no totally geodesic surfaces, which clearly are not constructed via interbreeding or inbreeding; see Example 6.3).

With the methods of this paper, we can give examples of Coxeter polyhedra whose associated non-arithmetic lattices are not commensurable with either the Gromov–Piatetski-Shapiro type lattices, or to the Agol type lattices. See §6.2, where we will prove the following, which perhaps further clarifies the subtle nature of Question 1.6.

Theorem 1.7. *There exist non-arithmetic lattices in $SO(5, 1)$ that are not commensurable with a lattice constructed by the methods of Gromov–Piatetski-Shapiro or Agol.*

Acknowledgments. Fisher was partially supported by NSF Grants DMS-1308291 and DMS-1607041. Lafont was partially supported by the NSF grant number DMS 1510640. Stover was partially supported by the NSF Grant Number DMS 1361000 and Grant Number 523197 from the Simons Foundation/SFARI. The authors acknowledge support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: GEometric structures And Representation varieties” (the GEAR Network).

2. BACKGROUND ON CONSTRUCTIONS OF HYPERBOLIC n -MANIFOLDS

This section gives the necessary background on the constructions of hyperbolic n -manifolds that are relevant for this paper. It also establishes much of the notation that we will use throughout.

2.1. Arithmetic manifolds constructed from quadratic forms. Here we recall the construction of arithmetic hyperbolic manifolds via quadratic forms over number fields. Our standing assumption throughout will be that K is a totally real number field of degree s over \mathbb{Q} with distinct real embeddings $\sigma_1, \dots, \sigma_s : K \rightarrow \mathbb{R}$. To simplify our discussion, we assume that K is a subfield of \mathbb{R} via $\sigma_1 = \text{Id}$.

Now, let V be a K -vector space of dimension $n + 1$ and q a nondegenerate K -defined quadratic form on V . Such a pair (V, q) is called a K -quadratic space, and associated with q we have the inner product

$$(1) \quad \langle u, v \rangle_q = \frac{1}{2} (q(u + v) - q(u) - q(v)).$$

We then have the K -algebraic group $SO(q, K)$ consisting of those determinant one linear transformations of V that preserve q . We use

$$V(\mathbb{R}) = V \otimes_K \mathbb{R} \cong \mathbb{R}^{n+1},$$

to denote the extension of V to a real vector space via our chosen real embedding of K , and q will denote the unique extension of q to a quadratic form on $V(\mathbb{R})$. For the nonidentity σ_i , q^{σ_i} will denote the extension of q to the real vector space $V \otimes_{\sigma_i(K)} \mathbb{R}$.

We assume throughout this paper that q has signature $(n, 1)$ on $V(\mathbb{R})$ and that q^{σ_i} is definite for all $i \neq 1$. Given this setup, there is a natural injection

$$\begin{aligned} \mathfrak{i} : \mathrm{SO}(q, K) &\hookrightarrow \prod_{i=1}^s \mathrm{SO}(q^{\sigma_i}, \mathbb{R}), \\ &\cong \mathrm{SO}(n, 1) \times \prod_{i=2}^s \mathrm{SO}(n+1), \end{aligned}$$

sending $g \in \mathrm{SO}(q, K)$ to

$$\mathfrak{i}(g) = (g, \sigma_2(g), \dots, \sigma_s(g)) \in \mathrm{SO}(n, 1) \times \prod_{i=2}^s \mathrm{SO}(n+1).$$

In other words, we identify $g \in \mathrm{SO}(q, K)$ with its image in $\mathrm{SO}(n, 1)$ under projection onto the first factor of the above product.

Now, let \mathcal{O}_K denote the ring of integers of K . The above identifications map $\mathrm{SO}(q, \mathcal{O}_K)$ to an arithmetic lattice Γ_q in $\mathrm{SO}(n, 1)$. All arithmetic lattices in $\mathrm{SO}(n, 1)$ considered in this paper will be commensurable with some Γ_q as above. We briefly note that in even dimensions this construction determines every commensurability class of arithmetic lattices in $\mathrm{SO}(n, 1)$.

Given a quadratic form q as above, we form the hyperboloid model of hyperbolic space associated with q by

$$\mathbb{H}_q^n = \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} \mid q(x_1, \dots, x_{n+1}) = -1, x_{n+1} > 0\}.$$

The inner product on \mathbb{R}^{n+1} defined by (1) determines the usual hyperbolic metric on \mathbb{H}_q^n . From here forward, when there is no possibility of confusion we will drop the subscript and identify \mathbb{H}_q^n with \mathbb{H}^n .

A finite-volume hyperbolic orbifold is given as a quotient of \mathbb{H}^n by a lattice Γ in $\mathrm{Isom}(\mathbb{H}^n) \cong \mathrm{PO}(n, 1)$. The group $\mathrm{Isom}^+(\mathbb{H}^n)$ of orientation-preserving isometries is the connected component $\mathrm{SO}_0(n, 1)$ of the identity in $\mathrm{SO}(n, 1)$. If Γ is torsion-free then \mathbb{H}^n/Γ is a manifold, and otherwise it is a hyperbolic orbifold. In particular, the arithmetic lattice Γ_q described above determines an arithmetic hyperbolic orbifold \mathbb{H}^n/Γ . Selberg's lemma then implies that there is a finite index, torsion-free subgroup $\Gamma' < \Gamma$ and hence there is always an arithmetic hyperbolic manifold \mathbb{H}^n/Γ' that finitely covers \mathbb{H}^n/Γ .

For the reader's convenience, we also state the commensurability classification for arithmetic hyperbolic manifolds arising from the above construction. See [18, 2.6] and [27] for further details.

Theorem 2.1. *Let Γ, Γ' be two arithmetic lattices in $\mathrm{SO}(n, 1)$ constructed from quadratic spaces $(V, q), (V', q')$ over the number fields K, K' , respectively. Then Γ' is commensurable with Γ if and only if $K \cong K'$ and q' is similar to q , i.e., there exists $\lambda \in K^*$ such that (V', q') is isometric to $(V, \lambda q)$.*

We now briefly recall the construction of finite-volume codimension k immersed totally geodesic submanifolds of arithmetic hyperbolic manifolds arising from quadratic forms. Let Γ be an arithmetic lattice in $\text{Isom}(\mathbb{H}^n)$ associated with the quadratic space (V, q) over the number field K . Let $W \subset V$ be a (K -defined) codimension k subspace such that the restriction of q to W has signature $(n - k, 1)$ on $W(\mathbb{R})$. Then the intersection of $W(\mathbb{R})$ with $\mathbb{H}^n \subset V(\mathbb{R})$ defines a totally geodesic embedding $f : \mathbb{H}^{n-k} \hookrightarrow \mathbb{H}^n$. The induced embedding of $\text{SO}(W, q|_W)$ into $\text{SO}(V, q)$ yields an embedding

$$G_W = \text{S}(\text{O}(k) \times \text{O}(n - k, 1)) \cap \text{SO}_0(n, 1) \hookrightarrow \text{Isom}(\mathbb{H}^n),$$

such that $\Gamma_W = G_W \cap \Gamma$ is a lattice in G_W . In particular, $\mathbb{H}^{n-k}/\Gamma_W$ maps in as an immersed finite-volume totally geodesic submanifold of \mathbb{H}^n/Γ . Moreover, it is well-known that all geodesic submanifolds of \mathbb{H}^n/Γ arise from the above construction, (see [27]).

Remark 2.2. It is also known that arithmetic manifolds of the above kind are the only arithmetic manifolds that contain a codimension one finite volume totally geodesic submanifold. In particular, this means that the arithmetic manifolds described above are precisely those that can be used in the constructions of non-arithmetic manifolds that follow.

2.2. Constructions of non-arithmetic manifolds following Gromov and Piatetski-Shapiro. In this section, we recall the construction by Gromov and Piatetski-Shapiro of non-arithmetic hyperbolic manifolds [18], along with generalizations by Raimbault [30] and Gelander–Levit [16].

First, we recall the notion of a *building block* and an *arithmetic building block* from the introduction. As previously, a hyperbolic manifold will always mean a connected and oriented quotient of hyperbolic space with finite-volume. If M is a hyperbolic n -manifold, then a building block $N \subset M$ is an n -dimensional submanifold with nonempty boundary such that $\pi_1(N) < \pi_1(M)$ is Zariski dense in $\text{Isom}(\mathbb{H}^n)$ under the holonomy of the hyperbolic structure on M . We call N arithmetic when M is an arithmetic hyperbolic manifold. The following is evident from [18, 1.6.A].

Proposition 2.3. *Suppose that M is an arithmetic hyperbolic manifold and $N \subset M$ is an arithmetic building block. If $N \subset M'$ for M' another arithmetic hyperbolic manifold, then M' is commensurable with M .*

In particular, if N_1, N_2 are arithmetic building blocks with associated arithmetic hyperbolic manifolds M_1 and M_2 , it makes sense to say that they are *similar* if M_1 and M_2 are commensurable and *dissimilar* otherwise. We now return to constructing non-arithmetic hyperbolic manifolds.

The most basic Gromov–Piatetski-Shapiro construction is a hybrid manifold defined by interbreeding two incommensurable arithmetic hyperbolic manifolds. In our language, one builds a hyperbolic manifold consisting of two dissimilar arithmetic building blocks. Specifically, suppose that M_1 and M_2 are incommensurable arithmetic hyperbolic n -manifolds and that Σ is a hyperbolic $(n - 1)$ -manifold that admits totally geodesic embeddings

$$f_i : \Sigma \hookrightarrow M_i,$$

$i = 1, 2$. Suppose that $f_i(\Sigma)$ separates M_i , and choose one side N_i . This is then a hyperbolic manifold with connected totally geodesic boundary isometric to Σ , and N_i is an arithmetic building block in the sense described above. Gluing N_1 to N_2 along their boundaries then defines a non-arithmetic manifold comprised of two incommensurable arithmetic building blocks.

Further variants of this construction are given by Raimbault [30] and Gelander–Levit [16], who use a generalization of this idea to prove asymptotic counts on the number of hyperbolic manifolds in terms of volume. We briefly describe the constructions so the reader can explicitly see that they satisfy the assumptions of Theorem 1.4.

Raimbault’s variant of the Gromov–Piatetski-Shapiro construction glues together arithmetic building blocks in a circular pattern. More specifically, start with a family of r arithmetic building blocks N_1, \dots, N_r that are pairwise dissimilar and such that ∂N_j has two connected components, each being isometric to some fixed hyperbolic $(n-1)$ -manifold Σ (with opposite orientations). Let Σ_j^+ (resp. Σ_j^-) denote the positively (resp. negatively) oriented boundary component. One then glues N_j to N_{j+1} by identifying Σ_j^+ to Σ_{j+1}^- , $1 \leq j \leq r-1$, and N_r to N_1 by Σ_r^+ to Σ_1^- to obtain a non-arithmetic hyperbolic manifold built from r building blocks.

Gelander–Levit also construct large families of non-arithmetic manifolds, in which they start with any 4-regular and 2-colored graph G on k vertices such that every vertex has the same color except for one and such that each edge is given a label from the set $\{a, a^{-1}, b, b^{-1}\}$. They then form a graph of spaces where the vertex set V corresponds to two fixed arithmetic building blocks $\{V_0, V_1\}$ each with totally geodesic boundary with four connected components, each isometric to a fixed hyperbolic $(n-1)$ -manifold Σ . Moreover, the edge set E corresponds to gluing together two of four arithmetic building blocks from $\{A^-, A^+, B^-, B^+\}$ each with totally geodesic boundary with two connected components, each also isometric to the same Σ as before (see [16, Rem. 3.2] for details). Provided the building blocks are dissimilar, this constructs a non-arithmetic hyperbolic manifold.

3. CLOSURE RIGIDITY

The purpose of this section is to show that any totally geodesic submanifold with boundary lying on a cutting hypersurface can be extended to a finite-volume totally geodesic submanifold of the arithmetic manifold associated with each of the adjacent building blocks. More specifically, we prove the following theorem.

Theorem 3.1 (Closure Rigidity). *Let $M = \mathbb{H}^n/\Gamma$ be a finite-volume arithmetic hyperbolic n -manifold, $n \geq 3$, $\pi : \mathbb{H}^n \rightarrow M$ be the universal covering, and $\Sigma = \mathbb{H}^{n-1}/\Gamma_0 \subset M$ be an embedded oriented finite volume totally geodesic hypersurface. For $1 < m \leq n-1$, suppose that $f : \mathbb{H}^m \rightarrow \mathbb{H}^n$ is a totally geodesic embedding such that the image $(\pi \circ f)(\mathbb{H}^m) \subset M$ contains a connected finite-volume m -manifold with totally geodesic boundary N such that $\partial N \cap \Sigma \neq \emptyset$ and N is not freely homotopic into Σ . Then $(\pi \circ f)(\mathbb{H}^m)$ is an immersed finite-volume totally geodesic submanifold of M of dimension m .*

Proof. Since M contains a finite volume hypersurface, we have that M is defined by a quadratic form q on the vector space V defined over a number field K and Σ




FIGURE 1. The case when N has one boundary component.

comes from a K -defined codimension one subspace $V_0 \subset V$. It will be convenient to set $k = n - m$ as the codimension of $f(\mathbb{H}^m)$ in \mathbb{H}^n .

If Σ is nonseparating then let $M_1 = M$ cut open along Σ and otherwise suppose that Σ divides M into two submanifolds with boundary $M_1 \cup M_2$, $\partial M_1 = \overline{\partial M_2} = \Sigma$. Set $\Gamma_i = \pi_1(M_i)$ and $\Gamma_0 = \pi_1(\Sigma)$, considered as a subgroup of Γ . Without loss of generality, we can assume that $N \subset M_1$. Indeed, if every such component of N were freely homotopic into Σ , then N itself would be freely homotopic into Σ . Then ∂N contains a nonempty union N_1, \dots, N_r of immersed totally geodesic submanifolds of Σ (note that the N_i might intersect if N is not embedded). Let $\Delta = \pi_1(N) \subset \Gamma_1$ and $\Delta_i = \pi_1(N_i) \subset \Gamma_0$. Since $m \geq 2$, the N_i are not points, and therefore they each have a nontrivial fundamental group that maps injectively into Δ (cf. [18, 2.10.A]).

First, suppose that there is some $g \in \Delta$ that is not Δ -conjugate into some Δ_i (i.e., not freely homotopic into Σ). For example, this is necessarily the case when N has a unique boundary component, since N is not freely homotopic into Σ (see Figure 1). We now need some arithmetic notation.

Since N_1 is a finite-volume totally geodesic submanifold with boundary in Σ , it is associated with a K -defined subspace $V_1 \subset V_0$ of codimension k . Similarly, associated with $f : \mathbb{H}^m \rightarrow \mathbb{H}^n$ is a codimension k subspace $W(\mathbb{R}) \subset V(\mathbb{R})$ with the property that $V_1(\mathbb{R}) \subset W(\mathbb{R})$. It follows that $W(\mathbb{R})$ is a Δ -invariant subspace of $V(\mathbb{R})$, since the image of f contains a lift of the map on universal coverings $\tilde{N} \rightarrow \tilde{M}_1$. In fact, there exists a vector $w \in V(\mathbb{R})$ so that $W(\mathbb{R})$ is spanned by w and $V_1(K)$. We will show that we can take $w \in V(K)$, hence W is a K -defined subspace of V and $(\pi \circ f)(\mathbb{H}^m)$ then must determine an immersed finite-volume totally geodesic submanifold of M .

We have that $g(W(\mathbb{R})) = W(\mathbb{R})$, but $g(V_1(\mathbb{R})) \neq V_1(\mathbb{R})$, since g is not contained in Γ_0 . However, the fact that $g \in \Gamma_1$ implies that $g(V(K)) = V(K)$. In particular, $g(V_1(K)) \subset W(\mathbb{R})$ contains a vector $v \in V(K)$ that is not contained in $V_1(\mathbb{R})$. Therefore, the K -span of $V_1(K)$ and v is a codimension 1 subspace of $V(K)$ contained in $W(\mathbb{R})$. Thus W is defined over K as claimed.

Therefore, to conclude the proof it suffices to establish the existence of a $g \in \Delta$ that is not Δ -conjugate into some Δ_i , and hence is not conjugate into Γ_0 . In our setting N is a finite-volume hyperbolic manifold of dimension $m \geq 2$ with totally

geodesic boundary consisting of connected components N_i . We fix a reference isometric embedding of the universal cover \tilde{N} of N into the unit ball model for hyperbolic space \mathbb{H}^m . From this embedding, it is easy to see that

$$\partial^\infty \Delta \cong \partial^\infty \tilde{N} \hookrightarrow \partial^\infty \mathbb{H}^m \cong S^{m-1}$$

is an $(m-2)$ -dimensional Sierpinski curve in the sense of Cannon [9]. The boundary at infinity contains a countably infinite collection $\mathcal{S} = \{S_j\}$ of canonical *peripheral* $(m-2)$ -spheres, so each S_j is homeomorphic to S^{m-2} . These are the boundaries at infinity of the various lifts of the boundary components N_i . In terms of the action of Δ on $\partial^\infty \tilde{N}$, an element $g \in \Delta$ is Δ -conjugate into some Δ_i if and only if its fixed points $\text{Fix}(g) = \{g^+, g^-\}$ on the boundary are contained in a single peripheral sphere S_j .

Thus it is sufficient to find an element g whose fixed set $\text{Fix}(g)$ is not contained in any S_j . To see this, take any two distinct peripheral spheres, say S_1 and S_2 . These are a pair of disjoint closed subsets of the compact space $\partial^\infty \tilde{N}$, so are a positive distance $\delta > 0$ apart in the visual metric on the boundary. Recall that the respective visual metrics naturally make all the S_i round codimension one spheres in the ambient round sphere $S^{m-1} = \partial^\infty \mathbb{H}^m$.

In terms of the ambient round metric on S^{m-1} , there are only finitely many disjoint S_i of diameter greater than any given $\epsilon > 0$. Take a pair of open $\delta/3$ -neighborhoods U, V , of the peripheral spheres S_1, S_2 . Note that $U \cap V = \emptyset$, and any S_j that intersects both U and V must have diameter $> \delta/3$. Since the collection of disjoint such S_j is finite, the index subset J for these is finite. We can now find some $\delta' < \delta/3$ so that the δ' -neighborhood $U' \subset U$ has the property that $S_j \cap U' = \emptyset$ for all j in the finite set J .

It is well-known that for the Γ considered here, the set of pairs of the form $\text{Fix}(g)$ are dense in $\partial^\infty \Gamma \times \partial^\infty \Gamma$ [17, 8.2.F, 8.2.G]. Applying this to the open set $U' \times V \subset \partial^\infty \Delta \times \partial^\infty \Delta$, one obtains an element $g \in \Delta$ with the property that $g^+ \in U', g^- \in V$. From the choice of the sets U', V , it now follows that $\text{Fix}(g)$ is not contained in any peripheral sphere, and hence that g is not Δ -conjugate into any Δ_i . This completes the proof. \square

Remark 3.2. The assumption that $1 < m$ is critical. Indeed, in the case where N is a geodesic arc in M_1 connecting two distinct points in Σ , the proof completely falls apart in that N and ∂N have trivial fundamental group. However, the first step of the proof does work in the following sense. For example, suppose that Γ is an arithmetic Kleinian group that contains arithmetic Fuchsian subgroups. If γ_1, γ_2 are two purely hyperbolic elements of Γ that mutually preserve a totally geodesic hyperbolic plane in \mathbb{H}^3 , then $\langle \gamma_1, \gamma_2 \rangle$ is contained in an arithmetic Fuchsian subgroup of Γ .

4. ANGLE RIGIDITY

In this section we prove the following.

Theorem 4.1 (Angle Rigidity). *Fix a non-arithmetic hyperbolic manifold M that is built from building blocks, and suppose that two adjacent building blocks N_1 and N_2 are arithmetic and dissimilar. Let $A \subset M$ be a connected totally geodesic submanifold of dimension at least 2 such that A intersects the interior of N_1 and N_2 , i.e., crosses a cutting hypersurface Σ . Then A meets Σ orthogonally.*




FIGURE 2. The proof of Angle Rigidity. Subspaces of \tilde{N}_i associated with geodesic subspaces of M_i (green) are matched only if they meet Σ orthogonally.

As remarked in the introduction, a special case of this result was recently and independently obtained by [6].

Proof. The basic idea of the proof is as follows. Each N_i is associated with an arithmetic manifold M_i . That the M_i share isometric codimension 1 submanifolds allows us to assume they are defined via quadratic forms over the same field K [27, §6]. Closure Rigidity allows us to assume that A is built from gluing submanifolds A_i of N_i , where $A_i = N_i \cap B_i$ for B_i a totally geodesic submanifold of M_i . We will show when M_1 is not commensurable with M_2 that such a matching can only occur when the B_i meet Σ orthogonally. See Figure 2. The key point is that such submanifolds B_i arise from certain K -defined subspaces of the K -vector space for M_i , but the identification between half-spaces in \mathbb{H}^n arising from gluing N_2 to N_1 is defined by a map of real vector spaces that preserves definition over K only for very special subspaces.

We first carefully explain why Closure Rigidity applies. Set $A_i = N_i \cap A \subset M_i$. This defines an immersed totally geodesic submanifold of M_i with totally geodesic boundary that meets the cutting hypersurface Σ nontrivially. If $f_i : \mathbb{H}^m \hookrightarrow \mathbb{H}^n$ denotes the extension to \mathbb{H}^m of the lift $\tilde{A}_i \rightarrow \tilde{M}_i$, we see that \tilde{A}_i satisfies the assumptions of Theorem 3.1. The existence of the totally geodesic submanifold B_i as above follows.

Next we set up some notation and make some simplifying assumptions. Suppose that N_1 and N_2 are dissimilar arithmetic building blocks glued together by an orientation-reversing isometry of the connected totally geodesic boundary component Σ . Being arithmetic, each N_i has an associated quadratic space (V_i, q_i) , which is well defined up to similarity. The isometry $\varphi : \Sigma \rightarrow \bar{\Sigma}$ determines a similarity between codimension one subspaces H_i of V_i on which the restriction of q_i has signature $(n - 1, 1)$. We can change the similarity class of q_2 and assume that the similarity $H_2 \rightarrow H_1$ is an isometry.

Therefore, we can assume there is a quadratic space (H, q) of dimension n and signature $(n - 1, 1)$ that is isometric to a codimension one subspace of each (V_i, q_i) .

Using Gram–Schmidt, we can write

$$(V_i, q_i) = (V_i, \langle \alpha_i \rangle \oplus q),$$

for some $\alpha_i \in K^*$. More specifically, we can choose coordinates $\{x_0, \dots, x_n\}$ on V_1 and $\{y_0, \dots, y_n\}$ on V_2 such that

$$\begin{aligned} q_1(x_0, \dots, x_n) &= \alpha_1 x_0^2 + q(x_1, \dots, x_n), \\ q_2(y_0, \dots, y_n) &= \alpha_2 y_0^2 + q(y_1, \dots, y_n). \end{aligned}$$

The isometry between the models $\mathbb{H}_{q_i}^n$ of hyperbolic space associated with q_2 and q_1 is then induced by the map $V_2(\mathbb{R}) \rightarrow V_1(\mathbb{R})$ defined by $y_0 \mapsto \sqrt{\alpha_2/\alpha_1} x_0$ and $y_i \mapsto x_i$ for $1 \leq i \leq n$ (compare with [18, §2.9]). Let Φ denote this map, and note that Φ determines a map of K -vector spaces $V_2 \rightarrow V_1$ if and only if α_2/α_1 is a square in K^* . By Witt cancellation, this holds if and only if q_1 is isometric to q_2 , which is definitely not the case if M_1 and M_2 are noncommensurable.

Consider a codimension k totally geodesic subspace B_i of the arithmetic manifold M_i associated with q_i . Then B_i arises from a codimension k subspace W_i of the K -vector space V_i on which the restriction of q_i has signature $(n-k, 1)$. Furthermore, suppose that the intersection of B_i with Σ contains a codimension k totally geodesic submanifold of Σ . Then, for the correct choice of base point, W_i intersects H in a codimension k subspace U_i of H on which the restriction has signature $(n-k-1, 1)$. Thus W_i is generated by U_i and a vector not in H .

Turning this around, we can construct all the submanifolds B_i as above by starting with a codimension k subspace U_i of H then taking the subspace of V_i generated by U_i and another vector ξ_i that is not contained in H . Note that ξ_i is not in H if and only if $y_0 \neq 0$. Let W_i be the span over K of ξ_i and U_i . To have B_1 glue to B_2 means precisely that Φ takes $W_2(\mathbb{R})$ to $W_1(\mathbb{R})$.

Notice that $\Phi(U_2) = U_1$, since $\Phi(H) = H$. If ξ_2 has coordinates y_i , then

$$\Phi(\xi_2) = \left(\sqrt{\alpha_2/\alpha_1} y_0, y_1, \dots, y_n \right) \in V_1(\mathbb{R}).$$

If α_2/α_1 fails to be a square in K^* , we see that $\Phi(W_2(\mathbb{R}))$ is $W_1(\mathbb{R})$ for W_1 a K -defined subspace of V_1 if and only if $y_1 = \dots = y_n = 0$. This proves that such a K -defined subspace of V_2 maps to a K -defined subspace of V_1 if and only if $W_i(\mathbb{R})$ is the \mathbb{R} -span of U_i and the vector $(1, 0, \dots, 0)$. The same argument shows that this again holds with the roles of V_1 and V_2 reversed.

Geometrically, this proves that the submanifolds B_i of M_i that could possibly give rise to A as in the statement of the theorem are associated with subspaces W_i of V_i generated by a codimension one subspace U_i of H along with the vector $(1, 0, \dots, 0)$. It remains to compute that such a B_i intersects Σ orthogonally.

The subspace of V_i associated with Σ is the q_i -orthogonal complement to $e = (1, 0, \dots, 0)$. The subspace associated with B_i is the orthogonal complement W_i to a subspace Z_i of V_i with dimension k . Since $(1, 0, \dots, 0) \in W_i$, Z_i must be a subspace of H . However, the angle θ_i between B_i and Σ satisfies

$$\cos^2 \theta_i = \sup_{z \in Z_i} \frac{\langle e, z \rangle_{q_i}^2}{q_i(e)q_i(z)}.$$

See [31, p. 71]. Since $Z_i \subseteq H$, z is q_i -orthogonal to e for all $z \in Z_i$, and we see that the above expression is identically zero. This shows that B_i meets Σ orthogonally, as desired. \square

Remark 4.2. We briefly note that the proof of Theorem 4.1 does not appear to use anything about the assumption that A has dimension at least two. This assumption is hiding in the application of Closure Rigidity in the very first step of the proof, since Closure Rigidity fails when A is a geodesic. See Remark 3.2.

Remark 4.3. With a bit more work, one can show that this approach proves a stronger result. Suppose M_1, M_2 are arithmetic hyperbolic n -manifolds containing an isometric totally geodesic submanifold Σ . Suppose that there is a further totally geodesic subspace $Z_0 \subset \Sigma$ of codimension $1 \leq k < n-1$ and a codimension k totally geodesic submanifold N_i of M_i that meets Σ at Z_0 with angle θ_i . If $\theta_2 = \pm\theta_1$, then M_2 is commensurable with M_1 . One simply uses the above decomposition of q_1 and q_2 to directly build an isometry between the quadratic spaces.

5. EQUIDISTRIBUTION, HOMOGENEOUS DYNAMICS, AND PROOFS OF THE MAIN THEOREMS

In this section, we complete the proof of Theorem 1.4. We will first give some geometric reductions in Section 5.1 that recast the problem as a dynamical one. For the reader's convenience, we then give a simple proof in Section 5.2 in the special case of closed hypersurfaces in closed manifolds. We then go on to give a proof of the general form of Theorem 1.4, first in the cocompact case in Section 5.3, and finally in the finite-volume case in Section 5.4.

5.1. Some geometric preliminaries. For a Riemannian manifold M , we denote by $\text{OF}(M)$ the bundle of *oriented* orthonormal frames on M and for any $1 \leq m \leq n$ we denote by $G^m M$ the bundle over M whose fiber over a point p is the Grassmannian of m -dimensional subspaces in $T_p M$. For each m , there is a natural bundle map $\text{OF}(M) \rightarrow G^m M$ obtained by sending an orthonormal frame (v_1, \dots, v_n) at a point p to the m -dimensional subspace spanned by v_1, \dots, v_m . The fiber of the bundle map $\text{OF}(M) \rightarrow G^m M$ can naturally be identified with

$$S(\text{O}(m) \times \text{O}(n-m)) = \{(A, B) \in \text{O}(m) \times \text{O}(n-m) : \det(A)\det(B) = 1\}.$$


Given an immersion $N \rightarrow M$ with $\dim(N) \leq m$ and a sufficiently small neighborhood $U \subset N$ of a point $x \in N$, the inverse function theorem induces a map $G^m U \rightarrow G^m M$, which then induces a map $G^m N \rightarrow G^m M$.

We now spend the rest of this subsection proving the following proposition.

Proposition 5.1. *Let M be an n -dimensional hyperbolic manifold built from building blocks containing two adjacent, arithmetic, and dissimilar building blocks. Then there is an open subset $\Omega_m \subseteq G^m M$ such that $G^m N \cap \Omega_m = \emptyset$ for any any finite-volume immersed totally geodesic submanifold $N \subset M$ of dimension $2 \leq m < n$.*

Proof. Fix $m \geq 2$. By assumption, there is a totally geodesic cutting hypersurface $\Sigma \subset M$ contained in the common boundary of the arithmetic building blocks. We know from Theorem 4.1 that any closed immersed totally geodesic m -dimensional submanifold $N \rightarrow M$ intersects Σ orthogonally. That the submanifold N intersects Σ orthogonally means that $(T_x N)^\perp \subseteq T_x \Sigma$ for every $x \in N \cap \Sigma$.

Choose a compact core C for M [22, p. 156] and $x \in \Sigma \cap C$. Fix a small neighborhood U of x in $\Sigma \cap C$, and let $\delta > 0$ be the injectivity radius of M on U and $N_{\delta/2}(U) \subset M$ denote the open $\delta/2$ -neighborhood of U . Then for any point $p \in \overline{N_{\delta/2}(U)}$, there is a unique minimal length geodesic γ_p from p to U .

FIGURE 3. Choosing ϵ_0 in the proof of Proposition 5.1.

We form the subset

$$\mathcal{V} = \{(p, W) \in G^m M \mid p \in N_{\delta/2}(U), \dot{\gamma}_p \in W\},$$

and denote by \mathcal{V}_p the subset of \mathcal{V} lying above a point $p \in N_{\delta/2}(U)$. Note that $\mathcal{V} \rightarrow N_{\delta/2}(U)$ is a fiber bundle over an open subset of M with fiber \mathcal{V}_p a closed subset of the corresponding Grassmannian fiber of $G^m M$. Indeed, \mathcal{V}_p is a copy of $Gr(m-1, \mathbb{R}^{n-1})$ lying within the $Gr(m, \mathbb{R}^n)$ fiber of $G^m M$.

Take a geodesic γ in \mathbb{H}^2 , a point A at distance $\delta/2$ from γ , and let B denote the projection of A onto γ . Set $0 < \epsilon_0 < \pi/2$ to be the angle between the geodesic \overline{AB} from A to B and the geodesic ray α from A to one of the endpoints of γ on $\partial^\infty \mathbb{H}^2$. See Figure 3. In particular, notice that any geodesic arc in \mathbb{H}^2 starting at A with initial angle $0 < \epsilon < \epsilon_0$ with \overline{AB} then must intersect γ in \mathbb{H}^2 .

Take $0 < \epsilon < \epsilon_0$, and consider the set $N_\epsilon(\mathcal{V})$ defined by taking the fiber-wise ϵ -neighborhood of the set \mathcal{V} , measured in the angular metric on the Grassmannian. Thus a pair (p, W) lies in $N_\epsilon(\mathcal{V})$ if and only if $p \in N_{\delta/2}(U)$ and there exists a $W' \in \mathcal{V}_p$ such that $\theta(W, W') < \epsilon$. Finally, we let

$$\Omega_m = N_\epsilon(\mathcal{V}) \setminus \mathcal{V},$$

and note that Ω_m is an open subset of $G^m M$.

We now check that $G^m N \cap \Omega_m = \emptyset$ for any finite-volume immersed m -dimensional totally geodesic submanifold N of M . By way of contradiction, assume that there exists a point $(p, W) \in G^m N \cap \Omega_m$. Since N has dimension m , this means that $p \in N$, $W = T_p N \subset T_p M$, and $0 < \theta(W, \mathcal{V}_p) < \epsilon$. This last inequality tells us that $0 < \theta(W, \dot{\gamma}_p) < \epsilon$, where γ_p is the unique geodesic from $p \in N_{\delta/2}(U)$ to $U \subset \Sigma$.

We let $w \in W$ denote the projection of $\dot{\gamma}_p$ to $T_p N$, and consider the subspace $Z = \text{span}\{w, \dot{\gamma}_p\}$ of $T_p M$. We have that $\dim(Z \cap T_p N) = \dim(Z \cap W) = 1$, and hence Z defines a (not necessarily closed) geodesic contained in the submanifold N . It also contains the direction vector $\dot{\gamma}_p$ corresponding to the minimal geodesic from p to Σ . Moreover, we have that

$$z = \dot{\gamma}_p - w \in (T_p N)^\perp,$$

is a nonzero vector, hence $\dim(Z \cap (T_p N)^\perp) = 1$.

The 2-dimensional subspace Z gives rise to an isometric immersion $f : \mathbb{H}^2 \rightarrow M$ whose image contains both geodesics γ_p and η . We let q denote the terminal point




FIGURE 4. The geodesic triangle in Proposition 5.1.

of γ_p in the open subset U of the hypersurface Σ . Since $\theta(\dot{\gamma}_p, \dot{\eta}) = \theta(\dot{\gamma}_p, w) < \epsilon$, our choice of $\epsilon < \epsilon_0$ implies that the geodesic η intersects Σ at a point q' . Since $\dot{\eta} \in T_p N$, we see that η is contained entirely in the submanifold N , and hence that $q' \in N \cap \Sigma$.

Notice that we now have a geodesic triangle contained in the isometrically immersed hyperbolic plane $f(\mathbb{H}^2)$, consisting of the geodesic segment γ_p , the geodesic segment η , and the geodesic ξ joining the two endpoints $q, q' \in \Sigma$. See Figure 4. Since Σ is totally geodesic, we see that ξ is contained entirely in Σ , and basic hyperbolic trigonometry implies that $\theta_{q'}(\eta, \xi) < \pi/2$, since $\theta_q(\eta, \gamma_p) = \pi/2$.

We claim that N fails to be orthogonal to Σ at the point q' . To see this, we need to exhibit a vector in $(T_{q'}N)^\perp$ that does not lie in the subspace $T_{q'}\Sigma$. Recall that $z \in Z$ was a vector orthogonal to $T_p N$, and it is tangent to the immersed totally geodesic hyperbolic plane $f(\mathbb{H}^2)$. We can thus parallel transport this orthogonal vector along $\eta \subset N$ to obtain a vector $z' \in T_{q'}M$. Since parallel transport preserves orthogonality, we see that $z' \in (T_{q'}N)^\perp$. Also, $z \in T_p f(\mathbb{H}^2)$, and η is contained in $f(\mathbb{H}^2)$, which forces $z' \in T_{q'} f(\mathbb{H}^2)$. Since we have $z \perp \dot{\eta}(p)$, it follows that $z' \perp \dot{\eta}(q')$.

Finally, focusing on the behavior at $T_{q'}M$, we have three vectors that all lie in the 2-dimensional subspace $T_{q'} f(\mathbb{H}^2) \subset T_{q'}M$: the vector $\dot{\xi}$, which spans the 1-dimensional subspace

$$T_{q'} f(\mathbb{H}^2) \cap T_{q'} \Sigma,$$

the vector $\dot{\eta}$, which we argued forms an angle $\theta_{q'}(\dot{\eta}, \dot{\xi}) < \pi/2$ with $\dot{\xi}$, and the vector $z' \in (T_{q'}N)^\perp$ that is orthogonal to $\dot{\eta}$. It immediately follows that z' cannot be parallel to $\dot{\xi}$, but $\dot{\xi}$ generates the intersection of $(T_{q'}N)^\perp$ with $T_{q'}\Sigma$, i.e., we have constructed an element of $(T_{q'}N)^\perp$ that cannot be in $T_{q'}\Sigma$, and hence N is not orthogonal to Σ .

This contradicts Theorem 4.1, so we conclude that the set $G^m N \cap \Omega_m$ is indeed empty, completing the proof of the proposition. \square

5.2. Codimension 1. In this subsection, we show how Proposition 5.1 can be used to complete the proof of Theorem 1.4 for the case of closed hypersurfaces in a compact manifold. For this, we use the following result of Shah [37, Thm. B] from homogeneous dynamics.

Theorem 5.2. *Suppose $n \geq 3$, $G = SO_0(n, 1)$, $\Gamma < G$ a cocompact lattice, and $W = SO_0(n - 1, 1)$. Then every W invariant subset of G/Γ is either dense or the union of finitely many closed W orbits.*

It is worth mentioning that one can also prove Theorem 5.2 by using Ratner's work on invariant measures for unipotent flows [32] and work of either Ratner or Dani–Margulis on equidistribution and orbit closures [33, 11]. In fact we will use this approach in Section 5.3. The proof given in Shah's paper follows instead Margulis' original proof of the Oppenheim conjecture using topological dynamics [23, 10]. It seems possible that those methods might prove all the results we need in this paper, but we do not pursue this here.

The following standard lemma shows how immersed totally geodesic submanifolds correspond to closed W orbits in G/Γ .

Lemma 5.3. *Let M be a closed hyperbolic manifold of dimension n and $N \subset M$ a closed immersed totally geodesic hypersurface. Then there are either one or two closed W orbits in G/Γ which project onto N . Furthermore, distinct immersed totally geodesic hypersurfaces give rise to distinct W orbits.*

Proof. Given a closed immersed totally geodesic hypersurface N , its lift \tilde{N} to the universal cover $\tilde{M} = \mathbb{H}^n$ has stabilizer

$$H = g^{-1} O^+(n - 1, 1)g,$$

where g is any isometry taking \tilde{N} to the standard \mathbb{H}^{n-1} stabilized by

$$O^+(n - 1, 1) = S(O(1) \times O(n - 1, 1)) \cap SO_0(n, 1).$$

Since N is closed, $H \cap \Gamma$ is a cocompact lattice. Writing $O^+(n - 1, 1)g = gH$ we see that the $O^+(n - 1, 1)$ orbit of $[g]$ is the g translate of $H/H \cap \Gamma$ in G/Γ . As $O^+(n - 1, 1)$ contains $SO_0(n - 1, 1)$ as a subgroup of index two, this $O^+(n - 1, 1)$ orbit is either one or two $SO_0(n - 1, 1)$ orbits giving the first claim. Since distinct choices of N give rise to distinct groups H , the last statement is clear. \square

We are now in a position to conclude Theorem 1.4 in our special case.

Theorem 5.4. *Let M be a hyperbolic manifold built from building blocks with two adjacent, arithmetic, and dissimilar building blocks. Then there are only finitely many closed immersed totally geodesic hypersurfaces in M .*

Proof. Denote by Ω_{n-1} the open set provided by Proposition 5.1 applied when $m = n - 1$ and by V its pre-image in G/Γ . By Proposition 5.1 and Lemma 5.3, the W orbits arising from all closed immersed totally geodesic hypersurfaces in M have trivial intersection with V . By Theorem 5.2 we thus have a finite collection of W orbits and hence we must only have finitely many totally geodesic closed immersed hypersurfaces in M . \square

5.3. Theorem 1.4 in the compact case. For simplicity, we write this subsection assuming that M is compact, that is to say that $\Gamma < G$ is a cocompact lattice. Throughout we assume M satisfies the hypotheses of Theorem 1.4 and use the conventions that $m = n - k$ and $W_m = SO_0(m, 1)$. Our choice of W_m ensures that it is connected and generated by one-parameter unipotent subgroups.

We begin by recalling some notation and a result of Dani–Margulis. Let \mathcal{H} be the collection of subgroups H of G such that $H \cap \Gamma$ is a lattice in H and $\text{Ad}(H \cap \Gamma)$ is

Zariski dense in $\text{Ad}(H)$. Given $H \in \mathcal{H}$ and a group $W \subset G$ generated by unipotent elements, define

$$X(H, W) = \{g \in G \mid Wg \subset gH\}.$$

We remark for later that the following is immediate from the definition.

Lemma 5.5. *If $x \in G$, then $X(H, xWx^{-1}) = xX(H, W)$.*

We now recall a special case of [11, Thm. 3], where the advantage of assuming that Γ is cocompact is that we may take $K = G/\Gamma$ in their notation. (We emphasize here that our K below will be the maximal compact subgroup of G .)

Theorem 5.6. *Let G be a Lie group and $\Gamma < G$ a cocompact lattice. Let μ be the Haar measure on G/Γ . Let $U = \{u_t\}$ be a one parameter Ad-unipotent subgroup of G and let ϕ be a bounded continuous function on G/Γ . Fix $\epsilon > 0$, then there exist finitely many subgroups $H_1, \dots, H_r \in \mathcal{H}$ such that for $x \notin \cup_{i=1}^r X(H_i, U)\Gamma/\Gamma$ there exists a $T_0 \geq 0$ such that*

$$\left| \frac{1}{T} \int_0^T \phi(u_t x) dt - \int_{G/\Gamma} \phi d\mu \right| < \epsilon,$$

for all $T > T_0$.

The theorem says that any u_t trajectory not contained in any $X(H_i, U)$ eventually has time average for ϕ equal to the space average for ϕ . Let $\pi_m : \text{OF}(M) \rightarrow G^m M$ be the bundle map defined in §5.1, Ω_m be the open set provided by Proposition 5.1, and let $V_m = \pi_m^{-1}(\Omega_m)$. For our purposes we will always take ϕ_m to be a compactly supported function of total integral 1 with support in V_m . Then any unipotent trajectory in $\text{OF}(M)$ corresponding to part of the horocycle flow of a closed immersed totally geodesic submanifold must have

$$\frac{1}{T} \int_0^T \phi_m(u_t x) dt = 0,$$

for all T . In particular, the entire trajectory is contained in

$$\bigcup_{i=1}^r X(H_i, U)\Gamma/\Gamma,$$

by Theorem 5.6.

In the remainder of this section, we show that all pre-images of immersed totally geodesic submanifolds are contained in

$$\bigcup_{i=1}^r X(H_i, W_m)\Gamma/\Gamma,$$

for some finite collection of subgroups $H_i \in \mathcal{H}$, then give a concrete description of each $X(H_i, W_m)$. This is the content of Lemma 5.7 and Proposition 5.11. The geometric translations in Corollaries 5.8 and 5.12 will then give the requisite finiteness statement. We now begin with a key reduction for our proof, noting that W_m is generated by unipotent subgroups.

Lemma 5.7. *Fix m between 2 and $n - 1$, let $W = W_m$, and let U_1, \dots, U_s be unipotent subgroups generating W . If $H \in \mathcal{H}$, then*

$$X(H, W) = \bigcap_{i=1}^s X(H, U_i).$$

Proof. It is clear that $X(H, W) \subset X(H, U_i)$ for all $1 \leq i \leq s$. Taking the derivative of the defining equation of $X(H, W)$ we see that $g \in X(H, W)$ if and only if $\text{Ad}(g^{-1})\mathfrak{w} \subset \mathfrak{h}$. Consequently if $g \in X(H, W)$, then $\text{Ad}(g^{-1})\mathfrak{u} \subset \mathfrak{h}$ for any connected subgroup $U < W$. As \mathfrak{h} is a subalgebra, if $\text{Ad}(g^{-1})\mathfrak{u}_i \subset \mathfrak{h}$ for all $1 \leq i \leq s$ then $\text{Ad}(g^{-1})\mathfrak{w} \subset \mathfrak{h}$. This gives the reverse containment. \square

Corollary 5.8. *Fix m between 2 and $n - 1$, and let $K \cong \text{SO}(n)$ denote the maximal compact subgroup of $G = \text{SO}_0(n, 1)$. Then there are finitely many subgroups $H_{1,m}, \dots, H_{j_m,m}$ of G such that the set of all closed immersed totally geodesic submanifolds of M of dimension m is contained in*

$$\mathcal{X}_m = K \backslash K \left(\bigcup_{i=1}^{j_m} X(H_{i,m}, W_m) \Gamma / \Gamma \right) \subset M.$$

In particular, the set of all closed immersed totally geodesic submanifolds of M of dimension between 2 and $n - 1$ is contained in the union of the \mathcal{X}_m .

Proof. Fix some m between 2 and $n - 1$. By definition, the pre-image in G/Γ of any closed immersed totally geodesic submanifold N of M with dimension m has empty intersection with V_m . Arguing as in Lemma 5.3, we see that $\tilde{N} \subset \tilde{M}$ is stabilized by a conjugate of

$$L = \text{S}(\text{O}(n - m) \times \text{O}(m, 1)) \cap \text{SO}_0(n, 1).$$

Again as in Lemma 5.3, there is a closed L orbit in G/Γ covering N which is clearly W_m invariant (since $W_m < L$). Therefore the set of all closed immersed totally geodesic submanifolds of M is contained in a W_m invariant set avoiding V_m . Picking any bounded continuous function ϕ_m with support in V_m and average $\int \phi_m d\mu = 1$, the statement is then immediate from Theorem 5.6 and Lemma 5.7 for totally geodesic submanifolds of dimension m . Repeating the argument in each dimension establishes the last statement in the corollary and completes the proof. \square

To complete the proofs of our main theorems, we now need to compute the sets $X(H, W_m)$ for any $H \in \mathcal{H}$. Note that by definition any H for which $X(H_i, W_m)$ is non-empty contains a conjugate of W_m or equivalently H is a conjugate of a group containing W_m . The following lemma is elementary and contained in, for example, Einsiedler–Wirth [14, Cor. 3.2].

Lemma 5.9. *Any closed subgroup $H < G = \text{SO}_0(n, 1)$ containing W_m is of the form*

$$\text{S}(K \times \text{O}(\ell, 1)) \cap G,$$

where $\ell \geq m$ and K is a compact subgroup of $\text{O}(n - \ell)$.

We also need an elementary algebraic lemma concerning W_ℓ , H , and G .

Lemma 5.10. *Fix m between 2 and $n - 1$, let $W = W_m$, and let H be any subgroup satisfying $W < H < G$. Then any subgroup W' of H isomorphic to W is conjugate to W in H . Furthermore, $\text{Aut}(W) < G$ and $\text{Aut}(W) < H$ unless $H = K \times \text{SO}(m, 1)$.*

Proof. For the first statement, we can assume $H = \text{SO}_0(\ell, 1)$ since any subgroup $W' \cong \text{SO}_0(m, 1)$ is obviously contained in a subgroup of H isomorphic to $\text{SO}_0(\ell, 1)$. In this case we identify $\text{Isom}(\mathbb{H}^\ell)$ with H by writing $\mathbb{H}^\ell = K \backslash H$, where $K = \text{SO}(\ell)$ is the maximal compact subgroup of H . The orbits of W and W' in \mathbb{H}^ℓ are

totally geodesic copies of \mathbb{H}^m . Since $\mathrm{SO}_0(\ell, 1)$ acts transitively on totally geodesic embeddings of \mathbb{H}^m , there is an isometry $g \in H$ carrying the W orbit to the W' orbit and hence $g^{-1}Wg < W'$. The last statement follows similarly as $\mathrm{Aut}(W)$ is simply the group of all isometries of \mathbb{H}^m , including the orientation-reversing isometries, and these can be realized in $\mathrm{Isom}(\mathbb{H}^\ell)$. The only case where these are not all in H is when H contains only $\mathrm{SO}_0(m, 1)$ and not $\mathrm{O}^+(m, 1)$. \square

We now describe the possible $X(H, W_m)$, where in what follows we write $Z(W)$ for the centralizer of a subgroup W in G .

Proposition 5.11. *Fix m between 2 and $n - 1$, let $W = W_m$, and fix $H \in \mathcal{H}$ for which $X(H, W)$ is nonempty. Then either there exists $x \in G$ such that $X(H, W) = Z(W)xH$ or there exists $x_1, x_2 \in G$ such that $X(H, W) = Z(W)x_1H \cup Z(W)x_2H$.*

Proof. Conjugating W by a suitable element $x \in G$ we may assume that $W' = x^{-1}Wx < H$. We first show the corresponding result for W' .

It is clear that $Z(W')H \subset X(H, W')$ and we need only prove the reverse inclusion. To this end, let $y \in X(H, W')$ and define $W'' = y^{-1}W'y$ to be the resulting subgroup of H . Then from Lemma 5.10, we know that W'' is conjugate to W' by an element $h \in H$. When $\mathrm{Aut}(W') < H$, we may moreover choose h so that conjugating W' by yh is the identity on W' . This immediately implies that $yh \in Z(W')$ and hence $y \in Z(W')H$. When $\mathrm{Aut}(W')$ is not contained in H , we can only guarantee that yh is either in $Z(W')$ or is of the form zf where $z \in Z(W')$ and $f \in G$ is any fixed element of G inducing the outer automorphism of W . Hence either $X(H, W') = Z(W')H$ or $X(H, W') = Z(W')H \cup Z(W')fH$ as claimed.

For the general case, we use Lemma 5.5 to compute $X(H, W)$. Since

$$Z(W') = Z(x^{-1}Wx) = x^{-1}Z(W)x,$$

in the first case we get that $X(H, W) = xx^{-1}Z(W)xH = Z(W)xH$ and in the second case we get that $X(H, W) = Z(W)xH \cup Z(W)xfH$. \square

Corollary 5.12. *For any fixed $H \in \mathcal{H}$ and m between 2 and $n - 1$, the subset $K \backslash KX(H, W_m)\Gamma/\Gamma \subset M$ is either one or two totally geodesic submanifolds of M .*

Proof. It is enough to see that $K \backslash KZ(W)xH\Gamma/\Gamma$ is as described. Since $Z(W) \subset K$, it can be dropped from the expression and we have $K \backslash KxH\Gamma/\Gamma$. By Lemma 5.9, $H = S(C \times \mathrm{O}(\ell, 1))$ where C is a subgroup of K and $H \cap \Gamma$ is a lattice. Hence we are reduced to the K, Γ double coset of xH which is precisely a closed immersed totally geodesic submanifold of M . \square

Theorem 1.4 in the case where M is compact is now immediate from Corollary 5.8 and Corollary 5.12.

5.4. Theorem 1.4 in the general finite-volume case. The goal of this subsection is to prove an analogue of Corollary 5.8 in the case when M has finite volume, after which one is reduced to repeating the proof in Section 5.3. There is some difficulty in extending the argument above to the case of finite covolume, since the general version of Theorem 5.6, stated below, is more complicated. Our main tool for circumventing this issue is the following simple geometric lemma. We briefly emphasize that, in this subsection we use ‘closed’ in the sense of orbits, so any finite-volume totally geodesic submanifold is closed in what follows.

Lemma 5.13. *Let M be a noncompact finite volume hyperbolic manifold of dimension n . Then there is a compact set $C \subset M$ such that any closed immersed totally geodesic submanifold N in M of dimension between 2 and $n - 1$ intersects C in an open set in N .*

Proof. Note that each N has fundamental group $\pi_1(N)$ which is a lattice in some $\mathrm{SO}(m, 1)$ and injects in $\pi_1(M)$. Moreover, the cusps of M have solvable fundamental groups and consequently $\pi_1(N)$ admits no embeddings into these groups. Let C_1 be a compact core of M [22, p. 156], so $M \setminus C_1$ is entirely contained in the cusps. Then the image of every N in M must intersect C_1 . Letting C be a compact subset of M containing a neighborhood of C_1 , gives the requisite compact set. \square

We now state the general form of the theorem of Dani–Margulis [11, Thm. 3], where in their language we assume that $K = C$, $F = \{x\}$, and the C_i are simply all of $X(H_i, U)$.

Theorem 5.14. *Let G be a connected Lie group and $\Gamma < G$ a lattice and μ be the G invariant probability measure on G/Γ . Let $U = \{u_t\}$ be any one parameter subgroup of G and ϕ a bounded continuous function on G/Γ . Let C be a compact subset of G/Γ and $\epsilon > 0$ be given. Then there exist finitely many proper closed subgroups H_1, \dots, H_r such that $H_i \cap \Gamma$ is a lattice in H_i and such that for any x in*

$$C \setminus \left(\bigcup_{i=1}^r X(H_i, U)\Gamma/\Gamma \right),$$

there exists $T_0 \geq 0$ such that

$$\left| \frac{1}{T} \int_0^T \phi(u_t x) dt - \int_{G/\Gamma} \phi d\mu \right| < \epsilon,$$

for all $T > T_0$.

The need for Lemma 5.13 comes from the fact that the H_i depend on C in this statement. We believe this is an artifact of the proof given in [11], but modifying their proof to avoid this dependence requires significant changes. We now show how to combine Lemma 5.13 and Theorem 5.14 to obtain our desired finiteness result.

Proposition 5.15. *Fix m between 2 and $n - 1$, let $W_m = \mathrm{SO}(m, 1)$, and let M be any finite volume hyperbolic manifold satisfying the hypotheses of Theorem 1.4. Then there are finitely many subgroups $H_{1,m}, \dots, H_{j_m,m}$ of G such that the set of all closed immersed totally geodesic submanifolds of M of dimension m is contained in*

$$\mathcal{X}_m = K \setminus K \bigcup_{i=1}^{j_m} X(H_{i,m}, W_m)\Gamma/\Gamma,$$

where K is the maximal compact subgroup of G . In particular, the set of all closed immersed totally geodesic submanifolds of M of dimension between 2 and $n - 1$ is contained in the union of the \mathcal{X}_m .

Proof. As in the proof of Corollary 5.8, we see that each totally geodesic submanifold of M of dimension m gives rise to a closed orbit of

$$L = \mathrm{S}(\mathrm{O}(n - m) \times \mathrm{O}(m, 1)) \cap \mathrm{SO}_0(n, 1).$$

By Lemma 5.13, there is a compact subset C in G/Γ such that any closed L orbit intersects C . We also know that there are finitely many unipotent subgroups U_1, \dots, U_{s_m} that generate $W_m = \text{SO}_0(m, 1) < L$, and so

$$X(H, W_m) = \bigcap_{j=1}^{s_m} X(H, U_j),$$

for any $H \in \mathcal{H}$.

We apply Theorem 5.14 to each U_j using C as the compact set, taking ϵ to be any fixed number $0 < \epsilon < 1$, and with ϕ_m a compactly supported function with integral 1 and support in $V_m = \pi^{-1}(\Omega_m)$. Given a closed L orbit \mathcal{O} , we pick a point $x \in \mathcal{O} \cap C$. For any U_j , the time average of ϕ_m over the x orbit is zero and so by the above, there are finitely many $H_{i,j,m}$ such that x is in one of the $X(H_{i,j,m}, U_j)$. Taking intersection over j and re-indexing the resulting collection of subgroups completes the proof for fixed m . Repeating the argument in each dimension establishes the last statement in the corollary and completes the proof. \square

Given Proposition 5.15, the proof of Theorem 1.4 for finite-volume manifolds proceeds exactly as in the compact case.

6. EXAMPLES

6.1. Hyperbolic 3-manifolds and link complements.

We begin with a few examples to give a sense of the known possibilities for existence of arithmetic/non-arithmetic hyperbolic 3-manifolds and their geodesic surfaces.

Example 6.1. The figure-eight knot complement contains infinitely many distinct immersed totally geodesic surfaces. They may be taken to be closed. [21, §9.6]

Example 6.2. The Weeks manifold [21, §13.5, Ex. 1] is arithmetic, and contains no totally geodesic surfaces.

Example 6.3. Performing $(-4, 1)$ -surgery on the sister of the figure-eight knot complement [21, §13.5, Ex. 8] determines a non-arithmetic closed hyperbolic manifold M of volume 1.42361... that satisfies the conditions of [21, Thm. 5.3.1], and hence contains no totally geodesic surfaces.

Example 6.4. Consider the three-twist knot 5_2 . It has (invariant) trace field the cubic extension of \mathbb{Q} with minimal polynomial $t^3 - t^2 + 1$. It follows from [21, Thm. 5.3.8] that the associated hyperbolic knot complement contains no *closed* totally geodesic surfaces. It does, however, contain an immersed totally geodesic 3-punctured sphere (see [34]).

Example 6.5. Calegari showed that fibered knots (more generally, fibered knots in rational homology spheres) with trace field of odd degree cannot contain immersed totally geodesic surfaces [8]. For instance, this holds for the knot 8_{20} in [35].

As further motivation, we recall the Menasco–Reid conjecture.

Conjecture 6.6 (Menasco–Reid [26]). *Let K be a hyperbolic knot. Then $S^3 \setminus K$ does not contain a closed, embedded totally geodesic surface.*

In [26], Menasco and Reid prove the conjecture for tunnel number one knots. Our results then have the following ‘almost’ version of the conjecture, which also allows one to promote embedded to immersed:




FIGURE 5. An essential 2-punctured disk from an unknotted component.

Theorem 6.7. *Let K be a hyperbolic knot for which $S^3 \setminus K$ satisfies the conditions of Theorem 1.4. Then $S^3 \setminus K$ contains only finitely many closed, immersed totally geodesic surfaces.*

Menasco and Reid also proved that alternating hyperbolic links cannot contain embedded closed totally geodesic surfaces [26, Thm. 1]. Such a link can certainly contain punctured totally geodesic surfaces, and, when Theorem 1.4 applies, we see that $S^3 \setminus L$ can only contain finitely many such surfaces. To prove Theorem 1.3, we must produce an infinite family of examples to which Theorem 1.4 will apply. We do this via the operation on links called the *belted sum*, which is justified by the following theorem of Adams.

Theorem 6.8 (Adams [2]). *Let M be a complete finite-volume hyperbolic 3-manifold. Then any incompressible and properly embedded 3-punctured sphere in M is isotopic to one that is totally geodesic.*

Let M_1, M_2 be noncommensurable arithmetic hyperbolic 3-manifolds each containing an embedded totally geodesic 3-punctured spheres. Since the hyperbolic structure on a 3-punctured sphere is unique, we can cut M_1 and M_2 open along these surfaces and obtain a Gromov and Piatetski-Shapiro non-arithmetic manifold. The proof of [2, Thm. 4.1] makes this connection very explicit. In particular, Theorem 1.4 applies and the hybrid manifold has finitely many totally geodesic surfaces.

We will apply this in the case where $M_i = S^3 \setminus L_i$ for L_i a link containing an unknotted component as in Figure 5. This contains an obvious properly embedded 2-punctured disk D_i , which is then isotopic to a totally geodesic 3-punctured sphere in M_i . One can then glue $M_1 \setminus D_1$ to $M_2 \setminus D_2$ to obtain a new link $L_{1,2}$ in S^3 , which is visibly built from arithmetic building blocks. This link is the belted sum of L_1 and L_2 .

The belted sum of two noncommensurable arithmetic links will be a non-arithmetic link built out of two dissimilar hyperbolic building blocks. To prove Theorem 1.3, we use an arithmetic link containing two trivial summands as in Figure 5. See Figure 6 for some arithmetic links with this property. We chose these links to be mutually incommensurable. The first is the Whitehead link, which is commensurable with $\mathrm{PSL}_2(\mathbb{Z}[i])$. The second is the three chain link, which is commensurable with $\mathrm{PSL}_2\left(\mathbb{Z}\left[\frac{1+\sqrt{-7}}{2}\right]\right)$. The third is a five component link commensurable with $\mathrm{PSL}_2\left(\mathbb{Z}\left[\frac{1+\sqrt{-15}}{2}\right]\right)$. See [21, §9.2].




FIGURE 6. Three incommensurable arithmetic links.




FIGURE 7. The belted sum of the Whitehead link and the three chain link.

See Figure 7 for the belted sum of the Whitehead link and the three chain link. This is the link 7_6^2 from Rolfsen’s tables [35]. This is a link complement built from dissimilar hyperbolic building blocks, hence it contains only finitely many immersed totally geodesic submanifolds. It is alternating, so it has no closed immersed totally geodesic submanifolds by [26, Thm. 1]. We note that it has invariant trace field $\mathbb{Q}(i, \sqrt{-7})$ by a theorem of Neumann–Reid (see [21, §5.6]). However, to prove Theorem 1.3, we use the five component link in Figure 6.

Proof of Theorem 1.3. Let L_1 be the Whitehead link and L_2 the five component link in Figure 6. Then L_2 contains a pair of totally geodesic 3-punctured spheres S_1 and S_2 associated with the, say, the red and magenta unknotted components.

Let N_0^+ and N_0^- be $S^3 \setminus L_1$ cut open along the totally geodesic 3-punctured sphere S_0 associated with the red unknotted component. For $1 \leq i \leq r$, let N_i be a copy of $S^3 \setminus L_2$, cut open along S_1 and S_2 . We can then perform an iterated belted sum, starting with N_0^+ belted to N_1 along S_1 , N_i belted to N_{i+1} by gluing S_1 to S_2 ($2 \leq i \leq r-1$), then N_r belted to N_0^- along S_2 .


$$N_0^+ \xrightarrow{S_0 \leftrightarrow S_1} N_1 \xrightarrow{S_2 \leftrightarrow S_1} N_2 \cdots N_{r-1} \xrightarrow{S_2 \leftrightarrow S_1} N_r \xrightarrow{S_2 \leftrightarrow S_0} N_0^-$$

This determines a link \mathcal{L}_r in S^3 that is built from arithmetic building blocks, and there are two adjacent blocks that are dissimilar, so $S^3 \setminus \mathcal{L}_r$ contains only finitely many immersed totally geodesic submanifolds.

To complete the proof of the theorem, we must make sure that the links \mathcal{L}_r are actually distinct. However, this follows immediately from the volume calculation

$$\text{vol}(S^3 \setminus \mathcal{L}_r) = 2 \text{vol}(S^3 \setminus L_1) + r \text{vol}(S^3 \setminus L_2)$$

found in [2, Cor. 5.2]. Mostow rigidity implies that hyperbolic volume is a link invariant, so this proves the theorem. \square

FIGURE 8. The Coxeter diagram of a 5-simplex in \mathbb{H}^5 .

6.2. Hyperbolic Coxeter lattices . A Coxeter polyhedron $P \subset \mathbb{H}^n$ is a finite-volume polyhedron with totally geodesic faces having the property that adjacent faces intersect at angles which are integral submultiples of π . Such finite-volume polyhedrons can only exist in dimensions $n \leq 995$ by Prokhorov [29], and compact ones can only exist in dimensions $n \leq 29$ by Vinberg [40]. Associated with such a polyhedron P , one can form the lattice $\Gamma_P < \text{SO}(n, 1)$ generated by reflections in the hyperplanes containing the faces of the polyhedron. We call these *Coxeter lattices*. Vinberg gave a simple criterion for whether such a lattice is arithmetic, and gave examples in dimensions $n = 3, 4, 5$ of non-arithmetic Coxeter lattices that pre-dated the Gromov–Piatetski-Shapiro constructions. This was extended by Ruzmanov [36] who produced non-arithmetic Coxeter lattices in dimension $6 \leq n \leq 10$.

In a recent paper, Vinberg [41] points out that the Gromov–Piatetski-Shapiro construction can also be applied to Coxeter polyhedra in a particularly simple manner. Given a pair of polyhedra P_1, P_2 , let us assume that they contain faces $F_i \subset P_i$ which are isometric to each other, and have the property that every other face which intersects F_i does so orthogonally. In that case, we can form a new Coxeter polyhedron P by gluing the P_i together along the F_i . If the original polyhedra P_i are not commensurable, then the resulting P is non-arithmetic. Vinberg used this procedure to create non-arithmetic Coxeter lattices in dimensions $n = 11, 12, 14, 18$ (as well as new examples in some lower dimensions).

Let us call a Coxeter polyhedron P *splittable* if one can find a totally geodesic hyperplane $H \subset \mathbb{H}^n$ with the property that every face of P which intersects H either is contained in H or intersects H orthogonally. Observe that the polyhedra P_i in Vinberg’s gluing construction are splittable, as is the resulting polyhedron P . Splittable polyhedra also play a key role in work of Allcock [4]. We call a polyhedron *unsplittable* if it is not splittable.

Lemma 6.9. *Let P be an unsplittable Coxeter polyhedron. Then the associated Coxeter lattice Γ_P is not commensurable with any Gromov–Piatetski-Shapiro type lattice.*

Proof. By way of contradiction, assume that Γ_P is commensurable with a Gromov–Piatetski-Shapiro type lattice Λ . Then there are isomorphic finite index torsion-free subgroups $\Gamma < \Gamma_P$ and $\Gamma < \Lambda$. The associated hyperbolic manifold $M = \mathbb{H}^n/\Gamma$ is then itself a Gromov–Piatetski-Shapiro type hyperbolic manifold, but is also tessellated by finitely many copies of the hyperbolic polyhedron P . Let $\Sigma \subset M$ be a totally geodesic cutting hypersurface for the Gromov–Piatetski-Shapiro decomposition of M .

In the tessellation of M , consider any of the copies of P that intersect Σ . Since P is unsplittable, and Σ is codimension one, there exists a face of this polyhedron that intersects Σ at an angle $0 < \theta < \pi/2$. But this face extends to a closed totally geodesic immersed hypersurface, which contradicts angle rigidity. \square




FIGURE 9. Coxeter diagrams for 3-simplices in \mathbb{H}^3 .

We now produce examples in dimension 3 and 5 to which Lemma 6.9 applies. First we briefly recall Vinberg’s notion of a *quasi-arithmetic* lattice. Let \mathcal{G} be a \mathbb{Q} -algebraic group such that $\mathcal{G}(\mathbb{Z})$ defines an arithmetic lattice in $\mathrm{SO}(n, 1)$. A lattice Γ in $\mathrm{SO}(n, 1)$ is quasi-arithmetic if it is contained in $\mathcal{G}(\mathbb{Q})$. Vinberg showed that there are quasi-arithmetic lattices that are not arithmetic (i.e., are not commensurable with $\mathcal{G}(\mathbb{Z})$) [39]. The inbreeding examples of Agol and Belolipetsky–Thomson are also quasi-arithmetic, but the Gromov–Piatetski-Shapiro type examples explained in §2.2 are not (see [38]).

The 5-dimensional example in the following will prove Theorem 1.7, i.e., that there are non-arithmetic lattices in $\mathrm{SO}(5, 1)$ that are not commensurable with a lattice constructed by the methods of Gromov–Piatetski-Shapiro or Agol.

Example 6.10. It is easy to see that if the Coxeter polyhedron $P \subset \mathbb{H}^n$ is combinatorially an n -simplex, then P is unsplittable. There are 72 finite volume Coxeter polyhedra with this combinatorial type. Of these, a few examples give rise to non-arithmetic lattices: one in dimension = 5 and seven in dimension = 3 (see [19, p. 128]). The 5-dimensional example arises from a non-compact, finite-volume, 5-simplex in \mathbb{H}^5 with Coxeter diagram given in Figure 8. For the 3-dimensional examples, six of the seven examples are non-compact, finite volume, while one is compact. Their Coxeter diagrams are listed out in Figure 9, with the upper left-most diagram corresponding to the compact 3-simplex polyhedron. For all these examples, our Lemma 6.9 applies and the associated lattices are not commensurable with any Gromov–Piatetski-Shapiro type lattices. Moreover, for the non-compact polyhedra, the associated lattices are not quasi-arithmetic (see [39, p. 442, Remark 3]). This proves Theorem 1.7.

Given a splittable Coxeter polyhedron $P \subset \mathbb{H}^n$, one can consider the hyperplane $H \subset \mathbb{H}^n$ giving rise to the splitting. Then the intersection $H \cap P$ yields a Coxeter polyhedron in $H \cong \mathbb{H}^{n-1}$ with faces of $H \cap P$ corresponding bijectively with the faces of the original P that were orthogonal to H . The number of faces of $H \cap P$ is strictly smaller than the number of faces of P . Indeed, H separates \mathbb{H}^n into two half-spaces H^+ and H^- , and in order for P to have finite volume, there needs to be at least one face of P on each side of H satisfying $H^\pm \cap \mathrm{Int}(P) \neq \emptyset$. Such faces will be disjoint from H , giving a generator for Γ_P which does not lie in $\Gamma_{P \cap H}$. Recall

that a *special subgroup* of a Coxeter group is a subgroup generated by a subset of the Coxeter generating set. This establishes the following:

Lemma 6.11. *If a Coxeter polyhedron $P \in \mathbb{H}^n$ is splittable, then the associated Coxeter group contains a proper special subgroup which is itself a Coxeter group for a Coxeter polyhedron $P' \subset \mathbb{H}^{n-1}$.*

In practice, Lemma 6.11 can be used to show that certain Coxeter polyhedron are unsplittable. For example, Andreev’s theorem gives an algorithm for deciding whether a Coxeter group arises from a polyhedron in \mathbb{H}^3 (see [12, §6.10]). Thus if one is given a Coxeter polyhedron $P \subset \mathbb{H}^4$, one can sometimes use Andreev’s theorem to algorithmically verify if the polyhedron is unsplittable.

Closely related to our unsplittable polyhedra are the *essential* polyhedra, introduced by Felikson and Tumarkin [15]. These are polyhedra which cannot be decomposed into smaller Coxeter polyhedra glued along faces. In their paper they give a commensurability classification of all the known (at the time) essential polyhedra. While the two notions of unsplittable and essential are related, neither implies the other.

We close by noting that it would be interesting to use results of this kind to produce infinitely many commensurability classes of lattices in $\mathrm{SO}(n, 1)$ for small n that cannot arise from the constructions of Gromov–Piatetski-Shapiro or Agol. While this seems possible in small dimensions, one still wonders if the “maybe for large n ” in Question 1.6 is related to the non-existence of Coxeter polytopes and related geometric objects in sufficiently high dimension.

REFERENCES

- [1] M. Abert, N. Bergeron, I. Biringer, T. Gelander, N. Nikolov, J. Raimbault, and I. Samet. On the growth of L^2 -invariants of locally symmetric spaces, II: exotic invariant random subgroups in rank one. arXiv:1612.09510.
- [2] C. C. Adams. Thrice-punctured spheres in hyperbolic 3-manifolds. *Trans. Amer. Math. Soc.*, 287(2):645–656, 1985.
- [3] I. Agol. Systoles of hyperbolic 4-manifolds. arXiv: 0612290.
- [4] D. Allcock. Infinitely many hyperbolic Coxeter groups through dimension 19. *Geom. Topol.*, 10:737–758, 2006.
- [5] M. V. Belolipetsky and S. A. Thomson. Systoles of hyperbolic manifolds. *Algebr. Geom. Topol.*, 11(3):1455–1469, 2011.
- [6] Y. Benoist and H. Oh. Geodesic planes in geometrically finite acylindrical 3-manifolds. *Preprint*, 2018.
- [7] A. Borel. Density and maximality of arithmetic subgroups. *J. Reine Angew. Math.*, 224:78–89, 1966.
- [8] D. Calegari. Real places and torus bundles. *Geom. Dedicata*, 118:209–227, 2006.
- [9] J. W. Cannon. A positional characterization of the $(n - 1)$ -dimensional Sierpiński curve in S^n ($n \neq 4$). *Fund. Math.*, 79(2):107–112, 1973.
- [10] S. G. Dani and G. A. Margulis. Values of quadratic forms at primitive integral points. *Invent. Math.*, 98(2):405–424, 1989.
- [11] S. G. Dani and G. A. Margulis. Limit distributions of orbits of unipotent flows and values of quadratic forms. In *I. M. Gel’fand Seminar*, volume 16 of *Adv. Soviet Math.*, pages 91–137. Amer. Math. Soc., 1993.
- [12] M. W. Davis. *The geometry and topology of Coxeter groups*, volume 32 of *London Mathematical Society Monographs Series*. Princeton University Press, 2008.
- [13] K. Delp, D. Hoffoss, and J. Manning. Problems in Groups, Geometry, and Three-Manifolds. arXiv:1512.04620.
- [14] M. Einsiedler and P. Wirth. Effective equidistribution of closed hyperbolic surfaces on congruence quotients of hyperbolic spaces. *In preparation*, 2018.

- [15] A. Felikson and P. Tumarkin. Essential hyperbolic Coxeter polytopes. *Israel J. Math.*, 199(1):113–161, 2014.
- [16] T. Gelander and A. Levit. Counting commensurability classes of hyperbolic manifolds. *Geom. Funct. Anal.*, 24(5):1431–1447, 2014.
- [17] M. Gromov. Hyperbolic groups. In *Essays in group theory*, volume 8 of *Math. Sci. Res. Inst. Publ.*, pages 75–263. Springer, New York, 1987.
- [18] M. Gromov and I. Piatetski-Shapiro. Nonarithmetic groups in lobachevsky spaces. *Inst. Hautes Études Sci. Publ. Math.*, (66):93–103, 1988.
- [19] N. W. Johnson, R. Kellerhals, J. G. Ratcliffe, and S. T. Tschantz. Commensurability classes of hyperbolic Coxeter groups. *Linear Algebra Appl.*, 345:119–147, 2002.
- [20] C. Maclachlan and A. W. Reid. Commensurability classes of arithmetic Kleinian groups and their Fuchsian subgroups. *Math. Proc. Cambridge Philos. Soc.*, 102(2):251–257, 1987.
- [21] C. Maclachlan and A. W. Reid. *The arithmetic of hyperbolic 3-manifolds*, volume 219 of *Graduate Texts in Mathematics*. Springer-Verlag, 2003.
- [22] A. Marden. *Outer circles*. Cambridge University Press, 2007. An introduction to hyperbolic 3-manifolds.
- [23] G. A. Margulis. Formes quadratiques indéfinies et flots unipotents sur les espaces homogènes. *C. R. Acad. Sci. Paris Sér. I Math.*, 304(10):249–253, 1987.
- [24] G. A. Margulis. *Discrete subgroups of semisimple Lie groups*, volume 17 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*. Springer-Verlag, 1991.
- [25] C. T. McMullen, A. Mohammadi, and H. Oh. Geodesic planes in hyperbolic 3-manifolds. *Invent. Math.*, 209(2):425–461, 2017.
- [26] W. Menasco and A. W. Reid. Totally geodesic surfaces in hyperbolic link complements. In *Topology '90 (Columbus, OH, 1990)*, volume 1 of *Ohio State Univ. Math. Res. Inst. Publ.*, pages 215–226. de Gruyter, 1992.
- [27] J. S. Meyer. Totally geodesic spectra of arithmetic hyperbolic spaces. *Trans. Amer. Math. Soc.*, 369(11):7549–7588, 2017.
- [28] J. J. Millson. On the first Betti number of a constant negatively curved manifold. *Ann. of Math. (2)*, 104(2):235–247, 1976.
- [29] M. N. Prokhorov. Absence of discrete groups of reflections with a noncompact fundamental polyhedron of finite volume in a Lobachevskii space of high dimension. *Izv. Akad. Nauk SSSR Ser. Mat.*, 50(2):413–424, 1986.
- [30] J. Rambault. A note on maximal lattice growth in $SO(1, n)$. *Int. Math. Res. Not. IMRN*, (16):3722–3731, 2013.
- [31] J. G. Ratcliffe. *Foundations of hyperbolic manifolds*, volume 149 of *Graduate Texts in Mathematics*. Springer-Verlag, 1994.
- [32] M. Ratner. On Raghunathan’s measure conjecture. *Ann. of Math. (2)*, 134(3):545–607, 1991.
- [33] M. Ratner. Raghunathan’s topological conjecture and distributions of unipotent flows. *Duke Math. J.*, 63(1):235–280, 1991.
- [34] A. W. Reid. Totally geodesic surfaces in hyperbolic 3-manifolds. *Proc. Edinburgh Math. Soc. (2)*, 34(1):77–88, 1991.
- [35] D. Rolfsen. *Knots and links*, volume 7 of *Mathematics Lecture Series*. Publish or Perish, 1990.
- [36] O. P. Ruzmanov. Examples of nonarithmetic crystallographic Coxeter groups in n -dimensional Lobachevskii space when $6 \leq n \leq 10$. In *Problems in group theory and in homological algebra (Russian)*, pages 138–142. Yaroslav. Gos. Univ., Yaroslavl’, 1989.
- [37] N. A. Shah. Closures of totally geodesic immersions in manifolds of constant negative curvature. In *Group theory from a geometrical viewpoint (Trieste, 1990)*, pages 718–732. World Sci. Publ., 1991.
- [38] S. Thomson. Quasi-arithmeticity of lattices in $PO(n, 1)$. *Geom. Dedicata*, 180:85–94, 2016.
- [39] E. B. Vinberg. Discrete groups generated by reflections in Lobachevskii spaces. *Mat. Sb. (N.S.)*, 72 (114):471–488; correction, *ibid.* 73 (115) (1967), 303, 1967.
- [40] E. B. Vinberg. The nonexistence of crystallographic reflection groups in Lobachevskii spaces of large dimension. *Funktsional. Anal. i Prilozhen.*, 15(2):67–68, 1981.
- [41] E. B. Vinberg. Non-arithmetic hyperbolic reflection groups in higher dimensions. *Mosc. Math. J.*, 15(3):593–602, 606, 2015.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405
E-mail address: `fisherdm@indiana.edu`, `nimimill@iu.edu`

DEPARTMENT OF MATHEMATICS, OHIO STATE UNIVERSITY, COLUMBUS, OH 43210
E-mail address: `jlafont@math.ohio-state.edu`

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA, PA 19122
E-mail address: `mstover@temple.edu`