

Isomorphism versus commensurability for a class of finitely presented groups

Goulnara Arzhantseva, Jean-François Lafont and Ashot Minasyan

Communicated by Alexander Olshanskii

Abstract. We construct a class of finitely presented groups within which the isomorphism problem is solvable but the commensurability problem is unsolvable. Conversely, we construct a class of finitely presented groups within which the commensurability problem is solvable but the isomorphism problem is unsolvable. These are the first examples of such a contrastive complexity behavior with respect to the isomorphism problem.

1 Introduction

The purpose of this paper is to study the relative algorithm complexities of the following two major group theoretical decision problems: the isomorphism problem and the commensurability problem.

Both of these problems have a long history [12, 27], a meaningful topological interpretation [5, 28], and a number of famous solutions for specific classes of groups [13, 20, 21]. It is also well known that these problems are undecidable within the class of all finitely presented groups. However their comparison from the algorithmic point of view seems not to have been done up to now. Moreover, there have been numerous results comparing decision problems dealing with elements in a single group, such as the word problem, conjugacy problem, power problem, etc. (see, for instance, [18, 21]). In contrast, there have so far been no comparative results involving the isomorphism problem. We remedy this situation, by establishing the following two complementary theorems:

Theorem 1.1. *There exists a recursively enumerable class \mathcal{C}_1 of finite presentations of groups, with uniformly solvable word problem, such that the isomorphism problem is solvable but the commensurability problem is unsolvable within this class.*

The research of the first author was partially supported by the ERC grant ANALYTIC no. 259527, and by the Swiss NSF, under Sinergia grant CRSI22-130435. The second author was partially supported by the NSF, under grants DMS-0906483, DMS-1207782, and by an Alfred P. Sloan research fellowship. The work of the third author was supported by the EPSRC grant EP/H032428/1.

Theorem 1.2. *There exists a recursively enumerable class \mathcal{C}_2 of finite presentations of groups such that the commensurability problem is solvable but the isomorphism problem is unsolvable within this class.*

These results can be contrasted with Thomas [31, Theorem 1.1], who showed that the isomorphism and commensurability problems have the same complexity from the viewpoint of descriptive set theory.

Let us now explain the terminology and the meaning of our main theorems.

A class \mathcal{C} of finite presentations of groups has *uniformly solvable word problem* if there is an algorithm which takes as input a presentation $P \in \mathcal{C}$ and a word in the generators of this presentation, and decides whether or not this word represents the identity element of the group given by P .

Two groups G_1, G_2 are *commensurable* if there exist two subgroups of finite index $H_i \leq G_i$ for $i = 1, 2$, such that H_1 and H_2 are isomorphic. It is not difficult to see that commensurability is an equivalence relation.

Given a class \mathcal{C} of finite presentations of groups, we say that the *isomorphism problem is solvable within \mathcal{C}* (*commensurability problem is solvable within \mathcal{C}*) if there is an algorithm, taking as input two group presentations from \mathcal{C} and deciding whether or not these presentations define isomorphic (commensurable) groups.

Often, when considering the isomorphism problem, one is looking at a certain class \mathcal{G} of finitely presented *groups*. This actually means the class of all finite presentations of groups from \mathcal{G} . At first glance, it might seem that our Theorems 1.1 and 1.2 are somewhat more restrictive, as we are only picking out some specific family of presentations. Let us clarify this issue.

Let \mathcal{G}_1 denote the collection of all groups defined via the presentations in the class \mathcal{C}_1 appearing in Theorem 1.1, and let $\widehat{\mathcal{C}}_1$ denote the class of *all* finite presentations of groups from \mathcal{G}_1 (so clearly $\mathcal{C}_1 \subset \widehat{\mathcal{C}}_1$). It follows immediately from Theorem 1.1 that the commensurability problem is unsolvable for the class $\widehat{\mathcal{C}}_1$ of finite presentations of groups, as it is already unsolvable within the subclass \mathcal{C}_1 . On the other hand, the isomorphism problem is still solvable within the class $\widehat{\mathcal{C}}_1$. Indeed, given any presentation $P \in \widehat{\mathcal{C}}_1$, one can start applying Tietze transformations to it; simultaneously we can start writing down the finite presentations from \mathcal{C}_1 , because the class \mathcal{C}_1 is recursively enumerable. At each step we can compare the transformations of P , obtained so far, with the presentations from the class \mathcal{C}_1 , written down by this step. After finitely many steps we will find a finite presentation $P' \in \mathcal{C}_1$ which defines the same group (up to isomorphism) as P (see [19, II.2.1]). This easily yields an algorithm that identifies a pair of presentations from \mathcal{C}_1 which define the same groups as the given pair of presentations in $\widehat{\mathcal{C}}_1$. Taking the resulting pair of presentations in \mathcal{C}_1 , we can then apply the algorithm for deciding the isomorphism problem within the subclass \mathcal{C}_1 .

As such, we view Theorem 1.1 as a statement about the corresponding class of groups \mathcal{G}_1 .

Similarly, let \mathcal{G}_2 denote the collection of all groups defined via the presentations in the class \mathcal{C}_2 appearing in Theorem 1.2, and let $\widehat{\mathcal{C}}_2$ denote the class of *all* finite presentations of groups from \mathcal{G}_2 (so again, we have $\mathcal{C}_2 \subset \widehat{\mathcal{C}}_2$). By an identical argument to the one in the previous paragraph, we have that the isomorphism problem is unsolvable in the class $\widehat{\mathcal{C}}_2$, but the commensurability problem is solvable. This allows us to view Theorem 1.2 as a statement about the corresponding class of groups \mathcal{G}_2 .

The fact that the isomorphism problem is unsolvable within the class \mathcal{C}_2 implies, in particular, that there are infinitely many pairwise non-isomorphic groups within \mathcal{G}_2 . More precisely, the set of representatives of isomorphism classes of groups from \mathcal{G}_2 is not recursively enumerable. This fact is of particular interest because it cannot be seen directly from our construction of the class \mathcal{C}_2 below.

The proofs of both theorems rely on a combination of various embedding theorems from combinatorial and geometric group theory involving finitely presented infinite simple groups and infinite groups with no finite quotients. The main idea is to start with a single group G and construct a class \mathcal{K} , of mapping tori of G , for which the isomorphism problem is directly related to the word problem in G . Similarly, the commensurability problem in \mathcal{K} will be directly related to the torsion problem in G . Thus the solvability/unsolvability of the word (resp. torsion) problem in G will yield the same for the isomorphism (resp. commensurability) problem in \mathcal{K} .

In Section 5 we prove that there exist no recursive classes of groups with decidable isomorphism or commensurability problems. This shows that the statements of Theorems 1.1 and 1.2 are optimal, as the recursively enumerable classes of groups we construct in these theorems cannot be recursive.

Besides isomorphism and commensurability, there are other natural equivalence relations on the class of finitely presented groups such as virtual isomorphism, bi-Lipschitz equivalence, quasi-isometry, etc. We discuss the corresponding algorithmic problems in the last section, where we also state some open questions.

2 Mapping tori of groups without proper finite index subgroups

Let G be a group and $\varphi \in \text{Aut}(G)$ be an automorphism of G . Let $G_\varphi := G \rtimes_\varphi \mathbb{Z}$ denote the associated *mapping torus*. As a set, $G_\varphi = G \times \mathbb{Z}$ and the group product is defined by $(g, n)(g', m) := (g \cdot \varphi^n(g'), n + m)$, where φ^n denotes the automorphism of G which is the n -fold composition of φ , with the convention that $\varphi^0 = \text{id}_G$, where $\text{id}_G \in \text{Aut}(G)$ the identity automorphism of G .

We shall consider the class of groups $\mathcal{K}_{G, \Phi} = \{G_\varphi \mid \varphi \in \Phi\}$, where Φ is some subset of $\text{Aut}(G)$, and analyze the isomorphism problem within the corresponding class of group presentations. We denote by $\bar{\varphi}$ the image of φ under the canonical epimorphism $\text{Aut}(G) \twoheadrightarrow \text{Out}(G) := \text{Aut}(G)/\text{Inn}(G)$ onto the quotient of $\text{Aut}(G)$ by the subgroup consisting of inner automorphisms.

Proposition 2.1. *Suppose that G is a group which has no epimorphisms onto \mathbb{Z} , and $\varphi, \psi \in \text{Aut}(G)$. Then the following are equivalent.*

- (i) G_φ is isomorphic to G_ψ .
- (ii) $\bar{\varphi} \in \text{Out}(G)$ is conjugate to one of the two elements $\bar{\psi}, (\bar{\psi})^{-1} \in \text{Out}(G)$.

Proof. Suppose that G_φ is isomorphic to G_ψ via an isomorphism

$$\rho: G \rtimes_\varphi \mathbb{Z} \rightarrow G \rtimes_\psi \mathbb{Z}.$$

Let $\tau: G \rightarrow \mathbb{Z}$ be the homomorphism defined by the composition

$$G \hookrightarrow G \rtimes_\varphi \mathbb{Z} \xrightarrow{\rho} G \rtimes_\psi \mathbb{Z} \twoheadrightarrow \mathbb{Z},$$

with the natural inclusion and epimorphism maps. It follows that τ is trivial as by hypothesis we know that G does not map onto \mathbb{Z} . Therefore, the restriction of ρ to G has image entirely contained in $G \leq G_\psi$. Applying the same argument to ρ^{-1} and recalling that $\rho^{-1} \circ \rho = \text{id}_{G_\varphi}$, we can conclude that ρ maps the G -factor in G_φ isomorphically onto the G -factor in G_ψ .

On the other hand, a generator t of the \mathbb{Z} -factor in $G_\varphi = G \rtimes_\varphi \mathbb{Z}$ has to map to a generator under the composition

$$\langle t \rangle = \mathbb{Z} \hookrightarrow G \rtimes_\varphi \mathbb{Z} \xrightarrow{\rho} G \rtimes_\psi \mathbb{Z} \twoheadrightarrow \mathbb{Z}.$$

Indeed, the composition map $G_\varphi \xrightarrow{\rho} G_\psi \twoheadrightarrow \mathbb{Z}$ is surjective as ρ is an isomorphism. Since $G \leq G_\varphi$ is contained in the kernel of this map, the image is determined by the image of the quotient group G_φ/G . However, such an image coincides with $\langle t \rangle$ through the short exact sequence $\{1\} \rightarrow G \hookrightarrow G_\varphi \twoheadrightarrow \mathbb{Z} \rightarrow \{1\}$. Thus, the surjectivity of $G_\varphi \xrightarrow{\rho} G_\psi \twoheadrightarrow \mathbb{Z}$ implies that $t \in G_\varphi$ maps to a generator $s^{\pm 1}$ of the \mathbb{Z} -factor in G_ψ .

Thus, in terms of the splittings, the isomorphism ρ is of the form:

$$\begin{aligned} (x, 0) &\xrightarrow{\rho} (\alpha(x), 0), \\ (e, 1) &\xrightarrow{\rho} (g, \pm 1) \end{aligned}$$

for any $x \in G$ and some fixed $\alpha \in \text{Aut}(G)$ and $g \in G$ ($e \in G$ is the identity element).

Let us now focus on the case where $(e, 1) \xrightarrow{\rho} (g, 1)$. Since the map ρ is assumed to be an isomorphism, it must preserve the relations of the group G_φ . Evaluating ρ on the relation $(e, 1)(x, 0)(e, 1)^{-1} = (\varphi(x), 0)$ yields the required constraint on the automorphisms. Indeed, evaluating the left hand side, we obtain

$$\rho((e, 1)(x, 0)(e, 1)^{-1}) = (g, 1)(\alpha(x), 0)(\psi^{-1}(g^{-1}), -1) = (g\psi(\alpha(x))g^{-1}, 0),$$

while evaluating the right hand side, we obtain

$$\rho((\varphi(x), 0)) = (\alpha(\varphi(x)), 0).$$

We deduce that the automorphism $\alpha \in \text{Aut}(G)$ and the element $g \in G$ are related to the given automorphisms $\varphi, \psi \in \text{Aut}(G)$ as follows:

$$\alpha \circ \varphi = c_g \circ \psi \circ \alpha,$$

where $c_g \in \text{Aut}(G)$ is the inner automorphism defined by $c_g(y) = gyg^{-1}$ for all $y \in G$.

Passing to the outer automorphism group, we see that we have to have

$$\bar{\alpha} \cdot \bar{\varphi} = \bar{\psi} \cdot \bar{\alpha},$$

that is, the classes $\bar{\varphi}$ and $\bar{\psi}$ are conjugate in $\text{Out}(G)$.

Conversely, if the classes $\bar{\varphi}, \bar{\psi} \in \text{Out}(G)$ are conjugate by some $\bar{\alpha} \in \text{Out}(G)$, then one can find an element $g \in G$ so that $\alpha \circ \varphi = c_g \circ \psi \circ \alpha$. It is now immediate that $G_\varphi \cong G_\psi$, via the isomorphism map defined by $(x, 0) \mapsto (\alpha(x), 0)$ and $(e, 1) \mapsto (g, 1)$.

A similar analysis can be done in the case $(e, 1) \xrightarrow{\rho} (g, -1)$. This yields the relation

$$\alpha \circ \varphi = c_g \circ \psi^{-1} \circ \alpha,$$

that is, the classes $\bar{\varphi}$ and $(\bar{\psi})^{-1}$ are conjugate in $\text{Out}(G)$. This finishes the proof. \square

In order to facilitate the notation let us give the following

Definition 2.2. We will say that a group G is NFQ (“No Finite Quotients”) if the only finite quotient of G is the trivial group.

Since every finite index subgroup contains a finite index normal subgroup, a group G is NFQ if and only if G has no proper subgroups of finite index. It is easy to see that any NFQ group G has no epimorphisms onto \mathbb{Z} , and thus it satisfies the assumptions of Proposition 2.1. Basic examples of NFQ groups are infinite simple groups.

To study the commensurability problem within the class $\mathcal{K}_{G,\Phi}$, we need to know the structure of subgroups of finite index in the corresponding mapping tori. The following observation shows that all such subgroups are “congruence subgroups”:

Proposition 2.3. *Let G be an NFQ group, and $\varphi \in \text{Aut}(G)$. Let $\pi: G_\varphi \twoheadrightarrow \mathbb{Z}$ be the canonical projection onto the \mathbb{Z} -factor of the mapping torus. Assume that $H \leq G_\varphi$ is a finite index subgroup of G_φ . Then $H = \pi^{-1}(k\mathbb{Z}) \cong G_{\varphi^k}$, where k is the index of H in G_φ (and in particular, H must be normal in G_φ).*

Proof. By the assumptions, $[G_\varphi : H] < \infty$, hence $[G : (G \cap H)] < \infty$, therefore $\ker \pi = G \leq H$ as G is NFQ. This forces H to be of the form $\pi^{-1}(k\mathbb{Z})$ for some k . The value of k can then be easily deduced:

$$k = [\mathbb{Z} : k\mathbb{Z}] = [G_\varphi : \pi^{-1}(k\mathbb{Z})] = [G_\varphi : H],$$

as stated in the proposition. □

Combining Propositions 2.1 and 2.3, we immediately obtain

Corollary 2.4. *Let G be an NFQ group, and $\varphi \in \text{Aut}(G)$. Then G_φ is commensurable with $G_{\text{id}_G} \cong G \times \mathbb{Z}$ if and only if the element $\bar{\varphi} \in \text{Out}(G)$ has finite order.*

Given two groups A and B , consider their free product $G = A * B$. For any $a \in A$ we can define a natural automorphism $\tau_a \in \text{Aut}(G)$ by $\tau_a(x) := a^{-1}xa$ for all $x \in A$ and $\tau_a(y) := y$ for all $y \in B$. Note that $(\tau_a)^k = \tau_{a^k}$ in $\text{Aut}(G)$ for all $k \in \mathbb{Z}$.

Lemma 2.5. *Suppose that $B \neq \{1\}$, $a \in A$ and $G = A * B$. Then $\tau_a \in \text{Inn}(G)$ if and only if a belongs to the center of A .*

Proof. Clearly, if a is central in A , then $\tau_a = \text{id}_G \in \text{Inn}(G)$. Conversely, suppose that there is an element $c \in A$ such that $a^{-1}ca \neq c$ in A . Take any $b \in B \setminus \{1\}$ and consider the element $g := cb \in G$. Then $\tau_a(g) = a^{-1}cab$ is not conjugate to g in $G = A * B$ by the criterion for conjugacy in free products (see [19, Section IV.1.4]). Hence $\tau_a \notin \text{Inn}(G)$, as required. □

Since the free product of two NFQ groups is again an NFQ group, we can put together Proposition 2.1, Corollary 2.4 and Lemma 2.5 to achieve

Corollary 2.6. *Let A and B be NFQ groups such that $B \neq \{1\}$ and A has trivial center. Then for $G = A * B$ and any $a \in A$ the following are true:*

- G_{τ_a} is isomorphic to G_{id_G} if and only if $a = 1$ in A .
- G_{τ_a} is commensurable with G_{id_G} if and only if a has finite order in A .

3 Word and torsion problems in NFQ groups

For a finite set X , we use X^* to denote the set of words with letters from $X^{\pm 1}$. Let R be a set of words from X^* and suppose that G is a group given by the presentation $P = \langle X \mid R \rangle$.

For a subset $Z \subseteq X^*$, we say the *word problem for Z in G is solvable* if there is an algorithm, which takes as input a word $w \in Z$ and decides whether or not this word represents the identity element of G . If $Z = X^*$, then the word problem for Z in G is simply known as *the word problem in G* . The word problem is one of the three fundamental group-theoretical decision problems introduced by Max Dehn [12] in 1911 (other two being the conjugacy and the isomorphism problems). It is well known that if the word problem for G is solvable with respect to one finite generating set, then it is solvable with respect to any other finite generating set of G .

For an arbitrary subset $Z \subseteq X^*$, one can also consider the *torsion problem for Z in G* , asking whether there exists an algorithm which inputs a word $w \in Z$, and decides whether or not w represents an element of finite order in G . This is closely related to some decision problems considered by Lipschutz and Miller in [18] (for instance, it is a special case of the *power problem*).

Proposition 3.1. *Every finitely presented group H can be embedded into a finitely presented NFQ group A with trivial center. Moreover, if the word problem in H is solvable, then it is also solvable in A .*

Proof. Take any infinite finitely presented simple group S (for instance, Thompson's group T or V , see [17] or see [6–9] for other such groups) and consider the free product $G = S * S$. Then G is NFQ and hyperbolic relative to these two copies of S . Therefore, by [3, Theorem 1.1], H can be isomorphically embedded into some quotient Q of G . Moreover, from the proof of this theorem, it follows that Q can be obtained from $H * G$ by adding only finitely many defining relations. Consequently, as both H and G are finitely presented, Q will also be finitely presented. The group Q is NFQ as a quotient of the NFQ group G . One can check that the center of the group Q , obtained from [3, Theorem 1.1], is in fact trivial. However, it is easy to bypass this, by setting $A := Q * S$ and observing that A is still finitely presented, NFQ, has trivial center (as a non-trivial free product – see [24, 6.2.6]) and contains a copy of H .

Now, suppose that the word problem in H is solvable. Note that the same is true in S , because the word problem is solvable in any recursively presented simple group ([19, IV.3.6]). By [3, Theorem 1.1] the group Q above is hyperbolic relative to the family of subgroups, consisting of H and two copies of S . Therefore Q has

solvable word problem (see [14, Theorem 3.7] or [22, Corollary 5.5]). Finally, the word problem is solvable in $A = Q * S$ by [19, IV.1.3]. \square

Remark. An alternative proof of Proposition 3.1 could use the theory of small cancellation over free products (see Schupp [25] and [19, Sections V.9, V.10]) instead of [3, Theorem 1.1].

Proposition 3.2. *There exists an NFQ group A_1 , with trivial center and finite presentation $P_1 = \langle X_1 \mid R_1 \rangle$, and a recursively enumerable subset of words $Z_1 = \{z_1, z_2, \dots\} \subset X_1^*$ such that the word problem in A_1 is solvable but the torsion problem for Z_1 in A_1 is unsolvable.*

Proof. Let H_0 be the center-by-metabelian group constructed by P. Hall in [15, p. 435]. Namely, H_0 is generated by two elements a, b , subject to the relations

$$[[b_i, b_j], b_k] = 1, \quad \text{for } i, j, k = 0, \pm 1, \pm 2, \dots,$$

where $b_i := a^{-i} b a^i$, $[x, y] := x^{-1} y^{-1} x y$, and

$$c_{i,j} = c_{i+k,j+k}, \quad \text{for } j > i, \quad i, j, k = 0, \pm 1, \pm 2, \dots,$$

where $c_{i,j} := [b_j, b_i]$. As Hall proved in [15, p. 435], the center of H_0 is the free abelian group with free abelian basis $\{d_1, d_2, \dots\}$, where

$$d_r := c_{0,r} = [a^{-r} b a^r, b], \quad r = 1, 2, \dots$$

Let $\langle a, b \mid R_0 \rangle$ be the above presentation for H_0 . Clearly this presentation is recursive. Now, consider a computable (recursive) function $f: \mathbb{N} \rightarrow \mathbb{N}$ with non-recursive range $f(\mathbb{N}) \subset \mathbb{N}$. Let H_1 be the quotient of H_0 by the central subgroup $\langle d_{f(n)}^n \mid n \in \mathbb{N} \rangle$ where $d_r, r \in \mathbb{N}$, are as above. Then H_1 has the presentation

$$\langle a, b \parallel R_0, ([a^{-f(n)} b a^{f(n)}, b])^n, n \in \mathbb{N} \rangle.$$

The group H_1 will be recursively presented since R_0 is recursively enumerable and f is computable.

By abusing notation, we will continue writing a, b, b_i, d_r for the images in H_1 of the corresponding elements of H_0 . We can solve the word problem in H_1 as follows. Given a word w , over the alphabet $\{a^{\pm 1}, b^{\pm 1}\}$, we want to determine whether $w = 1$ in H_1 . First we compute the sum $\varepsilon_a(w)$ of all exponents of a in w . If $\varepsilon_a(w) \neq 0$, then $w \neq 1$ in H_1 as there is a homomorphism $\alpha: H_1 \rightarrow \langle a \rangle$, whose kernel is generated by $b_i, i \in \mathbb{Z}$, such that $\alpha(w) = a^{\varepsilon_a(w)} \neq 1$. If $\varepsilon_a(w) = 0$, then $w \in B := \langle b_i, i \in \mathbb{Z} \rangle$ and we can re-write w as a word w_1 in letters $b_i, i \in \mathbb{Z}$.

If for some $i \in \mathbb{Z}$, $\varepsilon_{b_i}(w_1) \neq 0$, then, again, $w \neq 1$ in H_1 , because its image will be non-trivial in the abelianization of B . Otherwise, w will represent an element of the center $C := \langle d_r, r \in \mathbb{N} \rangle$ of H_1 , and we can re-write w_1 as a word $w_2 \equiv d_{r_1}^{n_1} d_{r_2}^{n_2} \cdots d_{r_l}^{n_l}$, where $l \geq 0$, $1 \leq r_1 < r_2 < \cdots < r_l$, and $n_j \in \mathbb{Z} \setminus \{0\}$ for $j = 1, \dots, l$. Note that $C = \bigoplus_{r \in \mathbb{N}} \langle d_r \rangle$ by definition. If $l = 0$ then $w = w_2 = 1$ in H_1 . If $l > 0$, then $w_2 = 1$ in C if and only if the order of d_{r_j} in H_1 divides n_j for all $j = 1, 2, \dots, l$. The latter can be verified as follows: for every positive divisor m of n_j , we compute $f(m)$ and check if it is equal to r_j . If this happens for some divisor m of n_j , then the order of d_{r_j} in H_1 is m , by construction, and so $d_{r_j}^{n_j} = 1$. If this is true for all $j = 1, \dots, l$, then $w = w_2 = 1$ in H_1 . As each n_j has only finitely many divisors, this can be checked in finitely many steps. Finally, if there is $j \in \{1, \dots, l\}$ such that for every positive divisor m of n_j , $f(m) \neq r_j$, then the order of d_{r_j} in H_1 does not divide n_j , and hence $w = w_2 \neq 1$ in H_1 .

Thus H_1 is a finitely generated recursively presented group with solvable word problem. By a theorem of Clapham [10, Theorem 6], H_1 can be embedded in a finitely presented group H_2 with solvable word problem. Now we can use Proposition 3.1 to embed H_2 into a finitely presented NFQ group A_1 , with trivial center and solvable word problem. Let $P_1 = \langle X_1 \parallel R_1 \rangle$ be some finite presentation for A_1 . Since $H_1 \leq A_1$, the generators a, b of H_1 can be represented by some words w_1, w_2 (respectively) in the alphabet $X_1^{\pm 1}$, and hence every word in letters from $\{a^{\pm 1}, b^{\pm 1}\}$ can be effectively re-written in letters from $X_1^{\pm 1}$. So, for every $r \in \mathbb{N}$ we can effectively compute a word $z_r \in X_1^*$ representing d_r in A_1 and set $Z_1 := \{z_r \mid r \in \mathbb{N}\} \subset X_1^*$. By construction, Z_1 is recursively enumerable.

Suppose that the torsion problem for Z_1 in A_1 is solvable. Then for any $r \in \mathbb{N}$ we can compute the word $z_r \in Z_1$, representing d_r in A_1 , and check if d_r has finite order in A_1 . But the latter happens if and only if $r \in f(\mathbb{N})$. Thus we would be able to determine whether or not r belongs to the range of f , contradicting the choice of f . Therefore the torsion problem for Z_1 in A_1 is unsolvable and the proposition is proved. □

The next statement suggests a construction which is in some sense opposite to the construction of Proposition 3.2.

Proposition 3.3. *There exists an NFQ group A_2 , with trivial center and finite presentation $P_2 = \langle X_2 \parallel R_2 \rangle$, and a recursively enumerable subset of words $Z_2 = \{z_1, z_2, \dots\} \subset X_2^*$ such that every word from Z_2 represents an element of order at most 2 in A_2 but the word problem for Z_2 in A_2 is unsolvable.*

Proof. Again, let us start with Hall’s group H_0 , used in the proof of Proposition 3.2, keeping the same notation as before. Let $f: \mathbb{N} \rightarrow \mathbb{N}$ be a computable

function with non-recursive range. We now let H_1 be the quotient of H_0 by the central subgroup $\langle d_n^2, d_{f(n)} \mid n \in \mathbb{N} \rangle$.

As before, H_1 will be finitely generated and recursively presented, however, the word problem in H_1 will be unsolvable (since the set $f(\mathbb{N})$ is not recursive). By the celebrated theorem of Higman [16], one can embed H_1 into a finitely presented group H_2 , and applying Proposition 3.1, we can embed H_2 into a finitely presented NFQ group A_2 with trivial center.

Let $P_2 = \langle X_2 \parallel R_2 \rangle$ be some finite presentation for the NFQ group A_2 . Fix some words $w_1, w_2 \in X_2^*$ representing the generators a, b (respectively) of H_1 in A_2 . Clearly there is an algorithm which takes as input a word in the alphabet $\{a^{\pm 1}, b^{\pm 1}\}$ and outputs a corresponding word in the alphabet $X_2^{\pm 1}$ (substituting every a -letter by w_1 and every b -letter by w_2). For each $r \in \mathbb{N}$, let $z_r \in X_2^*$ be the word representing $d_r \in H_1$, obtained this way, and set $Z_2 := \{z_r \mid r \in \mathbb{N}\} \subset X_2^*$. Evidently the set of words Z_2 is recursively enumerable and every word from this set represents an element d_r , which has order at most 2 in A_2 . By construction, $z_r = 1$ in A_2 if and only if $d_r = 1$ in H_1 , which happens if and only if $r \in f(\mathbb{N})$. Since $f(\mathbb{N})$ is non-recursive, we see that the word problem for Z_2 in A_2 is unsolvable. □

4 Proofs of the theorems

We are now ready to establish our two theorems.

Proof of Theorem 1.1. We start with the presentation $P_1 = \langle X_1 \parallel R_1 \rangle$ of the NFQ group A_1 , and the recursively enumerable set of words $Z_1 = \{z_1, z_2, \dots\} \subset X_1^*$, which were constructed in Proposition 3.2. Take some infinite finitely presented simple group B and fix some finite presentation $\langle Y \parallel S \rangle$ of it; recall that the word problem in B is solvable by [19, IV.3.6]. Let $z_0 \in X_1^*$ be the empty word. For each $r \in \mathbb{N} \cup \{0\}$, let d_r denote the element of A_1 represented by the $z_r \in Z_1$; let $G := A_1 * B$ and let C_{r+1} be the cyclic group of order $r + 1$. Then the group $K_r := G_{\tau_{d_r}} \times C_{r+1}$ has the presentation

$$P_{1,r} := \langle X_1, Y, t, u \parallel R_1, S, t^{-1}x^{-1}tz_r^{-1}xz_r, t^{-1}y^{-1}ty, u^{-1}x^{-1}ux, \\ u^{-1}y^{-1}uy, u^{-1}t^{-1}ut, u^{r+1}, \text{ for all } x \in X_1 \text{ and } y \in Y \rangle.$$

Since the sets X_1, Y, R_1 and S are finite, for every $r \in \mathbb{N} \cup \{0\}$, $P_{1,r}$ is a finite presentation of a group. Note that the presentation $P_{1,0}$ defines the group $K_0 \cong G_{\text{id}_G} \cong G \times \mathbb{Z}$.

Now, consider the class of group presentations $\mathcal{C}_1 := \{P_{1,r} \mid r \in \mathbb{N} \cup \{0\}\}$. We can make the following observations.

- (a) the class \mathcal{C}_1 is recursively enumerable by definition.
- (b) the word problem in \mathcal{C}_1 is uniformly solvable. This easily follows from the fact that the word problem in $G = A_1 * B$ is solvable and for each $r \in \mathbb{N} \cup \{0\}$, $G \triangleleft K_r$ and $K_r/G \cong \mathbb{Z} \times C_{r+1}$.
- (c) the isomorphism problem within \mathcal{C}_1 is trivially solvable. This is because for any $r \in \mathbb{N} \cup \{0\}$, the abelianization of the group K_r is isomorphic to $\mathbb{Z} \times C_{r+1}$ (as G is NFQ), hence for any $q \in \mathbb{N} \cup \{0\}$, $q \neq r$, the group K_r is not isomorphic to K_q since their abelianizations have different torsion subgroups. Thus any two distinct presentations from \mathcal{C}_1 define non-isomorphic groups.
- (d) the commensurability problem within \mathcal{C}_1 is unsolvable. Indeed, since the index $[K_r : G_{\tau_{d_r}}] = r + 1$ is finite, the group K_r is commensurable with the group $G_{\tau_{d_r}}$ for each $r \in \mathbb{N} \cup \{0\}$. So, if we could decide whether K_r is commensurable with K_0 , then we would be able to decide whether $G_{\tau_{d_r}}$ is commensurable with G_{id_G} , which, by Corollary 2.6, would imply that the torsion problem for Z_1 in $G = A_1 * B$ is solvable, contradicting to the claim of Proposition 3.2.

Thus the class of group presentations \mathcal{C}_1 satisfies all of the required properties. \square

Proof of Theorem 1.2. Now we start with the presentation $P_2 = \langle X_2 \parallel R_2 \rangle$ of the group A_2 , constructed in Proposition 3.3, and the recursively enumerable set of words $Z_2 = \{z_1, z_2, \dots\} \subset X_2^*$. Take some infinite finitely presented simple group B and fix some finite presentation $\langle Y \parallel S \rangle$ of it; then B will have solvable word problem ([19, IV.3.6]). Let $z_0 \in X_2^*$ be the empty word. For each $r \in \mathbb{N} \cup \{0\}$, let d_r denote the element of A_2 represented by the $z_r \in Z_2$ and let $G := A_2 * B$. Then the group $G_{\tau_{d_r}}$ has the presentation

$$P_{2,r} := \langle X_2, Y, t \parallel R_2, S, t^{-1}x^{-1}tz_r^{-1}xz_r, t^{-1}y^{-1}ty, \text{ for all } x \in X_2, y \in Y \rangle.$$

Since the sets X_2, Y, R_2 and S are finite, for every $r \in \mathbb{N} \cup \{0\}$, $P_{2,r}$ is a finite presentation of a group. As before, the presentation $P_{2,0}$ defines the group $G_{\text{id}_G} \cong G \times \mathbb{Z}$.

For the class of finite presentations $\mathcal{C}_2 := \{P_{2,r} \mid r \in \mathbb{N} \cup \{0\}\}$ we can observe the following:

- (a) the class \mathcal{C}_2 is recursively enumerable by definition.
- (b) the commensurability problem within \mathcal{C}_2 is trivially solvable, because any presentation from this class defines the group $G_{\tau_{d_r}}$, for some $r \in \mathbb{N} \cup \{0\}$, which is commensurable with G_{id_G} by Corollary 2.6, as the element $d_r \in A_2$ has finite order by construction. Thus any two presentations from \mathcal{C}_2 define commensurable groups.

- (c) the isomorphism problem within \mathcal{C}_2 is unsolvable. Indeed, according to Corollary 2.6, for any $r \in \mathbb{N}$, the group $G_{\tau_{d_r}}$, defined by $P_{2,r}$ is isomorphic to G_{id_G} , defined by $P_{2,0}$, if and only if $z_r = 1$ in G . Thus the isomorphism problem within \mathcal{C}_2 is equivalent to the word problem for Z_2 in A_2 , which is unsolvable by construction.

Thus the class \mathcal{C}_2 satisfies all of the needed properties. \square

5 Membership problem for some classes of groups

The purpose of this section is to show that the claims of Theorems 1.1 and 1.2 are optimal; that is, the recursively enumerable classes of groups we construct in these theorems cannot be recursive. More precisely, suppose that we are given a class of finitely presented groups \mathcal{K} , closed under isomorphism. The class \mathcal{K} is said to be *recursive*, or, equivalently, the *membership problem to \mathcal{K} is decidable* (within the class of all finitely presented groups) if there is an algorithm, which takes as input a finite presentation and decides whether or not the group defined by this presentation belongs to \mathcal{K} .

The next statement essentially shows that there is no recursive class of groups satisfying the claim of Theorem 1.1.

Proposition 5.1. *Let \mathcal{K} be a non-empty class of finitely presented groups with solvable isomorphism problem. Then \mathcal{K} is not recursive.*

Proof. Arguing by contradiction, suppose that \mathcal{K} is a non-empty recursive class of finitely presented groups such that the isomorphism problem is solvable within \mathcal{K} . Since \mathcal{K} is non-empty, we can assume that we possess a finite presentation of some group $G \in \mathcal{K}$. Let A be an arbitrary finitely presented group and set $H := G * A$. Then H also finitely presented and a finite presentation of H can be easily obtained from the finite presentations of G and A . Now, by the assumptions, we can decide whether or not H belongs to \mathcal{K} . If $H \notin \mathcal{K}$, then H is not isomorphic to G , hence the group A is non-trivial. If $H \in \mathcal{K}$, then one can apply the algorithm solving the isomorphism problem in \mathcal{K} to decide whether or not $H \cong G$. Evidently, if H is not isomorphic to G , then A is non-trivial. On the other hand, if $H \cong G$, then

$$\begin{aligned} \text{rank}(H) &= \text{rank}(G * A) \\ &= \text{rank}(G) + \text{rank}(A) = \text{rank}(G) \end{aligned}$$

by Grushko–Neumann theorem (see [19, IV.1.9]), where $\text{rank}(G)$ denotes the minimal number of elements required to generate the group G . Hence $\text{rank}(A) = 0$,

i.e., A is the trivial group. Thus we have described an algorithm which decides whether any given finitely presented group A is trivial. But it is well-known that the triviality problem is unsolvable within the class of all finitely presented groups (see, for example, [23, Theorem 2.2]). This contradiction proves the claim of the proposition. \square

Now we state a similar fact for the class of groups appearing in Theorem 1.2.

Proposition 5.2. *Let \mathcal{K} be a non-empty class of finitely presented groups with solvable commensurability problem. Then \mathcal{K} is not recursive.*

The proof of this statement will utilize the following lemma:

Lemma 5.3. *Let G and A be groups and let $H := G * (A * A)$, $F := G \times (A * A)$. Suppose that H and F are both commensurable to G . Then A is the trivial group.*

Proof. First suppose that G is finite. Since F is commensurable to G , it must also be finite, hence $A * A$ is finite, which can only happen if A is trivial.

Assume, now, that G is infinite. If A is non-trivial, then $A * A$ is infinite, hence any finite index subgroup of F itself contains a finite index subgroup that decomposes as a non-trivial direct product (of a finite index subgroup in G with a finite index subgroup in $A * A$). While any finite index subgroup of H decomposes in a non-trivial free product (by Kurosh theorem [19, IV.1.10]), and hence it cannot be isomorphic to a non-trivial direct product (see, for example, [19, Observation, p. 177]). Therefore F cannot be commensurable to H ; this contradicts with the assumption that they are both commensurable to G and the fact that commensurability is a transitive relation. Thus A must be trivial. \square

Proof of Proposition 5.2. As before, assume that there is a recursive class \mathcal{K} with solvable commensurability problem, and let $G \in \mathcal{K}$ be some group with a given finite presentation. Suppose we are given a finite presentation of any group A . Then we can easily produce the finite presentations for the groups $H := G * A * A$ and $F := G \times (A * A)$.

By the assumptions, we can decide whether or not $H \in \mathcal{K}$ and $F \in \mathcal{K}$. If at least one of these groups does not belong to \mathcal{K} , then A is non-trivial. So, assume that both H and F lie in \mathcal{K} . Since the commensurability problem is solvable within \mathcal{K} , we can decide if H and F are commensurable to G . Evidently, if at least one of these groups is not commensurable to G , then A is non-trivial. So, we can further suppose that both H and F are commensurable to G . Lemma 5.3 shows that the latter can happen only if A is trivial. Thus, again, we produced an algorithm deciding the triviality of A , which leads us to the required contradiction. \square

6 Decision problems in geometric group theory

From the viewpoint of geometric group theory, besides the isomorphism and commensurability relations, there are several other equivalence relations on groups which are of natural interest:

- Two finitely generated groups G_1, G_2 are *virtually isomorphic* (sometimes also called *commensurable up to finite kernels*) if there exist a pair of finite index subgroups $H_i \leq G_i$, and some further finite normal subgroups $N_i \trianglelefteq H_i$, $i = 1, 2$, with isomorphic quotients $H_1/N_1 \cong H_2/N_2$.
- Two finitely generated groups G_1, G_2 are *quasi-isometric* if there exists a map $f: G_1 \rightarrow G_2$ and a constant $K > 0$ so that for all $x, y \in G_1$

$$\begin{aligned} \frac{1}{K}d_1(x, y) - K &\leq d_2(f(x), f(y)) \\ &\leq K \cdot d_1(x, y) + K \end{aligned}$$

and the K -neighborhood of $f(G_1)$ is all of G_2 (the d_i are word metrics on the G_i , $i = 1, 2$).

- Two finitely generated groups are *bi-Lipschitz equivalent* if there is a bi-Lipschitz map between (G_1, d_1) and (G_2, d_2) , where again the d_i are word metrics (this is equivalent to the existence of a bijective quasi-isometry between them – see Whyte [33]).

We can now state the corresponding decision problems: the *virtual isomorphism problem* (respectively, *quasi-isometry problem* or *bi-Lipschitz problem*) asks whether there exists an algorithm which, given two finite presentations of groups, can decide whether or not they define virtually isomorphic (resp. quasi-isometric or bi-Lipschitz equivalent) groups. Several of these problems have been studied from the viewpoint of descriptive set theory by Thomas [29–32]. Note that a group is bi-Lipschitz equivalent to the trivial group if and only if it is trivial, and that it is virtually isomorphic, commensurable, or quasi-isometric to the trivial group if and only if it is finite. Since the problem of deciding whether a finitely presented group is finite (or trivial) is unsolvable (this follows from the famous Adian–Rabin theorem, see [1, 2] and [23]), we immediately obtain

Lemma 6.1. *Within the class of all finite presentations of groups, the virtual isomorphism, quasi-isometry, bi-Lipschitz, and commensurability problems are all unsolvable.*

It would be of some interest to study the relative complexity of these various decision problems. A straightforward consequence of the construction appearing in our proof of Theorem 1.2 is the following:

Corollary 6.2. *There exists a recursively enumerable class of finite presentations of groups within which the isomorphism problem is unsolvable, but the virtual isomorphism, quasi-isometry, and bi-Lipschitz problems are all (trivially) solvable.*

Proof. In the notation from the proof of Theorem 1.2, let

$$\Phi := \{\tau_{d_r} \mid r \in \mathbb{N} \cup \{0\}\} \subset \text{Aut}(\mathbf{G}).$$

Then the class of finite presentations \mathcal{C}_2 , constructed in the proof of Theorem 1.2, defines the class

$$\mathcal{K}_{G,\Phi} = \{G_{\tau_{d_r}} \mid r \in \mathbb{N} \cup \{0\}\}$$

of finitely presented groups. As we noticed above, any two groups from this class are commensurable. And since commensurable groups are automatically quasi-isometric, all the groups in $\mathcal{K}_{G,\Phi}$ are quasi-isometric to each other, and the quasi-isometry problem within \mathcal{C}_2 is (trivially) solvable.

Moreover, none of the groups $G_{\tau_{d_r}}$, $r \in \mathbb{N} \cup \{0\}$, can contain a non-trivial finite normal subgroup, for such a subgroup would have to map to the identity under the canonical projection $G_{\tau_{d_r}} \twoheadrightarrow \mathbb{Z}$, and hence it would have to be a normal subgroup in the group $G = A_2 * B$. But a non-trivial free product does not have any non-trivial finite normal subgroups. This tells us that within the class $\mathcal{K}_{G,\Phi}$, two groups are virtually isomorphic if and only if they are commensurable. Therefore the virtual isomorphism problem within \mathcal{C}_2 is also (trivially) solvable.

Finally, noting that $G = A_2 * B$ is non-amenable, as a non-elementary free product, and embeds into every $G_{\tau_{d_r}}$, $r \in \mathbb{N} \cup \{0\}$, we see that all the groups in the class $\mathcal{K}_{G,\Phi}$ are non-amenable. The work of Block and Weinberger [4, Theorem 3.1] implies that the groups in this class all have vanishing 0-dimensional uniformly finite homology. Whyte's thesis [33, Theorem 1.1] then implies that commensurability between any two groups from $\mathcal{K}_{G,\Phi}$ can be promoted to a bi-Lipschitz equivalence. We conclude that all the groups in $\mathcal{K}_{G,\Phi}$ are bi-Lipschitz equivalent to each other, so that the bi-Lipschitz problem within \mathcal{C}_2 is also (trivially) solvable. \square

More generally, we expect that these various decision problems are fundamentally unrelated to each other (with the possible exception of the bi-Lipschitz problem, in view of Whyte's thesis [33]). To be more precise, we suspect that given any two disjoint subsets of these decision problems, one can find a recursively enumerable class of finite presentations of groups such that any problem from the first of these subsets is solvable within this class, while problems from the second subset are all unsolvable.

In another vein, these algorithmic problems are also open for various natural classes of groups. For instance, one could focus on certain classes of lattices

within a fixed semi-simple Lie group G of non-compact type. If the \mathbb{R} -rank of G is ≥ 2 , and one restricts to uniform lattices (so that the quasi-isometry problem is trivially solvable), is the isomorphism problem or commensurability problem solvable? If one focuses on $G = \mathrm{SO}(n, 1)$, $n \geq 4$, and restrict to non-uniform lattices (so that the isomorphism problem is solvable, by Dahmani and Groves [11]), is the commensurability problem (equivalent to the quasi-isometry problem, by Schwartz [26]) solvable or not? Surprisingly, these questions do not seem to have been considered in the literature.

Acknowledgments. The authors thank Cornelia Druțu, Daniel Groves, and Mark Sapir for helpful conversations.

Bibliography

- [1] S. I. Adian, Finitely presented groups and algorithms, *Dokl. Akad. Nauk SSSR* **117** (1957) 9–12.
- [2] S. I. Adian, The unsolvability of certain algorithmic problems in the theory of groups, *Tr. Mosk. Mat. Obs.* **6** (1957), 231–298.
- [3] G. Arzhantseva, A. Minasyan and D. Osin, The SQ-universality and residual properties of relatively hyperbolic groups, *J. Algebra* **315** (2007), 165–177.
- [4] J. Block and S. Weinberger, Aperiodic tilings, positive scalar curvature and amenability of spaces, *J. Amer. Math. Soc.* **5** (1992), 907–918.
- [5] A. Borel, Commensurability classes and volumes of hyperbolic 3-manifolds, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (4)* **8** (1981), no. 1, 1–33.
- [6] M. Burger and S. Mozes, Finitely presented simple groups and products of trees, *C. R. Acad. Sci. Paris Sér. I Math.* **324** (1997), 747–752.
- [7] M. Burger and S. Mozes, Lattices in product of trees, *Publ. Math. Inst. Hautes Études Sci.* **92** (2000), 151–194.
- [8] P.-E. Caprace and B. Remy, Simplicité abstraite des groupes de Kac-Moody non affines, *C. R. Math. Acad. Sci. Paris* **342** (2006), 539–544.
- [9] P.-E. Caprace and B. Remy, Simplicity and superrigidity of twin building lattices, *Invent. Math.* **176** (2009), 169–221.
- [10] C. R. J. Clapham, An embedding theorem for finitely generated groups, *Proc. Lond. Math. Soc. (3)* **17** (1967), 419–430.
- [11] F. Dahmani and D. Groves, The isomorphism problem for toral relatively hyperbolic groups, *Publ. Math. Inst. Hautes Études Sci.* **107** (2008), 211–290.
- [12] M. Dehn, Über unendliche diskontinuierliche Gruppen, *Math. Ann.* **71** (1911), 116–144.

- [13] P. Deligne and G. D. Mostow, *Commensurabilities Among Lattices in $PU(1, n)$* , Ann. of Math. Stud. 132, Princeton University Press, Princeton, 1993.
- [14] B. Farb, Relatively hyperbolic groups, *Geom. Funct. Anal.* **8** (1998), 810–840.
- [15] P. Hall, Finiteness conditions for soluble groups, *Proc. Lond. Math. Soc. (3)* **4** (1954), 419–436.
- [16] G. Higman, Subgroups of finitely presented groups, *Proc. R. Soc. Lond. Ser. A* **262** (1961), 455–475.
- [17] W. R. Parry J. W. Cannon and W. J. Floyd, Introductory notes on Richard Thompson’s groups, *Enseign. Math. (2)* **42** (1996), 215–256.
- [18] S. Lipschutz and C. F. Miller III, Groups with certain solvable and unsolvable decision problems, *Comm. Pure Appl. Math.* **24** (1971), 7–15.
- [19] R. C. Lyndon and P. E. Schupp, *Combinatorial Group Theory*, Ergeb. Math. Grenzgeb. 89, Springer-Verlag, Berlin, 1977.
- [20] G. A. Margulis, Arithmeticity of nonuniform lattices, *Funktsional. Anal. i Prilozhen.* **7** (1973), no. 3, 88–89.
- [21] C. F. Miller III, Decision problems for groups—survey and reflections, in: *Algorithms and Classification in Combinatorial Group Theory* (Berkeley 1989), Math. Sci. Res. Inst. Publ. 23, Springer-Verlag, New York (1992), 1–59.
- [22] D. V. Osin, Relatively hyperbolic groups: intrinsic geometry, algebraic properties, and algorithmic problems, *Mem. Amer. Math. Soc.* **179** (2006), no. 843.
- [23] M. O. Rabin, Recursive unsolvability of group theoretic problems, *Ann. of Math. (2)* **67** (1958), 172–194.
- [24] D. J. S. Robinson, *A Course in the Theory of Groups*, 2nd ed., Grad. Texts in Math. 80, Springer-Verlag, New York, 1996.
- [25] P. E. Schupp, Small cancellation theory over free products with amalgamation, *Math. Ann.* **193** (1971), 255–264.
- [26] R. E. Schwartz, The quasi-isometry classification of rank one lattices, *Publ. Math. Inst. Hautes Études Sci.* **82** (1995), 133–168.
- [27] C. L. Siegel, Symplectic geometry, *Amer. J. Math.* **65** (1943), 1–86.
- [28] J. Stillwell, *Classical Topology and Combinatorial Group Theory*, 2nd ed., Grad. Texts in Math. 72, Springer-Verlag, New York, 1993.
- [29] S. Thomas, The virtual isomorphism problem for finitely generated groups, *Bull. Lond. Math. Soc.* **35** (2003), 777–784.
- [30] S. Thomas, On the complexity of the quasi-isometry and virtual isomorphism problems for finitely generated groups, *Groups Geom. Dyn.* **2** (2008), 281–307.
- [31] S. Thomas, The commensurability relation for finitely generated groups, *J. Group Theory* **12** (2009), 901–909.

- [32] S. Thomas and B. Velickovic, On the complexity of the isomorphism relation for finitely generated groups, *J. Algebra* **217** (1999), 352–373.
- [33] K. Whyte, Amenability, bi-Lipschitz equivalence, and the von Neumann conjecture, *Duke Math. J.* **99** (1999), 93–112.

Received December 19, 2012; revised May 9, 2013.

Author information

Goulnara Arzhantseva, Faculty of Mathematics, University of Vienna,
Oskar-Morgenstern-Platz 1, 1090 Vienna, Austria.
E-mail: goulnara.arzhantseva@univie.ac.at

Jean-François Lafont, Department of Mathematics, The Ohio State University,
100 Math Tower, 231 West 18th Avenue, Columbus, OH 43210-1174, USA.
E-mail: jlafont@math.ohio-state.edu

Ashot Minasyan, School of Mathematics, University of Southampton,
Highfield campus, Southampton, SO17 1BJ, United Kingdom.
E-mail: aminasyan@gmail.com