

RANDOM TRIGONOMETRIC POLYNOMIALS: UNIVERSALITY AND NON-UNIVERSALITY OF THE VARIANCE FOR THE NUMBER OF REAL ROOTS

YEN DO, HOI H. NGUYEN, AND OANH NGUYEN

ABSTRACT. It follows from a recent result by Bally, Caramellino, and Poly that for random trigonometric polynomials with iid coefficients satisfying certain continuousness and having bounded moments of all orders, the variance of the number of roots is asymptotically linear in terms of the expectation; furthermore the multiplicative constant in this linear relationship depends only on the kurtosis of the common distribution of the polynomial's coefficients. In this note, we show that this is a universal phenomenon under a more general condition, which in particular is satisfied by discrete random variables. Our method gives a fine comparison framework throughout Edgeworth expansion, asymptotic Kac-Rice formula, and a detailed analysis of characteristic functions.

1. INTRODUCTION

Universality for the distribution of roots of random polynomials is an exciting subject that has attracted the attention of many generations. When the degree of a polynomial is very large, it is often challenging (even numerically) to solve for the roots, and a very natural question is to obtain an accurate estimate for the number of roots in a given region (in particular in \mathbb{R}). There is a large body of studies in the past centuries dedicated to this task, showing that the typical size of the number of roots depends mostly on the underlying symmetries of the random polynomials and not on the particular distributions of the coefficients. These authors often assume a fairly minimal normalization condition, where the coefficients are independent with fixed means and fixed variances. Results of this type are known in the literature as universality results for the number of (real) roots.

Among many statistics about the number of real roots of random polynomials, denoted by N_n (or $N_{n,\mathbb{R}}$), the followings are often considered first by many authors: the expectation $\mathbf{E}N_n$, the variance $\text{Var}(N_n)$, and the limiting distribution of the standardization $N_n^* = \frac{N_n - \mathbf{E}N_n}{\sqrt{\text{Var}(N_n)}}$. One of the most studied random polynomials in the literature is perhaps Kac's polynomial, $P_n(x) = \xi_0 + \xi_1 x + \cdots + \xi_n x^n$ where ξ_j are iid copies of a common random variable ξ , often assumed to have zero mean and unit variance. The issue of estimating N_n for such polynomials was already raised by Waring as far back as 1782 ([59], [40]). In the early 1940s, Kac [37] (see also [55]) developed a magnificent formula for the expectation of number of real roots

$$\mathbf{E}N_n = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |y| p(t, 0, y) dy dt, \quad (1)$$

where $p(t, x, y)$ is the probability density for $P_n(t) = x$ and $P'_n(t) = y$. See for instance [1, 3, 19] for other variants of this Kac-Rice formula. When ξ is standard gaussian one can easily evaluate the right-hand side above and obtain $\mathbf{E}N_n = (\frac{2}{\pi} + o(1)) \log n$. Similarly, one can also show that $\text{Var}(N_n) = (\frac{4}{\pi}(1 - \frac{2}{\pi}) + o(1)) \log n$.

Kac-Rice formula, as stated in (1), is effective only when ξ is smooth. For general ξ , thanks to the contributions of Littlewood and Offord [42, 43, 44, 45], Erdős and Offord [21], Ibragimov and Maslova [29, 30, 31, 32] (among others), universality for the leading asymptotics (as $n \rightarrow \infty$) for $\mathbf{E}N_n$ and $\text{Var}(N_n)$ is now fairly well-understood. It is also known that N_n^* converges in distribution to standard Gaussian. Furthermore, for all these results to hold it suffices to assume $\mathbf{E}(|\xi|^{2+\varepsilon_0}) < \infty$ for a fixed $\varepsilon_0 > 0$, see [53].

The first author is supported in part by NSF grant DMS-1800855.

The second author is supported by NSF grant DMS-1752345.

Proving universality for these statistics about the number of real roots for other classical random ensembles (such as Weyl polynomial, Elliptic polynomial, polynomials with asymptotic growth of coefficients, etc) has become an active direction of research in recent years [11, 12, 34, 36, 61, 17, 25, 26]. There is also a distinction between local and global universality. The *global universality* concerns the limiting distribution of the empirical measure of all complex roots and has been established in several papers for many random polynomials, see for instance [36, 54, 15] and the references therein. The *local universality* concerns the distribution of the roots (complex, real, or both) in smaller/thinner sets and is developed in a series of work by Tao, Vu, and the current authors [61, 50, 17, 52].

In the opposite direction, it seems natural to search for “non-trivial” statistics that are not universal with respect to the standard assumptions of zero mean and unit variance. This non-universality was indeed observed for the expected value of the number of real roots of Kac polynomial in joint work of the current authors with Vu [18, 50]. Here for the Gaussian case, results by Wilkins [64] and later rediscovered by Edelman and Kostlan [19] show that $\mathbf{E}N_n = \frac{2}{\pi} \log n + C_G + o(1)$ where $C_G \approx .625738072$ is an explicit constant,

$$C_G = \frac{2}{\pi} \left(\log 2 + \int_0^\infty \sqrt{\frac{1}{x^2} - \frac{4e^{-2x}}{(1 - e^{-2x})^2}} - \frac{1}{x+1} dx \right).$$

It turns out that [18, 50] this type of refined estimates also extends to the Rademacher case ($\xi = \pm 1$ with probability 1/2) or the uniform case (among others).

Theorem 1.1. *Assume that $\mathbf{E}|\xi|^{2+\varepsilon_0} < \infty$ for some fixed $\varepsilon_0 > 0$. Then we have $\mathbf{E}N_n = \frac{2}{\pi} \log n + O_\xi(1)$. Furthermore, if ξ is uniform on $\{\pm 1, \dots, \pm K\}$ (for a given positive integer K) or ξ has p -integrable density for some $p > 1$, then there is a constant $C = C_\xi$ such that*

$$\mathbf{E}N_n = \frac{2}{\pi} \log n + C_\xi + o(1). \quad (2)$$

The constants C_ξ are numerically observed to be different from C_G for various cases. It is certainly interesting to pinpoint a more accurate formula for C_ξ and specify how C_ξ depends on ξ . Also, we believe that (2) holds under more general assumption on ξ .

One of our main goals in this note is to address a similar phenomenon, but for another important class of random functions where the picture seems to be much cleaner. Consider the random trigonometric polynomial

$$P_n(t, Y) = \frac{1}{\sqrt{n}} \sum_{i=1}^n y_{i1} \cos\left(\frac{it}{n}\right) + y_{i2} \sin\left(\frac{it}{n}\right), \quad (3)$$

with $Y = (Y_1, \dots, Y_n)$, where $Y_i = (y_{i1}, y_{i2})$ and Y_i are independent. In this note, we are interested in the roots of $P_n(t, Y)$ (although some other statistics of random trigonometric polynomials also play crucial role in some recent interesting studies, such as [4],[56]). We now redefine $N_n = N_n(Y)$ to be the number of roots of P_n for $t \in [-n\pi, n\pi]$. It is known from a result of Qualls [57] in the 1970s that when the y_{ij} are iid standard Gaussian then we have

$$\mathbf{E}N_n = 2\sqrt{(2n+1)(n+1)}/6.$$

Confirming a striking heuristic by Bogomolny, Bohigas, and Leboeuf [14], about ten years ago Granville and Wigman [27] proved the following.

Theorem 1.2. *When the y_{ij} are iid standard Gaussian, there exists an explicit positive constant c_G such that the variance satisfies*

$$\text{Var}(N_n) = (c_G + o(1))n.$$

Furthermore,

$$\frac{N_n - \mathbf{E}N_n}{\sqrt{c_G n}} \xrightarrow{d} \mathbf{N}(0, 1).$$

Here asymptotically $c_G \approx 0.55826$. More precisely,

$$c_G = \frac{4}{3\pi} \int_0^\infty \left(\frac{1 - g(t)^2 - 3g'(t)^2}{(1 - g(t)^2)^{3/2}} (\sqrt{1 - R^{*2}} + R^* \arcsin R^*) - 1 \right) dt + \frac{2}{\sqrt{3}},$$

where

$$g(t) = \frac{\sin(t)}{t}, \text{ and } R^* = R^*(t) = \frac{g''(t)(1 - g(t)^2) + g(t)g'(t)^2}{1/3(1 - g(t)^2) - g'(t)^2}.$$

Granville and Wigman established this beautiful result by a delicate method basing on Kac-Rice formula. More recently, Azaïs and León [6] provided an important alternative approach basing on Wiener chaos decomposition. Roughly speaking, they showed that $P_n(t, Y)$ converges in certain strong sense to the stationary Gaussian process of covariance $r(t) = \sin(t)/t$, from which variance and CLT can be deduced.

More relevant to our current note, the above result has been extended recently by Bally, Caramellino, and Poly [8] to more general distributions where certain continuousness is assumed. To discuss this extension, we first recall some notions. We say that Y_i satisfies the (two-dimensional) Doeblin's condition if there exists $a_i \in \mathbb{R}^2$ and $r, \eta \in (0, 1)$ such that for any $A \subset B_r(a_i)$

$$\mathbf{P}(Y_i \in A) \geq \eta \lambda(A).$$

Let $\mathcal{D}(r, \eta)$ denote the sequences of r.v. $Y_k = (y_{k1}, y_{k2})$ with the above property, with $\mathbf{E}y_{k_j1}y_{k_j2} = \delta_{j_1j_2}$, and uniformly bounded moments of all orders

$$\sup_k \mathbf{E}|Y_k|^p < \infty \quad \forall p,$$

where the Y_k are independent but not necessarily identically distributed.

Suppose that $Y \in \mathcal{D}(r, \eta)$ and for all $\alpha = (\alpha_1, \dots, \alpha_m) \in \{1, 2\}^m$ with $m = 3, 4$ the following limits exist

$$\lim_{n \rightarrow \infty} \mathbf{E} \left(\prod_{i=1}^m y_{n\alpha_i} \right) = y_\infty(\alpha).$$

The following result from [8] was formulated for $N_n([0, n\pi], Y)$ the number of roots inside $[0, n\pi]$ of $P_n(nt, Y)$ ¹. Let $N_n([0, n\pi], G)$ be the number of roots inside $[0, n\pi]$ of $P_n(nt, G)$, which is the random polynomial with coefficients y_{ij} being standard Gaussian.

Theorem 1.3. [8, Theorem 2.1] *We have*

$$\lim_n \frac{1}{n} \text{Var}(N_n([0, n\pi], Y)) = \lim_n \frac{1}{n} \text{Var}(N_n([0, n\pi], G)) + \frac{1}{60} y_*$$

with

$$y_* = (y(1, 1, 2, 2) - 1) + (y(2, 2, 1, 1) - 1) + (y(1, 1, 1, 1) - 3) + (y(2, 2, 2, 2) - 3).$$

In particularly if y_{ij} are iid copies of a random variable ξ of mean zero and variance one and satisfies the (one-dimensional) Doeblin's condition, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}(N_n([0, n\pi], Y)) = \lim_n \frac{1}{n} \text{Var}(N_n([0, n\pi], G)) + \frac{1}{30} \mathbf{E}(\xi^4 - 3). \quad (4)$$

This result implies strong concentration around the mean of N_n . More crucially, it says that the variance is not universal with respect to second order normalization of ξ (having mean zero and variance one). At the same time, it also suggests a possible universal picture that the limit ratio V_n/n asymptotically depends on y_* , and in particularly on the fourth moment in the iid case. In this note we confirm this phenomenon by working with the iid case without assuming the Doeblin's condition.

¹The authors of [8] considered the number of roots inside $[0, \pi]$ of $P_n(nt, Y)$, which is the same as our $N_n([0, n\pi], Y)$.

Theorem 1.4 (main theorem). *Assume that $y_{ij}, 1 \leq i \leq n, j = 1, 2$ are iid copies a random variable ξ of mean zero, variance one, and $\mathbf{E}|\xi|^{M_0} < \infty$ for a sufficiently large positive number M_0 . Then*

$$\lim_n \frac{1}{n} \text{Var}(N_n) = c_G + \frac{2}{15} \mathbf{E}(\xi^4 - 3),$$

where we recall that c_G is the constant from Theorem 1.2.

We thus obtain that for the case where $y_{i,j}$ are Rademacher random variables,

$$\lim_n \frac{1}{n} \text{Var}(N_n) = c_G - \frac{4}{15} \approx 0.29159.$$

Our numerical experiments appear to be in accordance with these results as shown in Figure 1.

Note that our result is stated for the number of roots over $[-n\pi, n\pi]$, but the approach automatically works for roots over $[0, n\pi]$ as well. As a matter of fact, most of our arguments work for random variables $|\xi|$ of bounded $(2 + \varepsilon_0)$ -moment, except at the Edgeworth expansion step (for instance Theorem 4.1) where we assume boundedness of moments.

We can view Theorem 1.3 and Theorem 1.4 as a mixture of universality and non-universality. The fact that the variance is linear in n indicates that there is no correlation (repulsion and attraction) among sufficiently far apart roots, and this phenomenon is universal in the sense that it suffices to assume $|\xi|$ to have bounded moments. However, the multiplicative constant, which is determined by the correlation of nearby roots, is affected by the kurtosis as seen.

Finally, we also invite the reader to Theorem 8.1 which says that under a very general setting (including the non-iid case) there is already a significant cancellation in the variance formula. More precisely, there exists a positive constant c such that

$$\text{Var}(N_n) = O(n^{2-c}). \tag{5}$$




FIGURE 1. Sample variance (divided by n) of the number of roots in $[-n\pi, n\pi]$ for Gaussian random variables (dashed line) and Rademacher random variables (solid line).

2. OUR METHODS

We first mention briefly the approach by Bally et al. to prove Theorem 1.3. Here, powerful tools such as Malliavin calculus and Wiener chaos theory (see [6] and the references therein) do not apply under the




FIGURE 2. Sample variance (divided by n) of the number of roots in $[0, n\pi]$ for Gaussian random variables (dashed line) and Rademacher random variables (solid line).

Doebelin’s condition. Instead, the authors above have developed a sophisticated method using the Edgeworth expansion and approximate Kac-Rice formulas basing on their previous results in [7].

Generally speaking, for Theorem 1.4 we will follow the same machinery. However, as we have to deal with discrete random variables, none of the results from [7] and [8] could be applied. For instance, in our opinion, it is a non-trivial problem to study the small ball probability for the random walks associated to $P_n(t)$ without Doebelin assumption.

We would also like to point out that, broadly speaking, using the Edgeworth expansion to study the distribution of normalized sums of independent random variables is a classical approach (see [9]) and this approach was also used by Bally et al. [7, 8]. The novelty in our argument is a more refined estimate for characteristics functions motivated by work of Konyagin-Schlag [41] that works for a large class of distributions (including the discrete cases). This is where we deviate from Bally et al. [7, 8], who used a completely different approach to deal with non-smooth distributions. More precisely, in their papers [7, 8], the authors use the Nummelin splitting (which requires Doebelin condition) to decompose non-smooth distribution into two parts: a smooth part that can be treated directly by Edgeworth expansion methods, and a noisy part that can be treated by Wiener chaos techniques. Our modified approach circumvents the need for Nummelin’s splitting and therefore avoids the need for anti-concentration conditions like the Doebelin condition in Bally et al. [7, 8].

One trade-off that we need to face in order to obtain the generality of our result is that we necessarily rule out a set of points that are well-approximated by the integer lattice (see Condition 2). To show that this set does not contribute significantly to the whole picture, we utilize a universality result in [52] (Theorem 7.7) which, roughly speaking, says that the difference between the variance of the number of roots of $P_n(\cdot, Y)$ and $P_n(\cdot, G)$ over small intervals is negligible.

In what follows we sketch the highlights, some of which are of independent interest. (For instance a variant of Theorem 2.2 finds some applications in [51].)

2.1. Small ball estimates and characteristic functions. Here we only assume ξ to have mean zero, variance one and bounded $(2 + \varepsilon_0)$ -moment for any $\varepsilon_0 > 0$.

For $t \in [-n\pi, n\pi]$, we define the vectors

$$\mathbf{u}_i(t) := \left(\cos\left(\frac{it}{n}\right), -\frac{i}{n} \sin\left(\frac{it}{n}\right) \right), \text{ and } \mathbf{u}'_i(t) := \left(\sin\left(\frac{it}{n}\right), \frac{i}{n} \cos\left(\frac{it}{n}\right) \right). \quad (6)$$

Assume that y_{ij} , $1 \leq i \leq n$, $j = 1, 2$ are iid copies of a random variable ξ of mean zero, variance one. Consider the random walk in \mathbb{R}^2

$$S_n(Y, t) := \sum_{i=1}^n y_{i1} \mathbf{u}_i + y_{i2} \mathbf{u}'_i. \quad (7)$$

This random walk can also be written as $S_n(t, Y) = \sum_{i=1}^n C_n(i, t) Y_i$, where $Y_i = (y_{i1}, y_{i2})$ and

$$C_n(i, t) = \begin{pmatrix} \cos\left(\frac{it}{n}\right) & \sin\left(\frac{it}{n}\right) \\ -\frac{i}{n} \sin\left(\frac{it}{n}\right) & \frac{i}{n} \cos\left(\frac{it}{n}\right) \end{pmatrix}. \quad (8)$$

Note that for some values of t such as $t = o(1)$, the random walk does not spread out in the Radamacher case. We will show that these are the only cases to cause this clustering.

Condition 1. Let τ be a constant to be chosen sufficiently small. A number $t \in [-n\pi, n\pi]$ is said to satisfy Condition 1 if there does not exist a non-zero integer l with $|l| \leq n^\tau$ such that

$$\left\| l \frac{t}{\pi n} \right\|_{\mathbb{R}/\mathbb{Z}} \leq n^{-1+8\tau}.$$

Here $\|\cdot\|_{\mathbb{R}/\mathbb{Z}}$ is the distance to the nearest integer. In other words, the above condition requires that $t/\pi n$ cannot be within a distance of $n^{-1+o(1)}$ from rational numbers of denominator $n^{o(1)}$.

Theorem 2.2. Let $C > 0$ be a given constant. Assume that t satisfies Condition 1 with sufficiently small τ . Then for $\delta = n^{-C}$ and any open ball $B(a, \delta)$, we have

$$\mathbf{P} \left(\frac{1}{\sqrt{n}} S_n(t, Y) \in B(a, \delta) \right) = O(\delta^2).$$

As mentioned before, our condition on t is almost optimal. Towards Theorem 1.4, as we will be dealing with pair correlations, we will need to work with vectors in \mathbb{R}^4 . Let s, t be given, define the vectors $\mathbf{v}_i, \mathbf{v}'_i$ as follows

$$\mathbf{v}_i(s, t) := \left(\cos\left(\frac{it}{n}\right), -\frac{i}{n} \sin\left(\frac{it}{n}\right), \cos\left(\frac{is}{n}\right), -\frac{i}{n} \sin\left(\frac{is}{n}\right) \right) \quad (9)$$

and

$$\mathbf{v}'_i(s, t) = \left(\sin\left(\frac{it}{n}\right), \frac{i}{n} \cos\left(\frac{it}{n}\right), \sin\left(\frac{is}{n}\right), \frac{i}{n} \cos\left(\frac{is}{n}\right) \right) \quad (10)$$

Here we are interested in the random walk

$$S_n(s, t, Y) := \sum_{i=1}^n y_{i1} \mathbf{v}_i + y_{i2} \mathbf{v}'_i. \quad (11)$$

Using (8), if we let $C_n(i, s, t)$ be the matrix size 4×2 as a joint of $C_n(i, t)$ and $C_n(i, s)$, then we can see that this random walk can also be written as $S_n(s, t, Y) = \sum_{i=1}^n C_n(i, s, t) Y_i$.

We will assume that $s/\pi n$ and $t/\pi n$ cannot be jointly well-approximated by rational numbers.

Condition 2. Let τ be a constant to be chosen sufficiently small. Two numbers $s, t \in [-n\pi, n\pi]$ are said to satisfy Condition 2 if there do not exist integers k, l with $|k|, |l| \leq n^\tau$, not both zero, such that

$$\left\| k \frac{s}{\pi n} + l \frac{t}{\pi n} \right\|_{\mathbb{R}/\mathbb{Z}} \leq n^{-1+8\tau}.$$

Note that if s, t satisfy Condition 2 then each of them satisfies Condition 1 separately. It is clear that the measure of $(s/n, t/n) \in [-\pi, \pi]^2$ that does not satisfy the above condition is $n^{-1+O(\tau)}$. We will show the following small ball probability.

Theorem 2.3. *Let $C > 0$ be a given constant. Assume that s, t satisfy Condition 2 with sufficiently small τ . Then for $\delta = n^{-C}$ we have*

$$\mathbf{P} \left(\frac{1}{\sqrt{n}} S_n(s, t, Y) \in B(a, \delta) \right) = O(\delta^4).$$

To prove these small ball estimates we will rely on the following results on the characteristic functions. First, for the random walk $S_n(t, Y)$, let

$$\phi_{\mathbb{R}^2}(x) = \prod_{i=1}^n \phi_i(x) = \prod_{i=1}^n \mathbf{E} e(y_{i1} \langle \mathbf{u}_i, x \rangle) \prod_{i=1}^n \mathbf{E} e(y_{i2} \langle \mathbf{u}'_i, x \rangle), x \in \mathbb{R}^2,$$

where $e(y) = e^{iy}$. We will show that this function decays very fast.

Theorem 2.4. *Let $C_* > 0$ be any given constant, and t satisfies Condition 1 for some sufficiently small constant τ . Then the following holds for sufficiently large n and sufficiently small τ_* (depending on C_*, τ). For any $n^{5\tau-1/2} \leq \|x\|_2 \leq n^{C_*}$ we have*

$$|\phi_{\mathbb{R}^2}(x)| \leq \exp(-n^{\tau_*}).$$

We note that this was also studied in [41] for the Radamacher case, covering up to $\|x\|_2 \leq n^{1/2+o(1)}$. This result has been improved to $\|x\|_2 \leq n^{1-o(1)}$ in [51] recently for any ξ of variance one. Our current approach to prove Theorem 2.4 goes deeper than those of [41, 51] where we need to solve certain inverse-type problems. (See Sections 10 and 9 for more details.)

Similarly to the case of \mathbb{R}^2 , to establish these results we will study the characteristic function

$$\phi_{\mathbb{R}^4}(x) = \prod_{i=1}^n \phi_i(x) = \prod_{i=1}^n \mathbf{E} e(z_i \langle \mathbf{v}_i, x \rangle) \prod_{i=1}^n \mathbf{E} e(z'_i \langle \mathbf{v}'_i, x \rangle).$$

Theorem 2.5. *Let $C_* > 0$ be any given constant, and assume that s, t satisfy Condition 2 for some sufficiently small constant τ . Then the following holds for sufficiently large n and sufficiently small τ_* (depending on C_*, τ). For any $n^{5\tau-1/2} \leq \|x\|_2 \leq n^{C_*}$ we have*

$$|\phi_{\mathbb{R}^4}(x)| \leq \exp(-n^{\tau_*}).$$

We note that this result implies Theorem 2.4 because with t under Condition 1, there exists $s \in [-n\pi, n\pi]$ so that s, t satisfies Condition 2. We then apply Theorem 2.5 with $x = (x_1, x_2, 0, 0)$. However, we will present a separate proof of Theorem 2.4 in Section 9 to serve as a preparation for our more technical treatment of Theorem 2.5 in Section 10.

2.6. Approximated Kac-Rice formula and proof conclusion. We next briefly recall the use of approximated Kac-Rice.

Consider a smooth function f on an interval $[a, b]$ where for all $t \in [a, b]$ we have $|f(t)| + |f'(t)| > 0$. Then according to a celebrated formula of Kac and Rice, the number of roots of f in $[a, b]$ is given by

$$\lim_{\delta \rightarrow 0} \frac{1}{2\delta} \int_a^b |f'(t)| 1_{|f(t)| < \delta} dt.$$

Using this approximated formula for our polynomial $P_n(\cdot)$ we will show that for

$$\delta = \delta_n = n^{-5}, \tag{12}$$

we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \text{Var}(N_n(Y)) = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Var} \left(\frac{1}{2\delta} \int_{-n\pi}^{n\pi} P'_n(t, Y) 1_{|P_n(t, Y)| < \delta} \right) dt.$$

After expanding out the integrals, we will need to compute

$$\frac{1}{\delta^2} \int_{-n\pi}^{n\pi} \int_{-n\pi}^{n\pi} \text{Cov}(P'_n(t, Y) 1_{|P_n(t, Y)| < \delta}, P'_n(s, Y) 1_{|P_n(s, Y)| < \delta}) ds dt.$$

Let us introduce a few notations to simplify the discussion. We define the following even functions

$$F_\delta(x) = \frac{1}{2\delta} 1_{|x| < \delta}, x \in \mathbb{R} \quad (13)$$

and

$$\Phi_\delta(x) = |x_2| F_\delta(x_1), x = (x_1, x_2) \in \mathbb{R}^2 \quad (14)$$

and

$$\Psi_\delta(x) = \Phi_\delta(x_1, x_2) \Phi_\delta(x_3, x_4) = |x_2| F_\delta(x_1) |x_4| F_\delta(x_3), x = (x_1, x_2, x_3, x_4) \in \mathbb{R}^4. \quad (15)$$

We have

$$\Phi_\delta \left(\frac{1}{\sqrt{n}} S_n(t, Y) \right) = |P'_n(t, Y)| \times \frac{1}{2\delta} 1_{|y| \leq \delta}(P_n(t, Y)) =: \phi_\delta(t, Y)$$

and

$$\Psi_\delta \left(\frac{1}{\sqrt{n}} S_n(s, t, Y) \right) = \phi_\delta(s, Y) \phi_\delta(t, Y).$$

Finally, for short we introduce

$$\begin{aligned} v_n(s, t, Y) &:= \text{Cov}(P'_n(s, Y) 1_{|P_n(s, Y)| < \delta}, P'_n(t, Y) 1_{|P_n(t, Y)| < \delta}) \\ &= \mathbf{E} \Psi_\delta \left(\frac{1}{\sqrt{n}} S_n(s, t, Y) \right) - \mathbf{E} \phi_\delta(s, Y) \mathbf{E} \phi_\delta(t, Y). \end{aligned} \quad (16)$$

For given $\varepsilon > 0$ we will decompose the interval $(-n\pi, n\pi)$ into subintervals of length ε ,

$$I_k := [k\varepsilon, (k+1)\varepsilon] \subset [-n\pi, n\pi]. \quad (17)$$

Let

$$D_{n, \varepsilon} := \bigcup_{(k, p) \in \mathcal{D}} I_k \times I_p \quad (18)$$

where \mathcal{D} is the set consisting of all (k, p) with $-n\pi/\varepsilon \leq k < p \leq n\pi/\varepsilon$ such that for all $s \in I_p$ and $t \in I_k$, s and t satisfy Condition (2).

Let $N_n(G)$ and $v_n(s, t, G)$ be the statistics when the y_{ij} are standard Gaussian. In our next lemma we show that, in comparison with the Gaussian part, the contribution R_n from $(s, t) \notin D_{n, \varepsilon}$ is negligible in the variance computation.

Lemma 2.7. *With δ as in (12) we have*

$$\text{Var} N_n(Y) = \text{Var} N_n(G) + 2 \int_{D_{n, \varepsilon}} (v_n(s, t, Y) - v_n(s, t, G)) ds dt + R_{n, \varepsilon} \quad (19)$$

where

$$\lim_n \frac{R_{n, \varepsilon}}{n} = 0.$$

Therefore, we will need to control $\int_{D_{n, \varepsilon}} (v_n(s, t, Y) - v_n(s, t, G)) ds dt$ from (19), for which we will use Proposition 2.10 to show the following (see also [8, Lemma 5.1]).

Proposition 2.8. *For every $\varepsilon > 0$ we have*

$$\lim_n \frac{1}{n} \int_{D_{n, \varepsilon}} (v_n(s, t, Y) - v_n(s, t, G)) ds dt = \frac{1}{15} \mathbf{E}(\xi^4 - 3) + r_\varepsilon$$

with $|r_\varepsilon| = O(\varepsilon)$.

Combining Lemma 2.7 and Proposition 2.8, with $\varepsilon \rightarrow 0$, we obtain Theorem 1.4. We will prove Lemma 2.7 in Section 7 and Proposition 2.8 in Section 6 respectively. Notice that for these results we will also need to incorporate other existing results in the literature (notably [52]). We will also justify (5) by the same way (see Section 8).

2.9. Edgeworth expansion. We now compare $v_n(s, t, Y)$ with $v_n(s, t, G)$ by using Edgeworth expansion of order three. This approach is originated from [8], but our proof is directly based on the study of characteristic functions.

If X_i are iid real random variables of mean zero and variance one, the Central Limit Theorem says that, with Φ being the C.D.F. of the standard Gaussian distribution, for any x we have

$$\lim_{n \rightarrow \infty} \left| \mathbf{P} \left(\frac{S_n}{\sqrt{n}} \leq x \right) - \Phi(x) \right| = 0$$

where $S_n = \sum_{k=1}^n X_k$.

The Edgeworth expansion by Edgeworth [22], Chebyshev [63] and Cramér [16] says that under the so-called Cramér condition, if the X_i has bounded s_0 moments then there exist explicit polynomials P_1, \dots, P_{s_0-1} with coefficients depend on the cumulants of $\frac{S_n}{\sqrt{n}}$ such that

$$\left| \mathbf{P} \left(\frac{S_n}{\sqrt{n}} \leq x \right) - \sum_{r=0}^{s_0-1} n^{-r/2} P_r(-D)(\Phi(x)) \right| = O(n^{-s_0/2})$$

where D is the differential operator.

To prove Proposition 2.8 we will carry out the Edgeworth expansion for $\mathbf{E}\Psi_\delta(\frac{1}{\sqrt{n}}S_n(s, t, Y))$ as well as for $\mathbf{E}\Phi_\delta(\frac{1}{\sqrt{n}}S_n(s, Y))$ and $\mathbf{E}\Phi_\delta(\frac{1}{\sqrt{n}}S_n(t, Y))$, where we recall $S_n(t, Y)$ and $S_n(s, t, Y)$ from (7) and (11).

In what follows we mention briefly our contribution, we invite the reader to Section 4 and Section 5 for more details.

We let $X_n(t, Y)$ be the vector $(C_n(k, t)Y_k)_{k=1}^n$ and $X_n(s, t, Y)$ be the vector $(C_n(k, s, t)Y_k)_{k=1}^n$. We also let $V_n(t) = \frac{1}{n} \sum_{k=1}^n C_n(k, t)C_n(k, t)^*$ and $V_n(s, t) = \frac{1}{n} \sum_{k=1}^n C_n(k, s, t)C_n(k, s, t)^*$ be the average covariance matrices. Finally, we defer our definition of $\Gamma_{n,2}$ to (43). We will show the following CLT type estimates.

Proposition 2.10. *Assume that ξ mean zero and variance one and $\mathbf{E}|\xi|^{M_0} < \infty$ for sufficiently large M_0 . Assume that s, t satisfy Condition 2. Then we have*

$$|\mathbf{E}F_\delta(P_n(t, Y)) - \mathbf{E}F_\delta(P_n(t, G))| \leq \frac{C}{n^{1/2}}, \quad (20)$$

and

$$\begin{aligned} & \left| \mathbf{E}\Phi_\delta \left(\frac{1}{\sqrt{n}}S_n(t, Y) \right) - \mathbf{E}\Phi_\delta \left(\frac{1}{\sqrt{n}}S_n(t, G) \right) - \frac{1}{n} \mathbf{E}[\Phi_\delta(I_2(\lambda)^{1/2}W_2)\Gamma_{n,2}(I_2(\lambda)^{-1/2}X_n(t, Y), W_2)] \right| \\ & \leq \frac{C}{n^{3/2}} + \frac{1}{n}r_n(t, \Phi_\delta), \end{aligned} \quad (21)$$

and

$$\begin{aligned} & \left| \mathbf{E}\Psi_\delta \left(\frac{1}{\sqrt{n}}S_n(s, t, Y) \right) - \mathbf{E}\Psi_\delta \left(\frac{1}{\sqrt{n}}S_n(s, t, G) \right) - \frac{1}{n} \mathbf{E}[\Psi_\delta(I_4(\lambda)^{1/2}W_4)\Gamma_{n,2}(I_4(\lambda)^{-1/2}X_n(s, t, Y), W_4)] \right| \\ & \leq \frac{C}{n^{3/2}} + \frac{1}{n}r_n(s, t, \Psi_\delta), \end{aligned} \quad (22)$$

where $I_2(\lambda)$ and $I_4(\lambda)$ are any invertible diagonal matrices² and W_2, W_4 are standard Gaussian vectors in \mathbb{R}, \mathbb{R}^2 and \mathbb{R}^4 respectively, and where the implied constants are allowed to depend on the M_0 -moment of ξ , on the constants in Conditions 1 and 2, and on a lower bound of the least singular values of $V_n(t), V_n(s, t)$ and $I_2(\lambda), I_4(\lambda)$. Furthermore we have the following bounds

$$r_n(t, \Phi_\delta) = O(\|V_n(t) - I_2(\lambda)\|_2) \text{ and } r_n(s, t, \Psi_\delta) = O(\|V_n(s, t) - I_4(\lambda)\|_2).$$

We also refer the reader to [8, Section 3] where a better error bound was obtained under the Doeblin's conditions. In application (Section 6) we will choose $I_2(\lambda)$ and $I_4(\lambda)$ so that $r_n \rightarrow 0$.

We will prove Proposition 2.10 by giving a general Edgeworth expansion result in Section 4, and then use it to conclude the proof in Section 5. Roughly speaking, our approach here is based on the work of Bhattacharya and Rao [9] (see also [2]) which relates Edgeworth expansion to the growth of characteristic functions of the corresponding random walks.

Notations. Throughout the note n is the parameter to be sent to ∞ . We write $X = O(Y)$, $Y = \Omega(X)$, $X \ll Y$, or $Y \gg X$ if $|X| \leq CY$ for some fixed C ; this C can depend on other fixed quantities such as the M_0 -moment of ξ . If $X \ll Y$ and $Y \ll X$, we say that $Y = \Theta(X)$ or $X \asymp Y$. We write $\omega(1)$ for a number that tends to ∞ as $n \rightarrow \infty$.

3. SMALL BALL PROBABILITY

In this section we address the small ball probabilities, we will just prove the \mathbb{R}^4 case (i.e. $d = 4$) because the \mathbb{R}^2 case can be proved similarly (by using Theorem 2.4 instead of Theorem 2.5).

Proof. (of Theorem 2.3) Let

$$t_0 = \delta^{-1}.$$

By a standard procedure (see for instance [2, Eq. 5.4]) we can bound the small ball probability by

$$\mathbf{P}\left(\frac{1}{\sqrt{n}} \sum_i z_i \mathbf{v}_i + z'_i \mathbf{v}'_i \in B(a, \delta)\right) \leq C_d \left(\frac{n}{t_0^2}\right)^{d/2} \int_{\mathbb{R}^d} \prod_i \phi_i(u) e^{-\frac{n\|u\|_2^2}{2t_0^2}} du.$$

Choose C_* to be sufficiently large compared to C . We break the integral into three parts, J_1 when $\|u\|_2 \leq r_0 = O(1)$, J_2 when $r_0 \leq \|u\|_2 \leq R = n^{C_*}$, and J_3 for the remaining part.

For J_1 , recall that

$$\left| \prod_i \phi_i(u) \right| \leq \exp\left(-\sum_i \|\langle \mathbf{v}_i, u \rangle\|_z^2 / 2\right).$$

So if $\|u\|_2 \leq c$ for sufficiently small c , then we have $\|\langle \mathbf{v}_i, u \rangle\|_{\mathbb{R}/\mathbb{Z}} = \|\langle \mathbf{v}_i, u \rangle\|_2$, and so because of Condition 2 (where we would need that $\sum_i \langle \mathbf{e}, \mathbf{v}_i \rangle^2 \geq c'n$) we have

$$\sum_i \|\langle \mathbf{v}_i, u \rangle\|_z^2 / 2 = \sum_i \|\langle \mathbf{v}_i, u \rangle\|_2^2 / 2 \geq c'n \|u\|_2^2.$$

²The vector parameter λ stands for the diagonal entries, see (47).

Thus

$$\begin{aligned}
J_1 &= C_d \left(\frac{n}{t_0^2}\right)^{d/2} \int_{\|u\|_2 \leq r_0} \prod_i \phi_i(u) e^{-\frac{n\|u\|_2^2}{2t_0^2}} du \\
&\leq C_d \left(\frac{n}{t_0^2}\right)^{d/2} \int_{\|u\|_2 \leq r_0} e^{-\frac{n\|u\|_2^2}{2t_0^2} - c'n\|u\|_2^2} du \\
&= C_d \left(\frac{n}{t_0^2}\right)^{d/2} \int_{\|u\|_2 \leq r_0} e^{-(\frac{n}{2t_0^2} + c'n)\|u\|_2^2} du \\
&= O_d\left(\frac{1}{(c'n t_0^2 + 1)^{d/2}}\right) \\
&= O_d(\delta^d).
\end{aligned}$$

For J_2 , recall by Theorem 2.5 that for $r_0 \leq \|u\|_2 \leq R = n^{C_*}$ we have

$$|\prod_i \phi_i(u)| \leq e^{-n^{-\tau*}}.$$

Thus

$$\begin{aligned}
J_2 &= C_d \left(\frac{n}{t_0^2}\right)^{d/2} \int_{r_0 \leq \|u\|_2 \leq R} \prod_i \phi_i(u) e^{-\frac{n\|u\|_2^2}{2t_0^2}} du \\
&\leq C_d \left(\frac{n}{t_0^2}\right)^{d/2} \int_{r_0 \leq \|u\|_2 \leq R} e^{-n^{\tau*}} du \\
&= O_d(n^d \left(\frac{n}{t_0^2}\right)^{d/2} e^{-n^{\tau*}}) \\
&= O_d(e^{-n^{\tau*/2}}).
\end{aligned}$$

For J_3 , we have

$$\begin{aligned}
J_3 &= C_d \left(\frac{n}{t_0^2}\right)^{d/2} \int_{\|u\|_2 \geq n^{C_*}} \prod_i \phi_i(u) e^{-\frac{n\|u\|_2^2}{2t_0^2}} du \\
&= O_d(e^{-n})
\end{aligned}$$

as we chose C_* sufficiently large compare to C . □

Before concluding this section we introduce some useful corollaries of our small ball estimates. For short, let $\mathcal{G} = \mathcal{G}_\tau$ be the collection of $t \in [-n\pi, n\pi]$ that satisfies Condition 1.

We first deduce from Theorem 2.2 a small ball estimate for $P_n(t, Y)$ alone, which will be useful later.

Corollary 3.1. *Let $C > 1$ be a given constant. Assume that y_{ij} are iid copies of a bounded random variable ξ of mean zero, variance one, and bounded $\mathbf{E}(|\xi|^{M_0}) < \infty$ for some even positive integer M_0 . Assume that $t \in \mathcal{G}$ with sufficiently small τ . Then for $\delta = n^{-C}$ and any open interval $(a - \delta, a + \delta)$ we have*

$$\mathbf{P}(P_n(t, Y) \in (a - \delta, a + \delta)) = O\left(n^{\frac{M_0}{M_0+1}} \delta^{\frac{M_0}{M_0+1}}\right).$$

Proof. Since the random variables y_{i1}, y_{i2} are uncorrelated with mean 0 and bounded M_0 moments, we easily have

$$\mathbf{E}(P'_n(t, Y))^{M_0} = O_{M_0}(n^{M_0}).$$

Thus, by Markov's inequality, for a positive constant $A > 1$ to be chosen,

$$\mathbf{P}(|P'_n(t, Y)| \geq n^A) = P(n^{M_0 - M_0 A}).$$

We have

$$\mathbf{P}(|P_n(t, Y)| < \delta) \leq \mathbf{P}(|P'_n(t, Y)| \geq n^A) + \mathbf{P}(|P_n(t, Y)| < \delta, |P'_n(t, Y)| \leq n^A).$$

Since the latter event is a subset of a union of $n^A \delta^{-1}$ events of the form $S_n(Y, t) \in B(a, \delta)$ for some $a \in \mathbf{C}$, we apply Theorem 2.2 to get

$$\mathbf{P}(|P_n(t, Y)| < \delta) = O(n^{M_0 - M_0 A} + n^A \delta^{-1} \delta^2).$$

By choosing $A = \frac{M_0 + C}{1 + M_0}$, this proves Corollary 3.1. \square

Our next corollary is the following analog of [8, Eq. 3.40].

Theorem 3.2. *Let θ and $\varepsilon < 1/2$ be given constants. Assume that y_{ij} are iid copies of a bounded random variable ξ of mean zero, variance one, and bounded $\mathbf{E}(|\xi|^{M_0}) < \infty$ for sufficiently large M_0 (in terms of θ). We have*

$$\mathbf{P}(\inf_{|t| \in \mathcal{G}} |\frac{1}{\sqrt{n}} S_n(t, Y)| \leq n^{-\theta + \varepsilon/2}) = O(n^{-\theta + 1 + \varepsilon}).$$

Proof. First of all, let $\mathcal{E}_{\mathbf{b}}$ be the event that $|y_{ij}| \leq n^\varepsilon$ for all i, j . Then as M_0 is sufficiently large, by a union bound and by Markov's inequality we have

$$\mathbf{P}(\mathcal{E}_{\mathbf{b}}^c) \leq 2n \mathbf{P}(|\xi| > n^\varepsilon) \leq 2n \mathbf{E}(|\xi|^{M_0}) / n^{\varepsilon M_0} = O(n^{-\varepsilon M_0 + 1}) = O(n^{-\varepsilon M_0/2}).$$

Hence it suffices to condition on $\mathcal{E}_{\mathbf{b}}$. Next, for any fixed t we control the magnitude of

$$d(\frac{1}{\sqrt{n}} S_n(t, Y))/dt = (f_1(t, Y), f_2(t, Y))$$

where

$$f_1(t, Y) = \frac{1}{\sqrt{n}} \sum_{i=1}^n y_{i1} (-\frac{i}{n} \sin(\frac{it}{n})) + y_{i2} (\frac{i}{n} \cos(\frac{it}{n})) \text{ and } f_2(t, Y) = -\frac{1}{\sqrt{n}} \sum_{i=1}^n y_{i1} (\frac{i}{n})^2 \cos(\frac{it}{n}) + y_{i2} (\frac{i}{n})^2 \sin(\frac{it}{n}).$$

For this, again as ξ has mean zero and variance one and $\mathbf{E}(|\xi|^{M_0}) < \infty$, a moment computation shows that as long as $|c_i|, |d_i| \leq 1$ we have

$$\mathbf{E}(|\frac{1}{\sqrt{n}} \sum_{i=1}^n c_i y_{i1} + d_i y_{i2}|^{M_0}) = O_{M_0}(1).$$

Therefore for any fixed t we have

$$\mathbf{P}(|f_1(t, Y)| \geq n^{\varepsilon/2}) = O(n^{-\varepsilon M_0/2}) \text{ and } \mathbf{P}(|f_2(t, Y)| \geq n^{\varepsilon/2}) = O(n^{-\varepsilon M_0/2}). \quad (23)$$

Notice that on $\mathcal{E}_{\mathbf{b}}$ we trivially have $\sup_{t \in [-n\pi, n\pi]} |f'_i(t, Y)| = O(n^{1/2})$. By a standard net argument we obtain from (23) that

$$\mathbf{P}(\sup_{t \in [-n\pi, n\pi]} \|d(\frac{1}{\sqrt{n}} S_n(t, Y))/dt\|_2 \geq n^{\varepsilon/2}) = O(n^{-\varepsilon M_0/4}). \quad (24)$$

We will condition the complement of this event. Decompose \mathcal{G} into $O(n^{1+\theta})$ intervals of length $n^{-\theta}$ each, whose midpoints satisfy Condition 1. For each such interval I we estimate the probability that $\inf_{t \in I} |S_n(t, Y)| \leq n^{-\theta}$. By (24), this implies that for the midpoint t_I we have

$$\frac{1}{\sqrt{n}} S_n(t_I, Y) \leq n^{-\theta} + n^{\varepsilon/2} n^{-\theta} = O(n^{\varepsilon/2 - \theta}).$$

However, by using Theorem 2.2, we can control this event by

$$\mathbf{P}(|\frac{1}{\sqrt{n}} S_n(t_I, Y)| \leq n^{-\theta + \varepsilon/2}) = O(n^{-2\theta + \varepsilon}).$$

Taking union bounds over the midpoints of the $O(n^{1+\theta})$ intervals we obtain the bound $O(n^{-\theta + 1 + \varepsilon})$ as claimed, provided that M_0 is sufficiently large. \square

4. EDGEWORTH EXPANSION INVOLVING TRIGONOMETRIC FUNCTIONS

Our goal in this section is to establish an Edgeworth expansion for several sums of random vectors that arise from random trigonometric functions. The results are formulated under very mild assumptions on the coefficient distribution(s), which hold in discrete settings (such as the Rademacher distribution) beyond the scope of the Cramér condition and known extensions [2].

Let $s, t \in \mathbb{R}$ be given. Let $d = 4$. Consider the following sequence of random vectors in \mathbb{R}^d

$$X_{n,k} := C_n(k)Y_k, \quad k = 1, \dots, n, \quad (25)$$

where (i) Y_k 's are random vectors in \mathbb{R}^2 and their coordinates are iid with mean zero and variance one (we'll actually assume in our result that furthermore $\mathbf{E}\|Y_j\|_2^{\ell+d+1} < \infty$ for some $\ell \geq 4$), and (ii) the deterministic $d \times 2$ matrices $C_n(k)$ are defined below. Recall from Subsection 2.1 that

$$C_n(k, t) = \begin{pmatrix} \cos(\frac{kt}{n}) & \sin(\frac{kt}{n}) \\ -\frac{k}{n} \sin(\frac{kt}{n}) & \frac{k}{n} \cos(\frac{kt}{n}) \end{pmatrix} \quad (26)$$

and

$$C_n(k) = \begin{pmatrix} C_n(k, t) \\ C_n(k, s) \end{pmatrix} \quad (27)$$

is the 4×2 matrix obtained as the joint of $C_n(k, t)$ and $C_n(k, s)$. Recall also $S_n(s, t, Y)$ from (11), and for short let

$$S_n := S_n(s, t, Y) =: X_{n,1} + \dots + X_{n,n}. \quad (28)$$

Let the average covariance matrix be

$$V_n := \frac{1}{n} \sum_{k=1}^n C_n(k)C_n(k)^*. \quad (29)$$

This is the same as the covariance for S_n/\sqrt{n} . Let \tilde{Q}_n denote the distribution of S_n/\sqrt{n} , and let $\tilde{Q}_n(x)$ denote the cumulative distribution function for this distribution.

The main result of this section, stated below, shows that \tilde{Q}_n is asymptotically $\tilde{Q}_{n,\infty}$, where for $\ell \geq 2$ let

$$\tilde{Q}_{n,\ell} := \sum_{r=0}^{\ell-2} n^{-r/2} P_r(-\Phi_{0,V_n}, \{\bar{\chi}_\nu\}). \quad (30)$$

and we will define the signed measure $P_r(-\Phi_{0,V_n}, \{\bar{\chi}_\nu\})$ below after fixing a few notations. For convenience, the density of $\tilde{Q}_{n,\ell}$ is denoted by $Q_{n,\ell}$ while the density of \tilde{Q}_n is denoted by Q_n .

First, let W be the standard Gaussian vector in \mathbb{R}^d , then for any covariance matrix V , $V^{1/2}W$ will be the Gaussian random variable in \mathbb{R}^d with mean zero and covariance V . Let $\phi_{0,V}$ denote the density of its distribution and let $\Phi_{0,V}$ denote the cumulative distribution function. If V is the identity matrix then we simply write ϕ and Φ , respectively. Note that this is consistent with our definition of Φ at the beginning Section 2.9.

Secondly, recall that the cumulants of a random vector X in \mathbb{R}^d are the coefficients in the following (multiple) power series expansion

$$\log \mathbf{E}[e^{z \cdot X}] = \sum_{\nu \in \mathbf{N}^d} \frac{\chi_\nu z^\nu}{\nu!}, \quad z \in \mathbf{C}^d. \quad (31)$$

Given that X has mean zero, it is standard that the cumulant χ_ν is bounded above by the $|\nu|$ th moment of X . In our situation, using independence of $X_{n,1}, \dots, X_{n,n}$, it follows that the cumulants of S_n are the sum of the corresponding cumulants of $X_{n,1}, \dots, X_{n,n}$. Let $\bar{\chi}_\nu := \chi_\nu(S_n)/n$, then $\bar{\chi}_\nu$ is also the average cumulant of $X_{n,1}, \dots, X_{n,n}$.

Now, note that cumulants of $V_n^{1/2}W$ matches with the cumulants of S_n/\sqrt{n} for any $|\nu| \leq 2$, at the same time the higher order cumulants of $V_n^{1/2}W$ vanish thanks to symmetries of centered Gaussian. Therefore,

$$\begin{aligned} \log \mathbf{E}[e^{z \cdot (S_n/\sqrt{n})}] &= \log \mathbf{E}[e^{z \cdot (V_n^{1/2}W)}] + \sum_{\nu \in \mathbf{N}^d: |\nu| \geq 3} (n\bar{\chi}_\nu) \frac{z^\nu}{\nu!} n^{-|\nu|/2} \\ &= \log \mathbf{E}[e^{z \cdot V_n^{1/2}W}] + \sum_{\ell \geq 1} \left(\sum_{\nu \in \mathbf{N}^d: |\nu| = \ell+2} \bar{\chi}_\nu \frac{z^\nu}{\nu!} \right) n^{-\ell/2}. \end{aligned}$$

Letting $\bar{\chi}_\ell(z) = \ell! \sum_{\nu \in \mathbf{N}^d: |\nu| = \ell} \bar{\chi}_\nu z^\nu$ for all $z \in \mathbf{C}^d$, we obtain

$$\begin{aligned} \mathbf{E}[e^{z \cdot (S_n/\sqrt{n})}] / \mathbf{E}[e^{z \cdot V_n^{1/2}W}] &= \exp\left[\sum_{\ell \geq 1} \frac{\bar{\chi}_{\ell+2}(z)}{(\ell+2)!} n^{-\ell/2}\right] \\ &= \sum_{m \geq 0} \frac{1}{m!} \left(\sum_{\ell \geq 1} \frac{\bar{\chi}_{\ell+2}(z)}{(\ell+2)!} n^{-\ell/2} \right)^m = \sum_{\ell \geq 0} \tilde{P}_\ell n^{-\ell/2}, \end{aligned}$$

where \tilde{P}_ℓ is obtained by grouping terms of the same order $n^{-\ell/2}$. It is clear that \tilde{P}_ℓ depends only on z and the average cumulants $\bar{\chi}_\nu, |\nu| \leq \ell+2$. We'll write $\tilde{P}_\ell(z, \{\bar{\chi}_\nu\})$ to stress this dependence. Replacing z by iz , we obtain the following expansion for the characteristic function of S_n/\sqrt{n} :

$$\mathbf{E}[e^{iz \cdot (S_n/\sqrt{n})}] = \mathbf{E}[e^{iz \cdot V_n^{1/2}W}] \sum_{\ell \geq 0} \tilde{P}_\ell(iz, \{\bar{\chi}_\nu\}) n^{-\ell/2}.$$

Now, let $D = (D_1, \dots, D_n)$ be the partial derivative operator and let $\tilde{P}_\ell(-D, \{\bar{\chi}_\nu\})$ be the differential operator obtained by formally replacing all occurrences of iz by $-D$ inside $\tilde{P}_\ell(iz, \{\bar{\chi}_\nu\})$. The signed measure $P_\ell(-\Phi_{0, V_n}, \{\bar{\chi}_\nu\})$ in the definition (30) of $\tilde{Q}_{n, \ell}$ now can be defined: it has the following density with respect to the Lebesgue measure:

$$P_\ell(-\phi_{0, V_n}, \{\bar{\chi}_\nu\})(x) := \left(\tilde{P}_\ell(-D, \bar{\chi}_\nu) \phi_{0, V_n} \right)(x).$$

For convenience of notation, for each $\ell > 0$, let $\rho_\ell = \frac{1}{n} \sum \mathbf{E} \|X_i\|_2^\ell$ and

$$M_\ell(f) := \sup_{x \in \mathbb{R}^d} \frac{|f(x)|}{1 + \|x\|_2^\ell}$$

for any measurable function f .

Theorem 4.1. *Let S_n be defined as above using (28) where we assume that the distribution of Y_j satisfies $\mathbf{E} \|Y_j\|_2^{\ell+d+1} < \infty$ for some $\ell \geq 4$. Let f be measurable such that $M_\ell(f) < \infty$.*

Suppose that:

- (1) all eigenvalues of V_n are larger than a constant $\sigma > 0$ independent of n ;
- (2) the parameters s, t in the definition of $C_n(1), \dots, C_n(n)$ satisfy Condition 2 for some sufficiently small τ .

Then the following estimate holds for $\varepsilon = n^{-C_*}$ where C_* is any given positive constant:

$$\begin{aligned} & \left| \int f(x) d\tilde{Q}_n - \int f(x) d\tilde{Q}_{n, \ell} \right| \\ & \leq CM_\ell(f) (n^{-(\ell-1)/2} + e^{-n^{-\tau_*}} + e^{-cn}) + \bar{\omega}_f(2\varepsilon : \sum_{r=0}^{\ell+d-2} n^{-r/2} P_r(-\phi_{0, V_n} : \{\bar{\chi}_\nu\})) \end{aligned}$$

where

$$\bar{\omega}_f(\varepsilon : \phi) = \int \left(\sup_{y \in B(x, \varepsilon)} f(y) - \inf_{y \in B(x, \varepsilon)} f(y) \right) d\phi(x),$$

and the implied constant C depends on $\{\rho_k, k \leq \ell\}$, σ , C_* , and the implicit constants from Condition 2, but not on f .

Notice that the verification of condition (1) on the invertibility of V_n follows from [8, Appendix C].

The general strategy of our proof follows the approach in [9], here we focus on the main differences while trying to keep the exposition self-contained. Here our goal is not about proving the sharpest possible version for Theorem 4.1 in terms of the number of bounded moments for Y_i , rather our aim is to present a simpler argument (compared to [9]) at the sake of a more stringent moment assumption.

Before starting the proof, we include some estimates that will be useful in the proof.

Lemma 4.2. *Let $l, c_0 > 0$ be any given constants. Assume that $\mathbf{E}\|X_{n,k}\|_2^{\ell+1} = O(1)$ uniformly over n and $k = 1, \dots, n$. Then for some sufficiently small $c_1 > 0$ the following holds for all $\|\eta\|_2 < c_1 n^{1/2}$ and all multi-index α :*

$$\begin{aligned} D_\eta^\alpha \left(\mathbf{E}[e^{i\eta \cdot (S_n/\sqrt{n})}] (\mathbf{E}[e^{i\eta \cdot V_n^{1/2} W}])^{-1} - \sum_{r=0}^{\ell-2} \tilde{P}_r(i\eta, \{\bar{\chi}_\nu\}) n^{-r/2} \right) \\ \leq C n^{-(\ell-1)/2} e^{c_0 \|\eta\|_2^2} (\|\eta\|_2^{\ell+1-|\alpha|} + \|\eta\|_2^{3\ell+1-|\alpha|}). \end{aligned}$$

Here the implicit constant may depend on c_0, ℓ, α and $\bar{\chi}_0, \dots, \bar{\chi}_{\ell+1}$

Proof. For brevity we will write $\tilde{P}_r(i\eta)$ as a shortcut of $\tilde{P}_r(i\eta, \{\bar{\chi}_\nu\})$.

Let

$$f_{\eta, \ell}(u) := \exp(g_{\eta, \ell}(u)) := \exp\left(\sum_{m=1}^{\ell-2} \frac{\bar{\chi}_{m+2}(i\eta)}{(m+2)!} u^m\right).$$

We first show that for any multi-index α

$$\left| D_\eta^\alpha \left(f_{\eta, \ell} \left(\frac{1}{\sqrt{n}} \right) - \sum_{r=0}^{\ell-2} \tilde{P}_r(i\eta) n^{-r/2} \right) \right| \leq C n^{-(\ell-1)/2} (\|\eta\|_2^{\ell+1-|\alpha|} + \|\eta\|_2^{3\ell-3-|\alpha|}) e^{c_0 \|\eta\|_2^2} \quad (32)$$

for all $\|\eta\|_2 \leq c_1 n^{1/2}$ and $c_1 > 0$ is sufficiently small.

Let $u \in \mathbb{R}$ (that may depend on n). As a function of $u \in \mathbb{R}$, the polynomial $\sum_{r=0}^{\ell-2} \tilde{P}_r(i\eta) u^r$ is the Taylor approximation of degree $\ell - 2$ for $f_{\eta, \ell}(u)$. Now, if $\|\eta\|_2 u \ll 1$ then

$$|g_{\eta, \ell}(u)| = O(\|\eta\|_2^3 |u|) < c_0 \|\eta\|_2^2$$

and similarly $|g_{\eta, \ell}^{(k)}(u)| = O(\|\eta\|_2^{k+2})$. Thus, using the chain rule and the generalized Leibniz rule, we may bound

$$\begin{aligned} |f_{\eta, \ell}^{(\ell-1)}(u)| &\leq C e^{g_{\eta, \ell}(u)} \left(\sum_{j_1+2j_2+\dots=\ell-1} \prod_{k \geq 1} |g_{\eta, \ell}^{(k)}(u)|^{j_k} \right) \\ &\leq C e^{g_{\eta, \ell}(u)} \left(\sum_{j_1+2j_2+\dots=\ell-1} \prod_{k \geq 1} \|\eta\|_2^{(2+k)j_k} \right) \\ &= O((\|\eta\|_2^{\ell+1} + \|\eta\|_2^{3\ell-3}) e^{c_0 \|\eta\|_2^2}). \end{aligned}$$

(Here the implicit constant may depend on c_0, ℓ, α and $\bar{\chi}_0, \dots, \bar{\chi}_{\ell+1}$.) We obtain, assuming $\|\eta\|_2 < c_1 |u|^{-1}$,

$$\left| f_{\eta, \ell}(u) - \sum_{r=0}^{\ell-2} \tilde{P}_r(i\eta) u^r \right| \leq C |u|^{\ell-1} (\|\eta\|_2^{\ell+1} + \|\eta\|_2^{3\ell-3}) e^{c_0 \|\eta\|_2^2}.$$

We now let $u = \frac{1}{\sqrt{n}}$. Using analytic dependence on η of $f_{\eta, \ell}(\frac{1}{\sqrt{n}}) - \sum_{r=0}^{\ell-2} \tilde{P}_r(i\eta) n^{-r/2}$ and Cauchy's theorem for analytic functions, we obtain (32).

Now, it remains to show that

$$\left| D_\eta^\alpha \left(f_{\eta, \infty} \left(\frac{1}{\sqrt{n}} \right) - f_{\eta, \ell} \left(\frac{1}{\sqrt{n}} \right) \right) \right| \leq C n^{-(\ell-1)/2} (\|\eta\|_2^{\ell+1-|\alpha|} + \|\eta\|_2^{3\ell+1-|\alpha|}) e^{2c_0 \|\eta\|_2^2} \quad (33)$$

As before it suffices to show the case $\alpha = 0$ of this estimate, and then the desired estimate follows from an application of Cauchy's theorem. Now, since $|f_{\eta, \ell}(\frac{1}{\sqrt{n}})| \leq e^{c_0 \|\eta\|_2^2}$ as proved above, it suffices to show that

$$|f_{\eta, \infty} \left(\frac{1}{\sqrt{n}} \right) f_{\eta, \ell} \left(\frac{1}{\sqrt{n}} \right)^{-1} - 1| \leq C n^{-(\ell-1)/2} (\|\eta\|_2^{\ell+1} + \|\eta\|_2^{3\ell+1}) e^{c_0 \|\eta\|_2^2}. \quad (34)$$

Let $u \in (-2, 2)$ and let $h(u) := g_{u\eta, \infty}(\frac{1}{\sqrt{n}}) - g_{u\eta, \ell}(\frac{1}{\sqrt{n}})$. It is clear that the first ℓ derivatives with respect to $u \in \mathbb{R}$ of h all vanish at $u = 0$. Thus, using the chain rule and the generalized Leibniz rule, it follows that the first ℓ derivatives with respect to u of $f_{u\eta, \infty}(\frac{1}{\sqrt{n}}) f_{u\eta, \ell}(\frac{1}{\sqrt{n}})^{-1} - 1$ also vanish at $u = 0$. With $|u| = O(1)$, we obtain

$$\begin{aligned} |f_{u\eta, \infty} \left(\frac{1}{\sqrt{n}} \right) f_{u\eta, \ell} \left(\frac{1}{\sqrt{n}} \right)^{-1} - 1| &\leq C \left| \left(\frac{d}{du} \right)^{\ell+1} \left(f_{u\eta, \infty} \left(\frac{1}{\sqrt{n}} \right) f_{u\eta, \ell} \left(\frac{1}{\sqrt{n}} \right)^{-1} - 1 \right) \right| \\ &= e^{h(u)} O \left(\sum_{j_1 + 2j_2 + \dots = \ell+1} \prod_{k \geq 1} |h^{(k)}(u)|^{j_k} \right) \end{aligned} \quad (35)$$

Now, as the first ℓ derivatives of h all vanish at $u = 0$, for any $k \leq \ell + 1$ we have (with $|u| = O(1)$)

$$|h^{(k)}(u)| \leq C \sup_{|t| \leq |u|} |h^{(\ell+1)}(t)| = C \sup_{|t| \leq |u|} \left| \left(\frac{d}{dt} \right)^{(\ell+1)} g_{t\eta, \infty} \left(\frac{1}{\sqrt{n}} \right) \right|.$$

By definition we have

$$g_{u\eta, \infty} \left(\frac{1}{\sqrt{n}} \right) = -\log(\mathbf{E} e^{i(\eta \cdot V_n^{1/2} W)u}) + \sum_{j=1}^n \log(\mathbf{E} e^{i(\eta \cdot X_{n,j})u/\sqrt{n}})$$

Using $\mathbf{E} \|X_{n,j}\|_2 = O(1)$ we obtain

$$|\mathbf{E} e^{i\eta \cdot X_{n,j} u/\sqrt{n}} - 1| = O(\|\eta\|_2/\sqrt{n}),$$

therefore using the given assumption we obtain $|\mathbf{E} e^{i\eta \cdot X_{n,j} u/\sqrt{n}}| > 1/2$. Consequently, using the chain rule and the Leibniz rule we have

$$\left| \left(\frac{d}{du} \right)^{\ell+1} \log(\mathbf{E} e^{i\eta \cdot X_{n,j} u/\sqrt{n}}) \right| = O \left(\sum_{j_1 + 2j_2 + \dots = \ell+1} \prod_{k \geq 1} \mathbf{E} [|n^{-1/2} \eta \cdot X_{n,j}|^{j_k}] \right) = O(n^{-(\ell+1)/2} \|\eta\|_2^{\ell+1}).$$

Since $\log(\mathbf{E} e^{i(\eta \cdot V_n^{1/2} W)u})$ is quadratic with respect to u and $\ell \geq 4$, we obtain

$$\left| \left(\frac{d}{du} \right)^{\ell+1} g_{u\eta, \infty} \left(\frac{1}{\sqrt{n}} \right) \right| = O(n^{-(\ell-1)/2} \|\eta\|_2^{\ell+1}).$$

Consequently,

$$|h^{(k)}(u)| \leq C n^{-(\ell-1)/2} \|\eta\|_2^{\ell+1} \leq C c_1^{(\ell-1)/2} \|\eta\|_2^2,$$

in particular by choosing c_1 small we can ensure that $|h(u)| \leq c_0 \|\eta\|_2^2$. Therefore, using (35), we obtain

$$\begin{aligned} |f_{u\eta, \infty} \left(\frac{1}{\sqrt{n}} \right) f_{u\eta, \ell} \left(\frac{1}{\sqrt{n}} \right)^{-1} - 1| &\leq C e^{c_0 \|\eta\|_2^2} \sum_{j_1 + 2j_2 + \dots = \ell+1} \prod_{k \geq 1} (n^{-(\ell-1)/2} \|\eta\|_2^{\ell+1})^{j_k} \\ &\leq C e^{c_0 \|\eta\|_2^2} (n^{-(\ell-1)/2} \|\eta\|_2^{\ell+1} + (n^{-(\ell-1)/2} \|\eta\|_2^{\ell+1})^{\ell+1}) \\ &\leq C e^{c_0 \|\eta\|_2^2} n^{-(\ell-1)/2} \|\eta\|_2^{\ell+1} (1 + \|\eta\|_2^{2\ell}) \end{aligned}$$

We then set $u = 1$ to obtain the desired estimate. \square

As a corollary, we obtain

Corollary 4.3. *Assume that $\mathbf{E}|X_{n,k}|^{\ell+1} = O(1)$ uniformly over n and $k = 1, \dots, n$. Assume that the eigenvalues of V_n are bounded below by some positive constant independent of n . Then for some sufficiently small constants $c_0, c_1 > 0$ the following holds for all $\|\eta\|_2 < c_1 n^{1/2}$ and all multi-index α :*

$$\begin{aligned} & D_\eta^\alpha \left(\mathbf{E}[e^{i\eta \cdot (S_n/\sqrt{n})}] - \mathbf{E}[e^{i\eta \cdot V_n^{1/2} W}] \sum_{r=0}^{\ell-2} \tilde{P}_r(i\eta) n^{-r/2} \right) \\ & \leq C n^{-(\ell-1)/2} e^{-c_0 \|\eta\|_2^2} (\|\eta\|_2^{\ell+1-|\alpha|} + \|\eta\|_2^{3\ell+1+|\alpha|}). \end{aligned}$$

This corollary follows from the fact that $\mathbf{E}[e^{i\eta \cdot V_n^{1/2} W}]$ is $e^{-c\langle \eta, V_n^{-1} \eta \rangle}$ for some $c > 0$, so with $c_0 > 0$ sufficiently small one has

$$|D_\eta^\alpha \mathbf{E}[e^{i\eta \cdot V_n^{1/2} W}]| \leq e^{-2c_0 \|\eta\|_2^2} (\|\eta\|_2^{|\alpha|} + 1),$$

and combining these estimates with the Leibniz rule we obtain the desired conclusion.

Proof. (of Theorem 4.1) We now begin the proof of the main estimate. Let

$$\varepsilon = \varepsilon_n = n^{-C_*}. \quad (36)$$

For convenience, denote

$$\tilde{H}_n = \tilde{Q}_n - \tilde{Q}_{n,\ell},$$

and let H_n be its density. As usual the characteristic function of H_n is $\widehat{H}_n(\eta) = \int_{\mathbb{R}^d} e^{it \cdot \eta} \tilde{H}_n(dt)$.

Let \tilde{K} be a probability measure supported inside the unit ball $B(0, 1) = \{x \in \mathbb{R}^d : \|x\| \leq 1\}$ (whose density is denoted by K) such that its characteristic function $\widehat{\tilde{K}}(\eta)$ satisfies

$$|D^\alpha \widehat{\tilde{K}}(\eta)| = O(e^{-\|\eta\|_2^{1/2}}), \quad |\alpha| \leq \ell + d + 1. \quad (37)$$

Such a measure could be constructed using elementary arguments, see for instance [9, Section 10]. We then let \tilde{K}_ε be the ε -dilation of \tilde{K} , namely $\tilde{K}_\varepsilon(A) = \tilde{K}(\varepsilon^{-1}A)$ and $\varepsilon^{-1}A := \{x/\varepsilon : x \in A\}$ for all measurable A . Note that \tilde{K}_ε is a probability measure on $B(0, \varepsilon)$ and it satisfies the dilated version of (37).

We will be using the following simple identity: for any two measures μ_1 and μ_2 of bounded variation, $|\mu_1|(\mathbb{R}^d), |\mu_2|(\mathbb{R}^d) < \infty$, and any bounded f , it holds that

$$\int \int f(x+y) d\mu_1(x) d\mu_2(y) = \int f(t) (d\mu_1 * d\mu_2)(t).$$

Now, for each $x \in B(0, \varepsilon)$ we have $f(y) \leq \sup_{z \in B(0, \varepsilon)} f(x+y+z)$, therefore using nonnegativity of $d\tilde{Q}_n$ we obtain

$$\begin{aligned} \int f(y) d\tilde{H}_n(y) &= \int_{x \in B(0, \varepsilon)} \int f(y) d\tilde{H}_n(y) d\tilde{K}_\varepsilon(x) \\ &\leq \int \left(\int \sup_{z \in B(0, \varepsilon)} f(x+y+z) d\tilde{Q}_n(y) - \int f(y) d\tilde{Q}_{n,\ell}(y) \right) d\tilde{K}_\varepsilon(x) \\ &= \int \int \sup_{z \in B(0, \varepsilon)} f(x+y+z) dH_n(y) dK_\varepsilon(x) + \\ &\quad + \int \int \left(\sup_{z \in B(0, \varepsilon)} f(x+y+z) - f(y) \right) d\tilde{Q}_{n,\ell}(y) d\tilde{K}_\varepsilon(x) \\ &= \int \sup_{z \in B(0, \varepsilon)} f(t+z) (H_n * K_\varepsilon)(t) dt + \int \left(\sup_{z \in B(0, \varepsilon)} f(x+y+z) - f(y) \right) d\tilde{Q}_{n,\ell}(y) d\tilde{K}_\varepsilon(x) \\ &\leq M_\ell(f) \int (1 + \|t\|_2 + \varepsilon)^\ell |H_n * K_\varepsilon|(t) dt + \int \int \left(\sup_{B(y, 2\varepsilon)} f(t) - \inf_{B(y, 2\varepsilon)} f(t) \right) |d\tilde{Q}_{n,\ell}(y)| d\tilde{K}_\varepsilon(x) \\ &\leq C_\ell M_\ell(f) \int (1 + \|t\|_2)^\ell |H_n * K_\varepsilon|(t) dt + \bar{\omega}_f(2\varepsilon : |\tilde{Q}_{n,\ell}|). \end{aligned}$$

By applying the above estimate for $-f$ in place of f , it follows immediately that $|\int f d\widehat{H}_n|$ is bounded above by the same right hand side. By standard Sobolev embedding estimates for Fourier transforms, we have

$$\begin{aligned} \int (1 + \|t\|_2)^\ell |H_n * K_\varepsilon|(t) dt &= O\left(\max_{0 \leq |\alpha| \leq d + \ell + 1} \int |D^\alpha \widehat{H}_n * \widehat{K}_\varepsilon(\eta)| d\eta\right) \\ &= O\left(\max\left\{\int |D^\alpha(\widehat{H}_n)(\eta) D^\beta(\widehat{K}_\varepsilon)(\eta)| d\eta : |\alpha| + |\beta| \leq \ell + d + 1\right\}\right). \end{aligned}$$

Using (37) we have $D^\alpha \widehat{K}_\varepsilon(\eta) = O(1)$ for all $|\alpha| \leq \ell + d + 1$. While this estimate is fairly generous, it is good enough to control the contribution of small η in the integrals. More specifically, let $B_n^2 = V_n^{-1}$, then by the given assumption the eigenvalues of B_n are $O(1)$, so $\mathbf{E}\|B_n X_{n,k}\|_2^{\ell+d+1} = O(1)$. For some $c_1 > 0$ sufficiently small, using Corolary 4.3, we obtain

$$\begin{aligned} \int_{\|\eta\|_2 \leq c_1 \sqrt{n}} |D^\alpha \widehat{H}_n(\eta) D^\beta \widehat{K}_\varepsilon(\eta)| d\eta &= O\left(\int_{\|\eta\|_2 \leq c_1 \sqrt{n}} |D^\alpha \widehat{H}_n(\eta)| d\eta\right) \\ &= O(n^{-(\ell+d-1)/2}). \end{aligned}$$

We now consider the range $\|\eta\|_2 \geq c_1 \sqrt{n}$. We estimate

$$\begin{aligned} \int_{\|\eta\|_2 \geq c_1 \sqrt{n}} |D^\alpha \widehat{H}_n(t) D^\beta \widehat{K}_\varepsilon| d\eta &\leq \int_{\|\eta\|_2 \geq c_1 \sqrt{n}} |D^\alpha \widehat{Q}_n(t) D^\beta \widehat{K}_\varepsilon| d\eta \\ &\quad + \int_{\|\eta\|_2 \geq c_1 \sqrt{n}} |D^\alpha \left(\sum_{r=0}^{\ell-2+d} n^{-r/2} P_r(i\eta : \{\chi_{\nu,n}\})\right) \exp(-1/2\langle \eta, B_n \eta \rangle)| d\eta, \end{aligned}$$

and it is clear that the second term can be controlled by $O(e^{-cn})$ thanks to the Gaussian decay of $\exp(-1/2\langle \eta, B_n \eta \rangle)$.

Let $\phi_i(\eta) = \mathbf{E}e^{i\eta \cdot X_i}$. Then for $|\alpha| \leq \ell + d + 1$ we have $D_\eta^\alpha(\phi_i(\eta/\sqrt{n})) = n^{-|\alpha|/2} O(\mathbf{E}\|X_{n,i}\|_2^{|\alpha|}) = O(1)$. Thus,

$$|D^\alpha \widehat{Q}_n(\eta)| = |D^\alpha \left(\prod_{i=1}^n \phi_i\left(\frac{\eta}{\sqrt{n}}\right)\right)| = O\left(\sum_{\gamma_1 + \dots + \gamma_n = \alpha} \left| \prod_{i=1, \gamma_i=0}^n \phi_i\left(\frac{\eta}{\sqrt{n}}\right) \right|\right),$$

while we also have

$$|D^\beta \widehat{K}_\varepsilon(\eta)| = O(\varepsilon^{|\beta|} e^{-(\varepsilon\|\eta\|_2)^{1/2}}) = O(e^{-(\varepsilon\|\eta\|_2)^{1/2}}).$$

Thus, it remains to control, for each $(\gamma_1, \dots, \gamma_n)$ with $|\gamma_1| + \dots + |\gamma_n| \leq \ell + d + 1$ and each $r > 0$ independent of n :

$$\begin{aligned} J_\gamma(n, \varepsilon) &= \int_{\|\eta\|_2 \geq r\sqrt{n}} \left| \prod_{i=1, \gamma_i=0}^n \phi_i\left(\frac{\eta}{\sqrt{n}}\right) \right| e^{-(\varepsilon\|\eta\|_2)^{1/2}} d\eta \\ &= n^{d/2} \int_{\|\eta\|_2 \geq r} \left| \prod_{i=1, \gamma_i=0}^n \phi_i(\eta) \right| e^{-(\varepsilon\sqrt{n}\|\eta\|_2)^{1/2}} d\eta \\ &= n^{d/2} \int_{\|\eta\|_2 \geq r} \left| \prod_{i=1, \gamma_i=0}^n \phi_i(\eta) \right| e^{-(n^{-C_*+1/2}\|\eta\|_2)^{1/2}} d\eta. \end{aligned}$$

Clearly it suffices to consider $r \leq \|\eta\|_2 \leq n^{C_*-1/2+\tau}$ because the integral for $\|\eta\|_2 \geq n^{C_*-1/2+\tau}$ is extremely small. Again, because α is fixed, by throwing away from the set $\{\mathbf{v}_i\}$ a fixed number of elements, let us assume that $\alpha = 0$ for simplicity ³. By Theorem 2.5 for sufficiently large n we have

$$\left| \prod_i \phi_i(\eta) \right| \leq e^{-n^{-\tau*}}.$$

Thus we just shown that, with $\varepsilon = n^{-C_*}$,

$$J_\gamma(n, \varepsilon) = O(e^{-n^{-\tau*}}).$$

³In the general case $\alpha \neq 0$ we use Theorem 10.6 instead of Theorem 2.5.

Putting the bounds together, we obtain the desired estimate:

$$|\int f d\tilde{H}_n| \leq CM_\ell(f)(n^{-(\ell-1)/2} + e^{-n^{-\tau_*}} + e^{-cn}) + \bar{\omega}_f(2\varepsilon : \sum_{r=0}^{\ell-2+d} n^{-r/2} P_r(-\phi_{0,V_n} : \{\bar{\chi}_\nu\}).$$

□

4.4. A useful corollary. Below we consider a consequence of Theorem 4.1 that will be convenient for our proof of Theorem 1.4 in subsequent sections.

With $Y_k = (y_{k1}, y_{k2})$ where y_{ij} are iid with mean zero and variance one, we recall the definition of $P_n(t, Y)$ from (3). Let $G_k = (g_{k1}, g_{k2})$ where g_{ij} are iid standard Gaussian. Recall the definition of $S_n(Y, s, t)$ from (28) and let $S_n(G, s, t)$ be its Gaussian analogue.

Clearly $\tilde{P}_0 = 1$ and by explicit computation we have

$$\tilde{P}_1(z, \{\bar{\chi}_\nu\}) = \sum_{|\nu|=3} \frac{\bar{\chi}_\nu}{\nu!} z^\nu, \quad \tilde{P}_2(z, \{\bar{\chi}_\nu\}) = \frac{\bar{\chi}_4(z)}{24} + \frac{\bar{\chi}_3^2(z)}{72}. \quad (38)$$

For convenience of notation let $e_j = (\dots, 0, 1, 0, \dots) \in \mathbb{R}^d$ where 1 is in the j th coordinate. Using (38) we obtain

$$\begin{aligned} P_1(-\phi, \{\bar{\chi}_\nu\}) &= \sum_{|\nu|=3} \frac{\bar{\chi}_\nu}{\nu!} (-D)^\nu \phi(x) = \\ &= \left[\frac{1}{6} \sum_{j=1}^4 \bar{\chi}_{3e_j} (x_j^3 - 3x_j) + \frac{1}{2} \sum_{i \neq j} \bar{\chi}_{2e_i + e_j} (x_i^2 x_j - x_j) + \sum_{i < j < k} \bar{\chi}_{e_i + e_j + e_k} x_i x_j x_k \right] \phi(x) \\ &= \left[\frac{1}{6} \sum_{j=1}^4 \bar{\chi}_{3e_j} h_3(x_j) + \frac{1}{2} \sum_{i \neq j} \bar{\chi}_{2e_i + e_j} h_2(x_i) h_1(x_j) + \sum_{i,j,k} \bar{\chi}_{e_i + e_j + e_k} h_1(x_i) h_1(x_j) h_1(x_k) \right] \phi(x), \end{aligned}$$

where $h_k(x) = (-1)^k e^{x^2/2} \frac{\partial^k}{\partial x^k} e^{-x^2/2}$ ($k = 0, 1, 2, \dots$) are the (one dimensional) Hermite polynomials.

Now for any multi-index $\alpha = (\alpha_1, \dots, \alpha_\ell) \in \{1, \dots, d\}^\ell$, we let $|\alpha| = \ell$ and let $n_j(\alpha) = |\{i : \alpha_i = j\}|$ for each $j = 1, \dots, d$. We then define

$$H_\alpha(x_1, \dots, x_d) := \prod h_{n_1}(x_1) \dots h_{n_d}(x_d). \quad (39)$$

For a random vector $Z = (Z_1, \dots, Z_d)$ as usual let $Z^\alpha = \prod_{j=1}^d Z_j^{\alpha_j}$. With $X = (X_{n,1}, \dots, X_{n,n})$ define

$$\Delta_\alpha(X_{n,k}) = \mathbf{E} X_{n,k}^\alpha - \mathbf{E} G_{n,k}^\alpha, \quad (40)$$

$$c_n(\alpha, X) := \frac{1}{n} \sum_{k=1}^n \Delta_\alpha(X_{n,k}) \quad (41)$$

$$\Gamma_{n,1}(X, x) := \frac{1}{6} \sum_{|\alpha|=3} c_n(\alpha, X) H_\alpha(x). \quad (42)$$

Note that if α' is a permutation of α then $H_{\alpha'} = H_\alpha$. Furthermore using (31) and explicit computations it follows that $\chi_\nu(X) = \mathbf{E}[X^\nu]$ for all $|\nu| = 2, 3$ if X is a random vector in \mathbb{R}^d with mean $\mathbf{E}X = 0$. Thus, for all distinct i, j, k ,

$$\bar{\chi}_{3e_j} = c_n((j, j, j), X), \quad \bar{\chi}_{2e_i + e_j} = 0 = c_n((i, i, j), X), \quad \chi_{e_i + e_j + e_k} = 0 = c_n((i, j, k), X).$$

Using these observations, we obtain

$$P_1(-\phi_{0,V_n}, \{\bar{\chi}_\nu\}) = \Gamma_{n,1}(X, x) \phi_{0,V_n}(x).$$

We also define

$$\Gamma_{n,2}(X, x) = \Gamma'_{n,2} + \Gamma''_{n,2} \quad (43)$$

where

$$\Gamma'_{n,2}(X, x) = \frac{1}{24} \sum_{|\beta|=4} c_n(\beta, X) H_\beta(x)$$

and

$$\Gamma''_{n,2}(X, x) = \frac{1}{72} \sum_{|\rho|=3} \sum_{|\beta|=3} c_n(\beta, X) c_n(\rho, X) H_{\beta,\rho}(x).$$

Via explicit computations, it can also be checked that

$$P_2(-\phi_{0,V_n}, \{\bar{X}_\nu\}) = \Gamma_{n,2}(X, x) \phi_{0,V_n}(x).$$

Finally, recall the definition of $\tilde{Q}_{n,2}$ from (30), which has density

$$Q_{n,2}(X, x) = 1 + n^{-1/2} P_1(-\Phi_{0,V_n}, \{\bar{X}_\nu\}) + n^{-1} P_2(-\Phi_{0,V_n}, \{\bar{X}_\nu\}).$$

It follows that

Fact 4.5.

$$Q_{n,2}(X, x) = 1 + \frac{1}{\sqrt{n}} \Gamma_{n,1}(X, x) + \frac{1}{n} \Gamma_{n,2}(B_n X, x).$$

Now by applying Theorem 4.1 and then swallow higher order terms in the Edgeworth expansion into the error terms (resulting into $O(n^{-3/2})$, keeping the first three terms), we obtain the following corollary.

Theorem 4.6. *With the same assumption as in Theorem 4.1 the following holds for $\varepsilon = n^{-C_*}$ (and C_* is any given positive constant):*

$$\begin{aligned} & |\mathbf{E}(f(S_n(s, t, Y))) - \mathbf{E}(f(V_n^{1/2} W) Q_{n,2}(X, W))| \\ & \leq C n^{-3/2} + C M_\ell(f) (n^{-(\ell-1)/2} + e^{-n^{-\tau_*}} + e^{-cn}) + \bar{\omega}_f(2\varepsilon : \sum_{r=0}^{\ell+d-2} n^{-r/2} P_r(-\phi_{0,V_n} : \{\bar{X}_\nu\})). \end{aligned}$$

where W is the standard Gaussian vector in \mathbb{R}^d .

5. PROOF OF PROPOSITION 2.10 : ASYMPTOTIC KAC-RICE FORMULA

We will show the following more precise statement.

Proposition 5.1. *Let ℓ be a fixed positive integer. Let δ be as in (12). Assume that η has mean zero and variance one and $\mathbf{E}|\eta|^{M_0} < \infty$ for sufficiently large M_0 . Assume that s, t satisfy Condition 2. Then for any $\varepsilon = n^{-C_*}$ (where $C_* > 0$ is any absolute constant), we have*

$$|\mathbf{E}F_\delta(P_n(t, Y)) - \mathbf{E}F_\delta(P_n(t, G))| \leq \frac{C}{n^{1/2}} + C\delta^{-1}(n^{-(\ell-1)/2} + e^{-n^{\tau_*}} + e^{-cn} + \varepsilon) \quad (44)$$

and

$$\begin{aligned} & \left| \mathbf{E}\Phi_\delta\left(\frac{1}{\sqrt{n}} S_n(t, Y)\right) - \mathbf{E}\Phi_\delta\left(\frac{1}{\sqrt{n}} S_n(t, G)\right) - \frac{1}{n} \mathbf{E}\left[\Phi_\delta(I_2(\lambda)^{1/2} W_2) \Gamma_{n,2}(I_2(\lambda)^{-1/2} X_n(t, Y), W_2)\right] \right| \\ & \leq \frac{C}{n^{3/2}} + \frac{1}{n} r_n(t, \Phi_\delta) + C\delta^{-1}(n^{-(\ell-1)/2} + e^{-n^{\tau_*}} + e^{-cn} + \varepsilon) \end{aligned} \quad (45)$$

and

$$\begin{aligned} & \left| \mathbf{E}\Psi_\delta\left(\frac{1}{\sqrt{n}} S_n(s, t, Y)\right) - \mathbf{E}\Psi_\delta\left(\frac{1}{\sqrt{n}} S_n(s, t, G)\right) - \frac{1}{n} \mathbf{E}\left[\Psi_\delta(I_4(\lambda)^{1/2} W_4) \Gamma_{n,2}(I_4(\lambda)^{-1/2} X_n(s, t, Y), W_4)\right] \right| \\ & \leq \frac{C}{n^{3/2}} + \frac{1}{n} r_n(s, t, \Psi_\delta) + C\delta^{-2}(n^{-(\ell-1)/2} + e^{-n^{\tau_*}} + e^{-cn} + \varepsilon), \end{aligned} \quad (46)$$

where I_2 and I_4 are any invertible diagonal matrices and W, W_2, W_4 are standard Gaussian vectors in \mathbb{R}, \mathbb{R}^2 and \mathbb{R}^4 respectively, and where the implied constants are allowed to depend on the M_0 -moment of η , on the constants in Conditions 1 and 2, and on a lower bound of the least singular values of $V_n(t), V_n(s, t)$ and I_2, I_4 . Furthermore, we have the following bounds

$$r_n(t, \Phi_\delta) = O(\|V_n(t) - I_2\|_2) \text{ and } r_n(s, t, \Psi_\delta) = O(\|V_n(s, t) - I_4\|_2).$$

Note that if we apply the above theorem for $\varepsilon = n^{-(\ell-1)/2}$ and for sufficiently large ℓ (for instance $\ell = 16$ would suffice), then all the error bounds are absorbed into $O(\frac{1}{n^{3/2}})$, and hence proving Proposition 2.10.

We now discuss the proof. We first note that if f is an even function then using the fact that the standard Gaussian distribution is symmetric and the fact that Hermite polynomials of odd degrees are odd functions we obtain

$$\mathbf{E}[f(V_n^{1/2}W)\Gamma_{n,1}(V_n^{-1/2}X, W)] = 0.$$

In our applications below the functions f are indeed even therefore we could ignore the contribution of $\Gamma_{n,1}$ in the estimates.

Now, recall (28) and recall that $\delta = n^{-5}$ and $\varepsilon = n^{-C_*}$ for some given constant $C_* > 0$. Recall also the definitions of the even functions $F_\delta : \mathbb{R} \rightarrow \mathbb{R}_+, \Phi_\delta : \mathbb{R}^2 \rightarrow \mathbb{R}_+, \Psi_\delta : \mathbb{R}^4 \rightarrow \mathbb{R}_+$ from Subsection 2.6. Now, using standard integration by parts (for details see [8, Eq. 3.23]) we may rewrite the Gaussian part in a more canonical form: with f being either F_δ, Ψ_δ or Φ_δ , and with W being either W_2 or W_4 we have

$$\mathbf{E}(f(V_n^{1/2}W)\Gamma_{n,2}(V_n^{-1/2}X, W)) = \mathbf{E}(f(I_d^{1/2}W)\Gamma_{n,2}(I_d^{-1/2}X, W)) + r_n(f) \quad (47)$$

with I_d being a diagonal matrix with diagonal entries at least σ and here

$$|r_n(f)| \leq C\|V_n - I_d\|_2.$$

For this proof, we will only work with Ψ_δ and prove (46) as (44) and (45) are similar and simpler. By Fact 4.5 and by (47), to prove Proposition 5.1 for this $f = \Psi_\delta$ it suffices to show

$$\left| \mathbf{E}f\left(\frac{1}{\sqrt{n}}S_n(s, t, Y) - \mathbf{E}(f(V_n^{1/2}W)Q_{n,2}(X, W))\right) \right| \leq Cn^{-3/2} + C\delta^{-2}(n^{-(\ell-1)/2} + e^{-n^{-\tau_*}} + e^{-cn} + \varepsilon). \quad (48)$$

Proof. (of Proposition 5.1) Let $\lambda \in (0, 1)$ and let $\varphi_\lambda : \mathbb{R} \rightarrow [0, 1]$ be a $C^\infty(\mathbb{R})$ function with support inside $[-\delta, \delta]$ such that

- (i) $\varphi_\lambda(x) = \delta^{-1}$ for $|x| \leq \delta(1 - \lambda)$.
- (ii) $|\varphi_\lambda^{(k)}(x)| = O(\delta^{-(k+1)}\lambda^{-k})$ for any $k \geq 0$.

Let $\tilde{\varphi}_\lambda : \mathbb{R}^4 \rightarrow \mathbb{R}$ be defined by $\tilde{\varphi}_\lambda(x) = \varphi_\lambda(x_1)\varphi_\lambda(x_3)$. Let

$$f_\lambda(x_1, x_2, x_3, x_4) = |x_2||x_4|\tilde{\varphi}_\lambda(x).$$

Then f_λ is locally Lipschitz, and its derivative (defined almost everywhere) satisfies

$$|\nabla f_\lambda| \leq C\frac{1}{\delta^2\lambda}(1 + |x|)^4.$$

Recall that $\bar{\omega}_f(\varepsilon : \phi) = \int(\sup_{y \in B(x, \varepsilon)} f(y) - \inf_{y \in B(x, \varepsilon)} f(y))\phi(x)dx$, and ϕ is the density of a Gaussian vector. Consequently, for any polynomial $p(x)$ with bounded degree and bounded coefficients we have

$$\bar{\omega}_{f_\lambda}(\varepsilon : p(x)\phi_{0, V_n}(x)) = O(\lambda^{-1}\delta^{-2}\varepsilon).$$

Here we are implicitly using the fact that the eigenvalues of V_n are bounded above by $O(1)$, which should follow from the fact that the singular values of $C_n(k)$ are bounded: they are bounded by the Hilbert Schmidt norm, which is bounded since the entries of $C_n(k)$ are bounded.

Note that one could write

$$\sum_{r=0}^{\ell+d-2} n^{-r/2} P_r(-\phi_{0, V_n} : \{\bar{\chi}_\nu\}) = p(x) \phi_{0, V_n}(x)$$

for some polynomial p with degree at most $d + \ell$ and coefficients bounded by the first $d + \ell$ moments of the random coefficients Y_1, \dots, Y_n of P_n . Therefore

$$\bar{\omega}_{f_\lambda}(2\varepsilon : \sum_{r=0}^{\ell+d-2} n^{-r/2} P_r(-\phi_{0, V_n} : \{\bar{\chi}_\nu\})) = O(\lambda^{-1} \delta^{-2} \varepsilon). \quad (49)$$

We will also use the following elementary estimate: given any a_1, \dots, a_n deterministic and η_1, \dots, η_n independent with mean 0 and bounded 4th moment, the following holds

$$\mathbf{E}|a_1 \eta_1 + \dots + a_n \eta_n|^4 \leq C(a_1^2 + \dots + a_n^2)^2.$$

Indeed, thanks to independent and the mean zero property, the left hand side is

$$\begin{aligned} \mathbf{E}(\sum_{i,j,k,l} a_i a_j a_k a_l \eta_i \eta_j \eta_k \eta_l) &= \sum_i a_i^4 \mathbf{E} \eta_i^4 + O(\sum_{i < j} a_i^2 a_j^2 \mathbf{E} \eta_i^2 \eta_j^2) \\ &= O(a_1^2 + \dots + a_n^2)^2. \end{aligned}$$

In particular,

$$\mathbf{E}|P'_n(t, Y)|^4 = n^{-2} O((\sum_{j=1}^n \sin^2(jt/n) + \cos^2(jt/n))^2) = O(1).$$

We next proceed to conclude Proposition 5.1 for $f = \Psi_\delta$. Recall that ϕ_{0, V_n} denotes the density of $V_n^{1/2} W$. Then using Hölder's inequality and Theorem 2.2 we obtain

$$\begin{aligned} &\mathbf{E}(f - f_\lambda)(\frac{1}{\sqrt{n}} S_n(s, t, Y)) \\ &\leq C \delta^{-2} \mathbf{E}[|P'_n(t, Y) P'_n(s, Y)| \mathbf{1}_{\|P_n(t, Y) - \delta\| \leq \lambda \delta} \mathbf{1}_{\|P_n(s, Y) - \delta\| \leq \lambda \delta}] \\ &\leq C \delta^{-2} (\mathbf{E}|P'_n(t, Y)|^4)^{1/4} (\mathbf{E}|P'_n(s, Y)|^4)^{1/4} \mathbf{P}(\|P_n(t, Y) - \delta\| \leq \delta \lambda)^{1/4} \mathbf{P}(\|P_n(s, Y) - \delta\| \leq \delta \lambda)^{1/4} \\ &\leq C \delta^{-2} (n \delta \lambda)^{2/5} \quad (\text{assume bounded 4th moment for coefficients of } P_n \text{ and Corollary 3.1}) \\ &\leq C \delta^{-2} \lambda^{2/5}. \end{aligned}$$

By Theorem 4.6 and (49), applying for $\varepsilon^{7/2}$ in place of ε , we obtain

$$\begin{aligned} &|\mathbf{E} f_\lambda(\frac{1}{\sqrt{n}} S_n(s, t, Y)) - \mathbf{E}(f_\lambda(V_n^{1/2} W) Q_{n,2}(X, W))| \\ &= O(n^{-3/2}) + M_\ell(f_\lambda) O(n^{-(\ell-1)/2} + e^{-n^{-\tau^*}} + e^{-cn}) + \bar{\omega}_{f_\lambda}(2\varepsilon^{7/2} : \sum_{r=0}^{\ell+d-2} n^{-r/2} P_r(-\phi_{0, V_n} : \{\bar{\chi}_\nu\})) \\ &= O(n^{-3/2}) + \delta^{-2} O(n^{-(\ell-1)/2} + e^{-n^{-\tau^*}} + e^{-cn}) + O(\delta^{-2} \lambda^{-1} \varepsilon^{7/2}), \end{aligned}$$

where we note that $M_\ell(f_\lambda) = O(\delta^{-2})$. Note that as a special case, this bound also holds for the Gaussian case. Consequently,

$$\begin{aligned} &|\mathbf{E} f(\frac{1}{\sqrt{n}} S_n(s, t, Y)) - \mathbf{E}(f(V_n^{1/2} W) Q_{n,2}(X, W))| \\ &\leq |\mathbf{E}(f - f_\lambda)(\frac{1}{\sqrt{n}} S_n(s, t, Y))| + |\mathbf{E}(f - f_\lambda)((V_n^{1/2} W) Q_{n,2}(X, W))| + \\ &\quad + |\mathbf{E} f_\lambda(\frac{1}{\sqrt{n}} S_n(s, t, Y)) - \mathbf{E}(f_\lambda(V_n^{1/2} W) Q_{n,2}(X, W))| \\ &\leq C n^{-3/2} + C \delta^{-2} (n^{-(\ell-1)/2} + e^{-n^{-\tau^*}} + e^{-cn}) + \lambda^{-1} \varepsilon^{7/2} + \lambda^{2/5} \end{aligned}$$

We then take $\lambda = \varepsilon^{5/2}$ and obtain the desired estimate as in (48). \square

6. COMPLETING THE PROOF OF PROPOSITION 2.8

Recall the definition of $D_{n,\varepsilon}$ from (18).

In this proof, we follow the proof of [8, Lemma 5.1]; the only differences are that the Lebesgue measure of our set $D_{n,\varepsilon}$ is $4 + o(1)$ times larger and that we use our version of the Edgeworth expansion, Proposition 2.10. The proof consists of several steps which we use a similar enumeration as in [8].

Step 1: Making use of the Edgeworth expansion in Proposition 2.10. In this step, we shall apply the formulas in Proposition 2.10. We shall choose the diagonal matrices I_2 and I_4 that appear in Proposition 2.10 to be the limit of $V_n(t)$ and $V_n(s, t)$. More precisely, by letting I_2 be the 2×2 diagonal matrix with diagonal entries $\lambda_1 = 1$, $\lambda_2 = 1/3$, and I_4 be the 4×4 diagonal matrix with diagonal entries $\lambda_1 = \lambda_3 = 1$, $\lambda_2 = \lambda_4 = 1/3$, one can check that for all s, t satisfying Condition 2,

$$\lim_{n \rightarrow \infty} \|V_n(t) - I_2\|_2 = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|V_n(s, t) - I_4\|_2 = 0.$$

Applying Proposition 2.10, we obtain the expansion

$$v_n(s, t, Y) - v_n(s, t, G) = \frac{1}{n} \gamma_n(s, t) + \bar{R}_n(s, t) \quad (50)$$

where the γ_n has the main terms and the \bar{R}_n contains all the error terms and their product with the main terms in Proposition 2.10. In particular,

$$\begin{aligned} \gamma_n(s, t) &= \mathbf{E} \left[\Psi_\delta(I_4^{1/2} W_4) \Gamma_{n,2}(I_4^{-1/2} X_n(s, t, Y), W_4) \right] \\ &\quad - \mathbf{E} \left[\Phi_\delta(I_2^{1/2} W_2) \right] \mathbf{E} \left[\Phi_\delta(I_2^{1/2} W_2) \Gamma_{n,2}(I_2^{-1/2} X_n(t, Y), W_2) \right] \\ &\quad - \mathbf{E} \left[\Phi_\delta(I_2^{1/2} W_2) \right] \mathbf{E} \left[\Phi_\delta(I_2^{1/2} W_2) \Gamma_{n,2}(I_2^{-1/2} X_n(s, Y), W_2) \right] \\ &= \mathbf{E} \left[\Psi_{\delta_n}(I_4^{1/2} W_4) \right. \\ &\quad \left. \times \left[\Gamma_{n,2}(I_4^{-1/2} X_n(s, t, Y), W_4) - \Gamma_{n,2}(I_2^{-1/2} X_n(t, Y), W_2') - \Gamma_{n,2}(I_2^{-1/2} X_n(s, Y), W_2'') \right] \right] \end{aligned}$$

where W_2', W_2'' are independent standard Gaussian vectors in \mathbb{R}^2 and $W_4 = (W_2', W_2'')$ is a standard Gaussian vector in \mathbb{R}^4 . We recall that

$$\Gamma_{n,2}(X, x) = \Gamma'_{n,2}(X, x) + \Gamma''_{n,2}(X, x)$$

which is the sum of the following fourth moment corrector

$$\Gamma'_{n,2}(X, x) = \frac{1}{24} \sum_{|\beta|=4} c_n(\beta, X) H_\beta(x)$$

and the following combined third moment corrector

$$\Gamma''_{n,2}(X, x) = \frac{1}{72} \sum_{|\rho|=3} \sum_{|\beta|=3} c_n(\beta, X) c_n(\rho, X) H_{\beta,\rho}(x). \quad (51)$$

We recall the definition of $c_n(\cdot, X)$ in (41) and the polynomials $H_\beta, H_{\beta,\rho}$ in (39).

Denote by $\gamma'_n(t, s)$ and $\gamma''_n(t, s)$ the corresponding quantities when replacing $\Gamma_{n,2}(X, x)$ by $\Gamma'_{n,2}(X, x)$ and $\Gamma''_{n,2}(X, x)$, respectively, in the definition of $\gamma_n(t, s)$. Proposition 2.8 is reduced to showing the following:

$$\lim_n \frac{1}{n^2} \int_{D_{n,\varepsilon}} \gamma''_n(s, t) ds dt = 0, \quad (52)$$

$$\lim_n \frac{1}{n^2} \int_{D_{n,\varepsilon}} \gamma'_n(s, t) ds dt = \frac{\mathbf{E} \xi^4 - 3}{15} + O(\varepsilon), \quad (53)$$

and

$$\lim_n \frac{1}{n} \int_{D_{n,\varepsilon}} \bar{R}_n(s, t) ds dt = 0. \quad (54)$$

To see (54), we simply note that for almost every $(s, t) \in [0, \pi]^2$, $\bar{R}_n(ns, nt) = o(\frac{1}{n})$ as $n \rightarrow \infty$. This is mainly because (via examination), as $n \rightarrow \infty$, the re-scaled covariance matrix $V_n(nt, ns)$ converges to the diagonal matrix $(1, 1/3, 1, 1/3)$ if $\frac{t}{\pi}, \frac{s}{\pi}, \frac{t-s}{\pi}, \frac{t+s}{\pi}$ are irrational.

The remaining identities will be established in the next steps. For convenience of notation, in the rest of the section for each multi-index α (of dimension 3 or 4) we let

$$c_n(\alpha, s, t) = c_n\left(\alpha, I_4^{-1/2} X_n(s, t, Y)\right), \quad c_n(\alpha, t) = c_n\left(\alpha, I_2^{-1/2} X_n(t, Y)\right),$$

and define $c_n(\alpha, s)$ similarly. Also, we will denote by $\alpha - 2$ the adjusted multi-index where each index in α will be subtracted by 2.

Step 2: Proving (52).

For each multi-index $\rho, \beta \in \{1, 2, 3, 4\}^3$, define $\gamma''_{n,\rho,\beta}(s, t) = \mathbf{E}\{\Psi_{\delta_n}(I_4^{1/2} W_4) \Delta c_n(\beta, \rho, s, t)\}$ where

$$\Delta c_n(\beta, \rho, s, t) = \begin{cases} c_n(\rho, s, t) c_n(\beta, s, t) H_{\beta,\rho}(W_4) - c_n(\rho, t) c_n(\beta, t) H_{\beta,\rho}(W_2'), & \text{if } \beta, \rho \in \{1, 2\}^3; \\ c_n(\rho, s, t) c_n(\beta, s, t) H_{\beta,\rho}(W_4) - c_n(\rho - 2, s) c_n(\beta - 2, s) H_{\beta,\rho}(W_2''), & \text{if } \beta, \rho \in \{3, 4\}^3; \\ c_n(\rho, s, t) c_n(\beta, s, t) H_{\beta,\rho}(W_4), & \text{otherwise.} \end{cases}$$

It is clear that $\gamma''_n(s, t) = \sum_{|\beta|=3} \sum_{|\rho|=3} \gamma''_{n,\rho,\beta}(s, t)$.

Step 2, the cancellations: We recall the following cancellations that were observed in [8].

- (First cancellation) If $\beta, \rho \in \{1, 2\}^3$ then by examination we have $c_n(\beta, s, t) = c_n(\beta, t)$, $c_n(\rho, s, t) = c_n(\rho, t)$, and $H_{\beta,\rho}(W_4) = H_{\beta,\rho}(W_2')$. Consequently, $\Delta c_n(\beta, \rho, s, t) = 0$ and thus

$$\gamma''_{n,\rho,\beta}(s, t) = 0.$$

Similarly, if $\beta, \rho \in \{3, 4\}^3$ then $\Delta c_n(\beta, \rho, s, t) = 0$ and so $\gamma''_{n,\rho,\beta}(s, t) = 0$.

- (Second cancellation) We now consider those (β, α) not part of above cancellation scenarios, i.e. where there is a mixed of elements from $\{1, 2\}$ and elements from $\{3, 4\}$. Then if an index $j \in \{1, 2, 3, 4\}$ appears an odd number of times inside (β, ρ) then

$$\mathbf{E}\{\Psi_{\delta_n}(I_4^{1/2} W_4) H_{\beta,\rho}(W_4)\} = 0,$$

since $\Psi_{\delta_n}(I_4^{1/2} W_4)$ is an even function of $W_{4,j}$ (the j th coordinate of W_4) and $H_{\beta,\rho}(W_4)$ is an odd function of $W_{4,j}$. Consequently, in this case we also have $\gamma''_{n,\rho,\beta}(s, t) = 0$.

For the remaining (β, ρ) , for almost every $(s, t) \in [0, \pi]^2$ (with respect to the Lebesgue measure) we have

$$c_n(\beta, ns, nt) c_n(\rho, ns, nt) \rightarrow 0, \quad \text{as } n \rightarrow \infty. \quad (55)$$

Indeed, the ‘‘mixed’’ nature of (β, ρ) implies that one of β, ρ will be mixed. Without loss of generality assume that β is mixed, say $\beta = \{i, i, j\}$ where $i \leq 2 < j$. Then via examination $c_n(\beta, s, t)$ is an average (over k) of term of the following type:

$$\sum_{\ell_1, \ell_2, \ell_3=1}^2 A_k(ns, nt, \ell_1, \ell_2, \ell_3) \mathbf{E}(Y_{k,\ell_1} Y_{k,\ell_2} Y_{k,\ell_3})$$

whre $A_k(s, t, \ell_1, \ell_2, \ell_3)$ is a product of two terms from $C_n(k, t)$ and one term from $C_n(k, s)$, and $Y_{k,1}, Y_{k,2}$ are the coordinates of Y_k . Now $\mathbf{E}(Y_{k,\ell_1} Y_{k,\ell_2} Y_{k,\ell_3})$ is constant with respect to k (due to iid - although this is not essential, it suffices to assume convergence as $k \rightarrow \infty$ of this term), the desired convergence follows from the limit $\frac{1}{n} \sum_k A_k(ns, nt, \ell_1, \ell_2, \ell_3) \rightarrow 0$, which can be checked using elementary trigonometric identities, provided that $\frac{t}{\pi}, \frac{s}{\pi}, \frac{t+s}{\pi}, \frac{t-s}{\pi}$ are all irrational.

Now, since the c_n are uniformly bounded, (55) and Lebesgue's dominated convergence theorem imply that

$$\lim_n \frac{1}{n^2} \int_{D_{n,\varepsilon}} \gamma''_{n,\rho,\beta}(s,t) ds dt = \lim_n \int_0^\pi \int_0^\pi 1_{D_{n,\varepsilon}}(nu, nv) \gamma''_{n,\rho,\beta}(nu, nv) dudv = 0.$$

which completes the proof of (52).

Step 3: Proving (53). This follows from similar reasoning as in Step 2. First, using cancellations similar to Step 2, we also obtain

$$\gamma'_n(s,t) = \frac{1}{24} \sum_\alpha \mathbf{E} \left(\Psi_{\delta_n}(I_4^{1/2}W) H_\alpha(W) \right) c_n(\alpha, s, t)$$

where the sum runs over all mixed α of the form $\alpha = (i, i, j, j)$ with $i \in \{1, 2\}$ and $j \in \{3, 4\}$, and their permutations. It is clear that $\gamma'_n(s, t) = O(1)$ uniformly over n and s, t .

Thus,

$$\frac{1}{n^2} \int_{D_{n,\varepsilon}} \gamma'(s,t) ds dt = O(\varepsilon) + \frac{1}{n^2} \frac{1}{24} \sum_\alpha \mathbf{E} \left(\Psi_{\delta_n}(I_4^{1/2}W) H_\alpha(W) \right) \int_{D_{n,\varepsilon}} c_n(\alpha, s, t) ds dt. \quad (56)$$

Now, if α is a permutation of (i, i, j, j) with $i \leq 2 < j$, by examination and using trigonometric identities as in Step 2, it follows that $c_n(\alpha, s, t)$ is an average of sums $\sum_{\ell_1, \ell_2, \ell_3, \ell_4=1}^2 A_k(s, t, \ell_1, \dots, \ell_4) \mathbf{E}(Y_{k,\ell_1} Y_{k,\ell_2} Y_{k,\ell_3} Y_{k,\ell_4})$, $k = 1, \dots, n$. Here $A_k(s, t, \ell_1, \dots, \ell_4)$ is a product of two terms from $C_n(k, s)$ and two terms from $C_n(k, t)$. Via examination, (for details see for instance the appendix in [8]), one could show that

Lemma 6.1. *For almost every $(s, t) \in [0, \pi]^2$ (with respect to the Lebesgue measure) it holds that*

$$\lim_n c_n(\alpha, ns, nt) = \frac{2 \cdot 3^{i+j-4} (\mathbf{E}\xi^4 - 3)}{2i + 2j - 4} \quad (57)$$

On the other hand, it is clear that, for the same α ,

$$\lim_n \mathbf{E} \left(\Psi_{\delta_n}(I_4^{1/2}W) H_\alpha(W) \right) = \frac{1}{3\pi^2} (-1)^{i+j}.$$

Plugging in these limits to (56), we obtain (53).

7. PROOF OF LEMMA 2.7

We recall the definition of the sub-intervals I_k in (17) and the sets $D_{n,\varepsilon}$ and \mathcal{D} in (18). Let $N_{I_k}(Y)$ be the number of roots in I_k .

Since $N_n(Y) = \sum_k N_{I_k}(Y)$, we have

$$\text{Var} N_n(Y) - \text{Var} N_n(G) = 2\mathcal{V}_1 + \mathcal{V}_2 \quad (58)$$

where

$$\mathcal{V}_1 := \sum_{(k,p) \in \mathcal{D}} [\text{Cov}(N_{I_k}(Y), N_{I_p}(Y)) - \text{Cov}(N_{I_k}(G), N_{I_p}(G))], \quad (59)$$

and

$$\mathcal{V}_2 := 2 \sum_{(k,p) \notin \mathcal{D}, k < p} [\text{Cov}(N_{I_k}(Y), N_{I_p}(Y)) - \text{Cov}(N_{I_k}(G), N_{I_p}(G))] + \sum_k [\text{Var} N_{I_k}(Y) - \text{Var} N_{I_k}(G)]. \quad (60)$$

Therefore, Lemma 2.7 follows from the following two results concerning \mathcal{V}_1 and \mathcal{V}_2 , respectively.

Lemma 7.1 (asymptotic estimate for \mathcal{V}_1). *We have*

$$\mathcal{V}_1 = \int_{D_{n,\varepsilon}} (v_n(t, s, Y) - v_n(t, s, G)) dsdt + R_{n,\varepsilon} \quad (61)$$

where

$$\lim_n \frac{R_{n,\varepsilon}}{n} = 0.$$

We recall the definition of v_n in (16):

$$v_n(s, t, Y) = \text{Cov}(P'_n(s, Y)\mathbf{1}_{|P_n(s, Y)| < \delta_n}, P'_n(t, Y)\mathbf{1}_{|P_n(t, Y)| < \delta_n}) = \mathbf{E}\Psi_{\delta_n} \left(\frac{1}{\sqrt{n}} S_n(s, t, Y) \right) - \mathbf{E}\phi_{\delta_n}(s, Y)\mathbf{E}\phi_{\delta_n}(t, Y).$$

Lemma 7.2 (\mathcal{V}_2 is negligible). *There exists a constant c such that*

$$\mathcal{V}_2 \ll n^{1-c}.$$

Remark 7.3. *As we shall show in the proof, for Lemma 7.2, we only need to assume that the random variables y_{ij} are independent (not necessarily identically distributed) with mean 0, variance 1, and bounded $(2 + \varepsilon_0)$ -moment, namely $\mathbf{E}|y_{ij}|^{2+\varepsilon_0} < C$ for some positive constants ε_0, C and for all i, j .*

The rest of this section is devoted to the proof of these lemmas.

7.4. Proof of Lemma 7.1. For this proof, we adapt the proof of [8, Lemma 4.2] using the new inequalities that we have obtained.

Recall that $\delta = n^{-5}$ in this proof. Let

$$\delta_{a,b,Y} := \min_{t \in [a,b]} \{|P_n(a, Y)|, |P_n(b, Y)|, |P_n(t, Y)| + |P'_n(t, Y)|\}.$$

By the Kac-Rice formula, for any interval $[a, b]$, the number of zeros of $P_n(\cdot, Y)$ in the interval $[a, b]$ is given by

$$N_n([a, b], Y) = \int_a^b |P'_n(t, Y)| \mathbf{1}_{|P_n(t, Y)| \leq \delta} \frac{dt}{2\delta} \quad \text{if } \delta \leq \delta_{a,b,Y}. \quad (62)$$

To prove Lemma 7.1, it suffices to show that for any $(k, p) \in \mathcal{D}$,

$$\mathbf{E}N_{I_k}(Y)N_{I_p}(Y) = \int_{I_k \times I_p} \mathbf{E}\phi_\delta(t, Y)\phi_\delta(s, Y) dt ds + O(\varepsilon_{k,p}) \quad (63)$$

and

$$\mathbf{E}N_{I_k}(Y)\mathbf{E}N_{I_p}(Y) = \int_{I_k \times I_p} \mathbf{E}\phi_\delta(t, Y)\mathbf{E}\phi_\delta(s, Y) dt ds + O(\varepsilon_{k,p}) \quad (64)$$

where

$$\sum_{(k,p) \in \mathcal{D}} \varepsilon_{k,p} = o(n).$$

Since the proof of (63) and (64) are similar, we shall now only prove (63). By the Kac-Rice formula (62),

$$\mathbf{E}N_{I_k}(Y)N_{I_p}(Y)\mathbf{1}_{\delta \leq \min\{\delta_{I_k,Y}, \delta_{I_p,Y}\}} = \int_{I_k \times I_p} \mathbf{E}\phi_\delta(t, Y)\phi_\delta(s, Y)\mathbf{1}_{\delta \leq \min\{\delta_{I_k,Y}, \delta_{I_p,Y}\}} dt ds.$$

Thus, by setting

$$\varepsilon'_{k,p} = \mathbf{E}N_{I_k}(Y)N_{I_p}(Y)\mathbf{1}_{\delta > \min\{\delta_{I_k,Y}, \delta_{I_p,Y}\}} \quad (65)$$

and

$$\varepsilon''_{k,p} = \int_{I_k \times I_p} \mathbf{E}\phi_\delta(t, Y)\phi_\delta(s, Y)\mathbf{1}_{\delta > \min\{\delta_{I_k,Y}, \delta_{I_p,Y}\}} dt ds, \quad (66)$$

we are left to show that

$$\sum_{(k,p) \in \mathcal{D}} \varepsilon'_{k,p} = o(n) \quad (67)$$

and

$$\sum_{(k,p) \in \mathcal{D}} \varepsilon''_{k,p} = o(n). \quad (68)$$

For (67), using the fact that the number of real roots inside $[-n\pi, n\pi]$ is at most $2n$ deterministically, we get that

$$\varepsilon'_{k,p} \ll n^2 \mathbf{P}(\delta > \min\{\delta_{I_k, Y}, \delta_{I_p, Y}\}) \leq n^2 \mathbf{P}(\delta > \delta_{I_k, Y}) + n^2 \mathbf{P}(\delta > \delta_{I_p, Y}). \quad (69)$$

Let a, b be the endpoints of I_k . We have

$$\mathbf{P}(\delta > \delta_{I_k, Y}) \leq \mathbf{P}(|P_n(a, Y)| < \delta) + \mathbf{P}(|P_n(b, Y)| < \delta) + \mathbf{P}\left(\min_{t \in I_k} |P_n(t, Y)| + |P'_n(t, Y)| < \delta\right). \quad (70)$$

Since $(k, p) \in \mathcal{D}$ and since any (s, t) that satisfies Condition 2, it is necessary that both s and t satisfy Condition 1. In other words, $I_k \subset \mathcal{G}$ in Theorem 3.2. Applying this theorem, we get

$$\mathbf{P}\left(\min_{t \in I_k} |P_n(t, Y)| + |P'_n(t, Y)| < \delta\right) \ll n^{-4+\varepsilon} \quad (71)$$

where we recall that in Theorem 3.2, $\frac{1}{\sqrt{n}} S_n(Y, t) = (P_n(t, Y), P'_n(t, Y))$ as defined in (8).

Applying Corollary 3.1 with $M_0 = 4$, for all t satisfying Condition 1, we have

$$\mathbf{P}(|P_n(t, Y)| < \delta) \ll n^{-3-\varepsilon}. \quad (72)$$

Applying (72) for $t = a, b$, we get

$$\mathbf{P}(|P_n(a, Y)| < \delta) + \mathbf{P}(|P_n(b, Y)| < \delta) \ll n^{-3-\varepsilon}.$$

Plugging this together with (71) to (70), we obtain

$$\mathbf{P}(\delta > \delta_{I_k, Y}) \ll n^{-3-\varepsilon}.$$

Similarly for I_p . Thus, from (69), we have $\varepsilon'_{k,p} \ll n^{-1-\varepsilon}$ which gives (67).

For (68), we argue similarly using the observation in [8, Inequality (4.2)] that, deterministically,

$$\int_{I_k} \phi_\delta(t, Y) dt \leq 1 + N_{I_k}(Y) \leq 2n + 1 \quad \text{and} \quad \int_{I_p} \phi_\delta(t, Y) dt \leq 1 + N_{I_p}(Y) \leq 2n + 1.$$

7.5. Proof of Lemma 7.2. As in Remark 7.3, in this subsection, we only assume that the random variables y_{ij} are independent (not necessarily identically distributed) with mean 0, variance 1, and bounded $(2 + \varepsilon_0)$ -moment. To prove Lemma 7.2, we shall use the following result.

Lemma 7.6. *There exists a constant c such that for all k, p that are not necessarily distinct,*

$$\mathbf{E}N_{I_k}(Y)N_{I_p}(Y) - \mathbf{E}N_{I_k}(G)N_{I_p}(G) \ll n^{-2c}, \quad (73)$$

$$\mathbf{E}N_{I_k}(Y) - \mathbf{E}N_{I_k}(G) \ll n^{-2c}, \quad (74)$$

and

$$\mathbf{E}N_{I_k}(Y) \ll 1, \mathbf{E}N_{I_k}(G) \ll 1. \quad (75)$$

Proof of Lemma 7.2. Assuming Lemma 7.6, we have for all k, p ,

$$\mathbf{E}N_{I_k}(Y) \cdot \mathbf{E}N_{I_p}(Y) - \mathbf{E}N_{I_k}(G) \cdot \mathbf{E}N_{I_p}(G) \ll n^{-2c}$$

where we used the triangle inequality, (74), and (75). Combining this with (73), we obtain

$$\text{Cov}(N_{I_k}(Y), N_{I_p}(Y)) - \text{Cov}(N_{I_k}(G), N_{I_p}(G)) \ll n^{-2c} \quad \text{for all } k, p, \quad (76)$$

and in particular when $k = p$, we have

$$\text{Var}N_{I_k}(Y) - \text{Var}N_{I_k}(G) \ll n^{-2c}.$$

Plugging these estimates into (60), we obtain

$$\mathcal{V}_2 \ll n^{-2c} \#\{(k, p) \notin \mathcal{D}\}. \quad (77)$$

Observe that for each k , the number of p such that $(k, p) \notin \mathcal{D}$ is $O_\varepsilon(n^{11\tau})$. Indeed, by the definition of \mathcal{D} and Condition 2, for each $l_1, l_2 \neq 0$ with $|l_1|, |l_2| \leq n^\tau$, it suffices to show that the number of p such that there exist $t \in I_k, s \in I_p$ with

$$\left| \frac{l_1 t + l_2 s}{n} \right|_{\mathbb{R}/\pi\mathbb{Z}} \leq n^{-1+8\tau} \quad (78)$$

is $O_\varepsilon(n^{9\tau})$. The inequality (78) is equivalent to

$$l_1 t + l_2 s \in [na - n^{8\tau}, na + n^{8\tau}] \quad \text{for some } a \in \pi\mathbb{Z}, |a| \leq 2\pi n^\tau;$$

in other words,

$$s \in \frac{1}{l_2} ([na - n^{8\tau}, na + n^{8\tau}] - l_1 I_k) \quad \text{for some } a \in \pi\mathbb{Z}, |a| \leq 2\pi n^\tau.$$

For each a , the right-hand side is contained in an interval of length $O(n^{8\tau})$ which corresponds to $O_\varepsilon(n^{8\tau})$ values of p . Taking union bound over $O(n^\tau)$ choices of a gives the stated claim.

Using this observation, the right-hand side of (77) is $O(n^{-2c+1+11\tau}) = O(n^{1-c})$ by choosing τ to be sufficiently small compared to c . \square

To prove Lemma 7.6, we denote the roots of $P_n(\cdot, Y)$ by $\zeta_1(Y), \dots, \zeta_n(Y)$ and the roots of $P_n(\cdot, G)$ by $\zeta_1(G), \dots, \zeta_n(G)$. We shall use the following result in [52].

Theorem 7.7. [52, Theorem 3.3] *There exist constants c, C' such that for any real numbers x_1, x_2 and for any function $F : \mathbb{R}^2 \rightarrow \mathbb{R}$ supported on $[x_1 - 1, x_1 + 1] \times [x_2 - 1, x_2 + 1]$ with continuous derivatives up to order 8 and $\|\nabla^a F\|_\infty \leq 1$ for all $0 \leq a \leq 8$, we have*

$$\left| \mathbf{E} \sum F(\zeta_i(Y), \zeta_j(Y)) - \mathbf{E} \sum F(\zeta_i(G), \zeta_j(G)) \right| \leq C' n^{-c},$$

where the first sum runs over all pairs $(\zeta_i(Y), \zeta_j(Y)) \in \mathbb{R}^2$ of the roots of $P_n(\cdot, Y)$ and the second sum runs over all pairs $(\zeta_i(G), \zeta_j(G)) \in \mathbb{R}^2$ of the roots of $P_n(\cdot, G)$.

Proof of Lemma 7.6. Let x_k, x_p be the midpoint of I_k, I_p , respectively. Let $\gamma = n^{-s}$ for $s = c/100$ and c be the constant in Theorem 7.7.

The inequalities in (75) follow from [52, Theorem 3.6]. For the rest of the proof, we show (73). The proof of (74) is similar (and simpler).

We approximate the indicator function on the interval $[-\varepsilon/2, \varepsilon/2]$ by a smooth function ϕ satisfying

$$\mathbf{1}_{[-\varepsilon/2+\gamma, \varepsilon/2-\gamma]} \leq \phi \leq \mathbf{1}_{[-\varepsilon/2, \varepsilon/2]}$$

and

$$\|\nabla^a \phi\|_\infty \ll \gamma^{-a}, \quad \forall 0 \leq a \leq 8.$$

Let

$$F(x, y) := \phi(x + x_k)\phi(y + x_p).$$

Let

$$M_k(Y) := \left(\sum_{i=1}^n \phi(\zeta_i(Y) - x_k) \right), \quad M_p(Y) := \left(\sum_{i=1}^n \phi(\zeta_i(Y) - x_p) \right).$$

Denote by $M_k(G)$ and $M_p(G)$ the corresponding terms for the Gaussian case, i.e., with $\zeta_i(G)$ in place of $\zeta_i(Y)$. Applying Theorem 7.7 to the function $\gamma^8 F$, we obtain

$$\left| \mathbf{E} \sum F(\zeta_i(Y), \zeta_j(Y)) - \mathbf{E} \sum F(\zeta_i(G), \zeta_j(G)) \right| \leq C' n^{-c},$$

and so

$$|\mathbf{E} M_k(Y) M_p(Y) - \mathbf{E} M_k(G) M_p(G)| \leq C' n^{-c/2}, \quad (79)$$

We shall show that

$$\mathbf{E} N_{I_k}(Y) N_{I_p}(Y) - \mathbf{E} M_k(Y) M_p(Y) = O\left(n^{-s/10}\right). \quad (80)$$

The same argument applied to the Gaussian case will show that

$$\mathbf{E} N_{I_k}(G) N_{I_p}(G) - \mathbf{E} M_k(G) M_p(G) = O\left(n^{-s/10}\right). \quad (81)$$

Combining (79), (80), and (81), we obtain (73) as desired (by choosing the c in (73) to be $s/10$).

To prove (80), by Holder's inequality, we have

$$(\mathbf{E} N_{I_k}(Y) N_{I_p}(Y) - \mathbf{E} M_k(Y) N_{I_p}(Y))^2 \ll \mathbf{E} (N_{I_k}(Y) - M_k(Y))^2 \mathbf{E} N_{I_p}^2(Y). \quad (82)$$

Let $N_\gamma(Y)$ be the number of roots of $P_n(\cdot, Y)$ in the union of the intervals $[x_k + \varepsilon/2 - \gamma, x_k + \varepsilon/2]$, $[x_k - \varepsilon/2, x_k - \varepsilon/2 - \gamma]$, $[x_p + \varepsilon/2 - \gamma, x_p + \varepsilon/2]$, and $[x_p - \varepsilon/2, x_p - \varepsilon/2 - \gamma]$. We observe that $N_\gamma(Y)$ is at least $|N_{I_k}(Y) - M_k(Y)|$.

By [52, Formula (28), page 32], there exists an $x \in I_k$ such that

$$\mathbf{P}\left(\log |P_n(x, Y)| \leq -n^{s/10}\right) \ll n^{-100}.$$

By [52, Lemma 9.4],

$$\mathbf{P}\left(\log \max_{x \in B(x, \varepsilon)} |P_n(z, Y)| \geq n^{s/10}\right) \ll n^{-100}.$$

By Jensen's inequality (see, for example, [52, Formula (8), page 22]), under the event that $\log |P_n(x, Y)| \geq -n^{s/10}$ and $\log \max_{x \in B(x, \varepsilon)} |P_n(z, Y)| \leq n^{s/10}$, we have $N_{I_k}(Y) \leq n^{s/10}$. Thus,

$$\mathbf{P}\left(N_{I_k}(Y) \geq n^{s/10}\right) \ll n^{-100}. \quad (83)$$

And by [52, Lemma 8.6],

$$\mathbf{P}(N_\gamma(Y) \geq 2) \ll n^{-3s/2}. \quad (84)$$

When $N_\gamma(Y) < 2$, we have $N_\gamma(Y)^2 = N_\gamma(Y)$. Thus,

$$\begin{aligned} \mathbf{E} N_\gamma(Y)^2 &\leq \mathbf{E} N_\gamma(Y) + \mathbf{E} N_\gamma(Y)^2 \mathbf{1}_{2 \leq N_\gamma(Y) \leq n^{s/10}} + \mathbf{E} N_\gamma(Y)^2 \mathbf{1}_{n^{s/10} \leq N_\gamma(Y) \leq n} \\ &\ll n^{-s/2} + \mathbf{E} N_\gamma(Y)^2 \mathbf{1}_{2 \leq N_\gamma(Y) \leq n^{s/10}} + \mathbf{E} N_\gamma(Y)^2 \mathbf{1}_{n^{s/10} \leq N_\gamma(Y) \leq n} \quad \text{by [52, Corollary 3.7]} \\ &\ll n^{-s/2} + n^{-s} + n^{-90} \ll n^{-s/2} \quad \text{by (83) and (84)}. \end{aligned} \quad (85)$$

Similarly,

$$\mathbf{E} N_{I_p}^2(Y) \leq \mathbf{E} N_{I_p}^2(Y) \mathbf{1}_{N_{I_p}(Y) \leq n^{s/10}} + \mathbf{E} N_{I_p}^2(Y) \mathbf{1}_{n^{s/10} \leq N_{I_p}(Y) \leq n} \ll n^{s/5} + n^{-90} \ll n^{s/5}. \quad (86)$$

Plugging (85) and (86) into (82), we get

$$(\mathbf{E} N_{I_k}(Y) N_{I_p}(Y) - \mathbf{E} M_k(Y) N_{I_p}(Y))^2 \ll n^{-s/2} \mathbf{E} N_{I_p}^2(Y) \ll n^{-s/10}.$$

Similarly,

$$(\mathbf{E} M_k(Y) N_{I_p}(Y) - \mathbf{E} M_k(Y) M_p(Y))^2 \ll n^{-s/2}.$$

Combining these two inequalities gives (80) and completes the proof. \square

8. VARIANCE ESTIMATE UNDER THE NON-IID REGIME

Theorem 8.1. *Assume that the coefficients y_{ij} are independent (but not necessarily identically distributed) of mean zero, variance one, and bounded $(2 + \varepsilon_0)$ -moment: $\mathbf{E}|y_{ij}|^{2+\varepsilon_0} < C$ for some positive constants ε_0, C and for all i, j . Then there exists a constant $c > 0$ such that*

$$\mathbf{E}N_n = \left(\frac{2}{\sqrt{3}} + O(n^{-c}) \right) n \quad (87)$$

and

$$\text{Var}(N_n) = O(n^{2-c}), \quad (88)$$

where the implied constants depend on C and ε_0 .

Proof. Equation (87) is simply [52, Corollary 3.7] in which the c_i are all 1 and $u_n = 0$.

For Equation (88), as in Section 7, we recall the definition of the sub-intervals I_k in (17) and we denote by $N_{I_k}(Y)$ the number of roots of $P_n(\cdot, Y)$ in I_k . By (76), there exists a constant $c > 0$ such that for all indices k, p ,

$$\text{Cov}(N_{I_k}(Y), N_{I_p}(Y)) - \text{Cov}(N_{I_k}(G), N_{I_p}(G)) \ll n^{-2c}.$$

As mentioned in Remark 7.3 and the beginning of Subsection 7.5, this inequality was proven under the more general assumption that the y_{ij} are independent with mean zero, variance one, and bounded $(2 + \varepsilon_0)$ -moment.

Since $N_n(Y) = \sum_{k \in \mathbb{Z} \cap [-n\pi/\varepsilon, n\pi/\varepsilon]} N_{I_k}(Y)$ and similarly for $N_n(G)$, we have

$$\text{Var}(N_n(Y)) - \text{Var}(N_n(G)) = \sum_{k, p \in \mathbb{Z} \cap [-n\pi/\varepsilon, n\pi/\varepsilon]} \text{Cov}(N_{I_k}(Y), N_{I_p}(Y)) - \text{Cov}(N_{I_k}(G), N_{I_p}(G)) \ll n^{2-2c}.$$

By Theorem 1.2,

$$\text{Var}(N_n(G)) \ll n^{2-2c}.$$

Combining these bounds gives (88). □

9. CHARACTERISTIC FUNCTIONS IN \mathbb{R}^2 , PROOF OF THEOREM 2.4

Given a real number w and a variable z , we define the z -norm of w by

$$\|w\|_\xi := (\mathbf{E}\|w(\xi_1 - \xi_2)\|_{\mathbb{R}/\mathbb{Z}}^2)^{1/2},$$

where ξ_1, ξ_2 are two iid copies of ξ . For instance if z is Bernoulli with $\mathbf{P}(z = \pm 1) = 1/2$ (which is our main focus), then $\|w\|_\xi^2 = \|2w\|_{\mathbb{R}/\mathbb{Z}}^2/2$.

The following works for general \mathbb{R}^d : consider the random walk $\sum_i y_{i1} \mathbf{w}_i + y_{i2} \mathbf{w}'_i$, where $\mathbf{w}_i, \mathbf{w}'_i$ are vectors in \mathbb{R}^d . Then the corresponding characteristic function can be bounded by (see [60, Section 5])

$$\begin{aligned} |\prod \phi_i(x)| &= |\prod |\mathbf{E}e(y_{i1} \langle \mathbf{w}_i, x \rangle)| \prod |\mathbf{E}e(y_{i2} \langle \mathbf{w}'_i, x \rangle)|| \leq \prod_i [|\mathbf{E}e(y_{i1} \langle \mathbf{w}_i, x \rangle)|^2/2 + 1/2] \prod_i [|\mathbf{E}e(y_{i2} \langle \mathbf{w}'_i, x \rangle)|^2/2 + 1/2] \\ &\leq \exp\left(-\left(\sum_i \|\langle \mathbf{w}_i, x/2\pi \rangle\|_\xi^2 + \|\langle \mathbf{w}'_i, x/2\pi \rangle\|_\xi^2\right)/2\right). \quad (89) \end{aligned}$$

Hence if we have a good lower bound on the exponent $\sum_i \|\langle \mathbf{w}_i, x/2\pi \rangle\|_{\xi}^2 + \|\langle \mathbf{w}'_i, x/2\pi \rangle\|_{\xi}^2$ then we would have a good control on $|\prod \phi_i(x)|$. Furthermore, by definition

$$\begin{aligned} \sum_i \|\langle \mathbf{w}_i, x/2\pi \rangle\|_{\xi}^2 + \|\langle \mathbf{w}'_i, x/2\pi \rangle\|_{\xi}^2 &= \sum_i \mathbf{E} \|\langle \mathbf{w}_i, x/2\pi \rangle(\xi_1 - \xi_2)\|_{\mathbb{R}/\mathbb{Z}}^2 + \mathbf{E} \|\langle \mathbf{w}'_i, x/2\pi \rangle(\xi_1 - \xi_2)\|_{\mathbb{R}/\mathbb{Z}}^2 \\ &= \mathbf{E} \left(\sum_i \|\langle \mathbf{w}_i, x/2\pi \rangle(\xi_1 - \xi_2)\|_{\mathbb{R}/\mathbb{Z}}^2 + \sum_i \|\langle \mathbf{w}'_i, x/2\pi \rangle(\xi_1 - \xi_2)\|_{\mathbb{R}/\mathbb{Z}}^2 \right) \\ &= \mathbf{E}_y \left(\sum_i \|y \langle \mathbf{w}_i, x/2\pi \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 + \sum_i \|y \langle \mathbf{w}'_i, x/2\pi \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 \right), \end{aligned} \quad (90)$$

where $y = \xi_1 - \xi_2$. As ξ have mean zero, variance one and bounded $(2 + \varepsilon_0)$ -moment, there exist positive constants $c_1 \leq c_2, c_3$ such that $\mathbf{P}(c_1 \leq |y| \leq c_2) \geq c_3$, and so

$$\mathbf{E}_y \sum_i \|y \langle \mathbf{w}_i, x/2\pi \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 + \sum_i \|y \langle \mathbf{w}'_i, x/2\pi \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 \geq c_3 \inf_{c_1 \leq |y| \leq c_2} \sum_i \|y \langle \mathbf{w}_i, x/2\pi \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 + \sum_i \|y \langle \mathbf{w}'_i, x/2\pi \rangle\|_{\mathbb{R}/\mathbb{Z}}^2. \quad (91)$$

Hence for Theorem 2.4 (and similarly for Theorem 2.5) it suffices to show that for any $\mathbf{D} = (d_1, d_2)$ (which plays the role of $(y/2\pi)x$ such that $c_1 r \leq \|\mathbf{D}\|_2 \leq c_2 n^{C^*}$ we have

$$\sum_i \|\langle \mathbf{u}_i, \mathbf{D} \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 + \|\langle \mathbf{u}'_i, \mathbf{D} \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 \geq n^{2\tau}. \quad (92)$$

For $t \in [-n\pi, n\pi]$, we define $\psi_i(t), \psi'_i(t)$ by

$$\psi_i = d_1 \cos(it/n) - d_2 \frac{i}{n} \sin(it/n) \text{ and } \psi'_i = d_1 \sin(it/n) + d_2 \frac{i}{n} \cos(it/n). \quad (93)$$

In other words,

$$\psi_i = \langle \mathbf{D}, \mathbf{v}_i \rangle \text{ and } \psi'_i = \langle \mathbf{D}, \mathbf{v}'_i \rangle.$$

Let \mathbf{e} be the unit vector in the direction of D , $\mathbf{e} = \frac{\mathbf{D}}{\|\mathbf{D}\|_2}$. Define

$$T := n^{\tau*}.$$

Our key ingredient in the proof of Theorem 2.4 is the following refined analog of [41, Lemma 4.3].

Proposition 9.1. *Assume that t satisfies Condition 1. We have*

$$\sum_j \|\psi_j\|_{\mathbb{R}/\mathbb{Z}}^2 + \sum_j \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}}^2 \geq T.$$

It is clear this result implies Theorem 2.4 via (89). We note that [41] treated mainly with Bernoulli, and up to $\|\mathbf{D}\|_2 = n^{1/2+o(1)}$. This was generalized to $\|\mathbf{D}\|_2 = n^{1-o(1)}$ in [51] for general ensembles. Our starting point is similar to that paper, but the innovative of the current note is that we can also assume $n \ll \|\mathbf{D}\|_2$.

Before proving this main result of the section, we introduce a useful bound as follows.

Claim 9.2. *Assume that τ_* is sufficiently small given τ , and assume that t satisfies Condition 1. Let $I \subset [n]$ be any arithmetic progression of length $n^{1-6\tau_*}$. Then*

(1) *For all $\varepsilon_1, \varepsilon_2 \in \{-1, 1\}$, and any positive integer $A_0 = O(n^{\tau_*})$ there exists $i \in I$ so that*

$$\varepsilon_1 \sin(iA_0 s/n), \varepsilon_2 \cos(iA_0 s/n) > 0.$$

(2) *For any unit vector $\mathbf{e} \in \mathbb{R}^2$ we have*

$$\sum_{i \in I} \langle \mathbf{e}, \mathbf{u}_i \rangle^2 \geq n^{1-\tau} \text{ and } \sum_{i \in I} \langle \mathbf{e}, \mathbf{u}'_i \rangle^2 \geq n^{1-\tau}. \quad (94)$$

Proof. See the proof of Claim 10.1. □

For the rest of this section we prove Proposition 9.1 by contradiction: assume the opposite that we have

$$\sum_j \|\psi_j\|_{\mathbb{R}/\mathbb{Z}}^2 + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}}^2 \leq T. \quad (95)$$

We will then show that this is impossible as long as t satisfy Condition 1. We will do so by many steps. First, it follows from (95) that

$$|\{j \in [0, n) \cap \mathbb{Z} : \|\psi_j\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}} > 1/T\}| \leq 2T^3$$

and so for large n there exists an interval $J = [a, b] \subset [n]$ of length n/T^6 so that for $j \in J$

$$\|\psi_j\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}} < 1/T. \quad (96)$$

Differencing. Let A, k be chosen later so that

$$4\|\mathbf{D}\|_2 \frac{(4\pi)^k}{A^{(k-3)/2}} + 4 \times 2^k \frac{1}{T} < 1. \quad (97)$$

By pigeonholing we can find $p_0 \in \mathbb{Z}, p_0 \neq 0$ and t_0 so that

$$p_0 \frac{t}{2\pi} - t_0 \in \mathbb{Z}, 1 \leq |p_0| \leq A, |t_0| \leq \frac{4}{A}. \quad (98)$$

From the approximation we infer that

$$|e^{\sqrt{-1}p_0 \frac{t}{n}} - 1| = |e^{-\sqrt{-1}(2\pi t_0)} - 1| \leq |2 \sin(\pi t_0)| \leq 4\pi/A. \quad (99)$$

Next, for a sequence $\{g_j\}_{j \in [n]}$ we define the discrete differentials with step p_0 by

$$\Delta^k g_{j, p_0} := \sum_{i=0}^k \binom{k}{i} (-1)^i g_{j+ip_0}.$$

Let m_j and m'_j be the integers closest to ψ_j and ψ'_j respectively. Thus for $j \in J$ by (117) we have $|\psi_j - m_j| \leq 1/T$ and $|\psi'_j - m'_j| \leq 1/T$ and

$$|m_j|, |m'_j| \leq 2\|\mathbf{D}\|_2.$$

We show that

Lemma 9.3. *We have*

$$|\Delta^k m_{j, p_0}| + |\Delta^k m'_{j, p_0}| \leq 4\|\mathbf{D}\|_2 \frac{(4\pi)^k}{A^{(k-3)/2}} + 4 \times 2^k \frac{1}{T} \quad (100)$$

provided that $[j, j + kp_0] \subset J$.

Proof. (of Lemma 9.3) Recall that $\psi_j = d_1 \cos(jt/n) - d_2 \frac{j}{n} \sin(jt/n)$ and $\|\psi_j\|_{\mathbb{R}/\mathbb{Z}} \leq 1/T$ over $j \in J$. Consider

$$\Delta^k \psi_{j, p_0} = \sum_{i=0}^k \binom{k}{i} (-1)^i \psi_{j+ip_0}.$$

We first have

$$\begin{aligned} \sum_{i=0}^k \binom{k}{i} (-1)^i \cos\left(\frac{ip_0 t}{n} + \frac{jt}{n}\right) &= \operatorname{Re} \sum_{i=0}^k \binom{k}{i} (-1)^i e^{i\left(\frac{ip_0 t}{n} + \frac{jt}{n}\right)} \\ &= \operatorname{Re} \left[\left(\sum_{i=0}^k \binom{k}{i} (-1)^i e^{i\frac{ip_0 t}{n}} \right) e^{i\frac{jt}{n}} \right] \\ &= \operatorname{Re} \left[\left(1 - e^{i\frac{p_0 t}{n}} \right)^k e^{i\frac{jt}{n}} \right] \\ &\leq (4\pi/A)^k < (4\pi/\sqrt{A})^k \end{aligned} \quad (101)$$

where we used (99) in the last estimate. It also follows that

$$\begin{aligned}
\sum_{i=0}^k \binom{k}{i} (-1)^i \frac{j + ip_0}{n} \sin\left(\frac{ip_0 t}{n} + \frac{jt}{n}\right) &= \frac{\partial}{\partial t} \left(\sum_{i=0}^k \binom{k}{i} (-1)^i \cos\left(\frac{ip_0 t}{n} + \frac{jt}{n}\right) \right) \\
&= \frac{\partial}{\partial t} \left(\operatorname{Re} \left((1 - e(\frac{p_0 t}{n}))^k e(\frac{jt}{n}) \right) \right) \\
&= \operatorname{Re} \left(\frac{\partial}{\partial t} \left((1 - e(\frac{p_0 t}{n}))^k e(\frac{jt}{n}) \right) \right) \\
&= \operatorname{Re} \left(-k \sqrt{-1} \frac{p_0}{n} (1 - e(\frac{p_0 t}{n}))^{k-1} e(\frac{jt}{n}) \right) \\
&\quad + \sqrt{-1} \frac{j}{n} (1 - e(\frac{p_0 t}{n}))^k e(\frac{jt}{n}) \\
&\leq A(4\pi/A)^{k-1} + (4\pi/A)^k < (4\pi)^k / A^{(k-3)/2}. \tag{102}
\end{aligned}$$

Putting the bounds together we obtain

$$|\Delta^k \psi_{j,p_0}| = \left| \sum_{i=0}^k \binom{k}{i} (-1)^i \psi_{j+ip_0} \right| \leq |d_1| \left(\frac{4\pi}{\sqrt{A}} \right)^k + |d_2| \frac{(4\pi)^k}{A^{(k-3)/2}} < 4 \|\mathbf{D}\|_2 \frac{(4\pi)^k}{A^{(k-3)/2}}.$$

One can also obtain similarly estimates for $|\Delta^k \psi'_{j,p_0}|$ where we recall that $\psi'_i = d_1 \sin(it/n) + d_2 \frac{i}{n} \cos(it/n)$. More precisely, as in (101)

$$\begin{aligned}
\sum_{i=0}^k \binom{k}{i} (-1)^i \sin\left(\frac{ip_0 t}{n} + \frac{jt}{n}\right) &= \operatorname{Im} \sum_{i=0}^k \binom{k}{i} (-1)^i e\left(\frac{ip_0 t}{n} + \frac{jt}{n}\right) \\
&= \operatorname{Im} \left[(1 - e(\frac{p_0 t}{n}))^k e(\frac{jt}{n}) \right] \\
&\leq (4\pi/\sqrt{A})^k. \tag{103}
\end{aligned}$$

Also, as in (102)

$$\begin{aligned}
\sum_{i=0}^k \binom{k}{i} (-1)^i \frac{j + ip_0}{n} \cos\left(\frac{ip_0 t}{n} + \frac{jt}{n}\right) &= \frac{\partial}{\partial t} \left(\sum_{i=0}^k \binom{k}{i} (-1)^i \sin\left(\frac{ip_0 t}{n} + \frac{jt}{n}\right) \right) \\
&= \operatorname{Im} \left(\frac{\partial}{\partial t} \left((1 - e(\frac{p_0 t}{n}))^k e(\frac{jt}{n}) \right) \right) \\
&< (4\pi)^k / A^{(k-3)/2}. \tag{104}
\end{aligned}$$

It thus follows that

$$\begin{aligned}
|\Delta^k m_{j,p_0}| &\leq |\Delta^k \psi_{j+l_{p_0}}| + |\Delta^k (\psi_{j+l_{p_0}} - m_{j+l_{p_0}})| \\
&\leq 4 \|\mathbf{D}\|_2 \frac{(4\pi)^k}{A^{(k-3)/2}} + 4 \times 2^k \frac{1}{T},
\end{aligned}$$

and similarly for $|\Delta^k m'_{j,p_0}|$. □

Now, with a sufficiently small τ_* (given τ) we choose

$$A = n^{\tau_*} \text{ and } k = 16 \lfloor \frac{\log \|\mathbf{D}\|_2}{\varepsilon \log n} \rfloor. \tag{105}$$

We note that $k = O(1)$ because $\|\mathbf{D}\|_2 \leq n^{C^*}$. (We restrict $\|\mathbf{D}\|_2$ to grow polynomially here so that k is bounded, which significantly simplifies our later analysis). We also recall that $p_0 \leq A \leq n^{\tau_*}$.

With these choices we see that (97) is fulfilled, and hence the RHS terms in Lemma 9.3 are strictly smaller than 1. But because the numbers m_j, m'_j are integers, it follows that as long as $\{j, j + p_0, \dots, j + kp_0\} \subset$

$J = [a, b]$ we must have

$$\Delta^k m_{j,p_0} = 0 \text{ and } \Delta^k m'_{j,p_0} = 0. \quad (106)$$

Now we consider the sequence of integers $\{m_{j+ip_0}, 0 \leq i \leq (b-j)/p_0\}$. By repeatedly applying (122) for $j = j, j+p_0, j+2p_0, \dots$ we see the k -differential of any $k+1$ consecutive terms of this sequence is zero. We thus deduce from here a crucial conclusion below.

Lemma 9.4. *For given $j \in J$, there exists a polynomial of degree at most k so that for any $0 \leq i \leq (b-j)/p_0$ we have*

$$m_{j+ip_0} = P_{j,p_0}(i).$$

We also have a similarly conclusion for m'_{j+ip_0} where the polynomial can be different.

We note that this result holds for any j such that $[j, j+kp_0] \subset J$. Now we will exploit these polynomial properties furthermore by specifying the choices of parameters. We first consider the case that $\|\mathbf{D}\|_2$ is small.

Case 1. Assume that $\|\mathbf{D}\|_2 \leq n^{1-4\tau}$. It suffices to work with the ψ_j sequence, the treatment for ψ'_j is identical. Here we will show that P_j is a constant. Indeed, otherwise then as P'_j has at most $k-2$ roots, there is an interval of length $|J|/k$ where P_j is strictly monotone. But on this interval (of length $|J|/k \geq n/kT^6 \geq n^{1-6\tau_* - o(1)}$ at least), $m_j \in [-2n^{1-4\tau}, 2n^{1-4\tau}]$ because $|m_j| \leq 2\|\mathbf{D}\|_2$, so this is impossible if τ_* is sufficiently small compared to τ . Thus we have shown that the polynomials are constant,

$$m_{j+lp_0} = m_j, \forall j, l \in \mathbb{Z}, [j+lp_0, j+(l+k)p_0] \subset J = [a, b].$$

Let's next fix j , then the range for l is $I = (a-j)/p_0 \leq l \leq (b-j)/p_0 - O(1)$, which is an interval of length of order $n^{1-7\tau_*}$. On this range of l , the condition of t in 1 shows that $\psi_{j+lp_0} = d_1 \cos((j+lp_0)t) - d_2 \frac{i}{n} \sin((j+lp_0)t)$ changes size. But as $m_{j+lp_0} = m$ is the common integral part for all l , this is impossible unless $m_{j+lp_0} = 0$ for all l in the range I above.

Argue similarly for ψ'_j , we have thus obtained in this case that all the integral parts are zero, and hence

$$\sup_{l \in I_L} |\psi_{j+lp_0}| = \sup_{l \in I} |\psi_{j+lp_0}|_{\mathbb{R}/\mathbb{Z}}, \sup_{l \in I} |\psi'_{j+lp_0}| = \sup_{l \in I} |\psi'_{j+lp_0}|_{\mathbb{R}/\mathbb{Z}} \leq 1/T = n^{-\tau_*}.$$

It follows that

$$\begin{aligned} \sum_i \|\langle \mathbf{u}_i, \mathbf{D} \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 + \|\langle \mathbf{u}'_i, \mathbf{D} \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 &\geq \sum_{l \in I} \|\langle \mathbf{u}_{j+lp_0}, \mathbf{D} \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 + \sum_{l \in I} \|\langle \mathbf{u}'_{j+lp_0}, \mathbf{D} \rangle\|_2^2 \\ &= \sum_{l \in I} \|\langle \mathbf{u}_{j+lp_0}, \mathbf{D} \rangle\|_2^2 + \sum_{l \in I} \|\langle \mathbf{u}'_{j+lp_0}, \mathbf{D} \rangle\|_2^2 \\ &= \|\mathbf{D}\|_2^2 \sum_{l \in I} \langle \mathbf{u}_{j+lp_0}, \mathbf{e} \rangle^2 + \langle \mathbf{u}'_{j+lp_0}, \mathbf{e} \rangle^2 \geq r^2 n^{1-8\tau} > n^{2\tau}, \end{aligned}$$

where we used Claim 9.2. This bound clearly contradicts (95).

Case 2. Assume that $n^{1-4\tau} \leq \|\mathbf{D}\|_2 \leq n^{C_*}$. Roughly speaking, our approach here is of inverse-type in the sense that we will try to gain as much as possible information for t given the obtained bounds; and our final result on t is almost optimal.

Recall that $\sum_{j \in J} \|\psi_j\|_{\mathbb{R}/\mathbb{Z}}^2 + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}}^2 \leq T$ and by Cauchy-Schwarz we have

$$\sum_{j \in J} \|\psi_j\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}} \leq 2\sqrt{nT}.$$

We will reapply the process from (101) and (102) with $q_0 = lp_0$ (for a given positive integer l). By the polynomial properties of the $m_{j+ip_0} = P(i)$ for a polynomial P of degree at most $k-1$, we also have that

$$\sum_{i=0}^k \binom{k}{i} (-1)^i m_{j+iq_0} = \sum_{i=0}^k \binom{k}{i} (-1)^i m_{j+ilp_0} = \sum_{i=0}^k \binom{k}{i} (-1)^i P_{j,p_0}(il) = 0.$$

Set

$$z_{q_0} := z(t, q_0) = e(q_0 t/n). \quad (107)$$

We then write as follows

$$\begin{aligned} |\Delta^k \psi_{j, q_0}| &= \left| \sum_{i=0}^k \binom{k}{i} (-1)^i \psi_{j+iq_0} \right| = \left| \sum_{i=0}^k \binom{k}{i} (-1)^i (\psi_{j+iq_0} - m_{j+iq_0}) + \sum_{i=0}^k \binom{k}{i} (-1)^i m_{j+iq_0} \right| \\ &= \left| \sum_{i=0}^k \binom{k}{i} (-1)^i (\psi_{j+iq_0} - m_{j+iq_0}) \right| \leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned}$$

Similarly,

$$\begin{aligned} |\Delta^k \psi'_{j, q_0}| &= \left| \sum_{i=0}^k \binom{k}{i} (-1)^i \psi'_{j+iq_0} \right| = \left| \sum_{i=0}^k \binom{k}{i} (-1)^i (\psi'_{j+iq_0} - m'_{j+iq_0}) + \sum_{i=0}^k \binom{k}{i} (-1)^i m'_{j+iq_0} \right| \\ &= \left| \sum_{i=0}^k \binom{k}{i} (-1)^i (\psi'_{j+iq_0} - m'_{j+iq_0}) \right| \leq 2^k \sum_{i=0}^k \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \Delta^k \psi_{j, q_0} &= \sum_{i=0}^k \binom{k}{i} (-1)^i \psi_{j+iq_0} = d_1 \sum_{i=0}^k \binom{k}{i} (-1)^i \cos\left(\frac{iq_0 t}{n} + \frac{jt}{n}\right) + d_2 \sum_{i=0}^k \binom{k}{i} (-1)^i \frac{j+iq_0}{n} \sin\left(\frac{iq_0 t}{n} + \frac{jt}{n}\right) \\ &= d_1 \operatorname{Re}\left((1-z_{q_0})^k e(jt/n)\right) + d_2 \operatorname{Re}\left(-k\sqrt{-1} \frac{q_0}{n} ((1-z_{q_0})^{k-1} e(jt/n)) + \sqrt{-1} j/n ((1-z_{q_0})^k e(jt/n))\right) \end{aligned}$$

and

$$\begin{aligned} \Delta^k \psi'_{j, q_0} &= \sum_{i=0}^k \binom{k}{i} (-1)^i \psi'_{j+iq_0} = d_1 \sum_{i=0}^k \binom{k}{i} (-1)^i \sin\left(\frac{iq_0 t}{n} + \frac{jt}{n}\right) + d_2 \sum_{i=0}^k \binom{k}{i} (-1)^i \frac{j+iq_0}{n} \cos\left(\frac{iq_0 t}{n} + \frac{jt}{n}\right) \\ &= d_1 \operatorname{Im}\left((1-z_{q_0})^k e(jt/n)\right) + d_2 \operatorname{Im}\left(-k\sqrt{-1} \frac{q_0}{n} ((1-z_{q_0})^{k-1} e(jt/n)) + \sqrt{-1} j/n ((1-z_{q_0})^k e(jt/n))\right). \end{aligned}$$

By the triangle inequality, the above then implies that as long as $[j, j+kq_0] \subset J$ we have

$$\begin{aligned} &\left| (1-z_{q_0})^{k-1} \left(d_1 (1-z_{q_0}) + \sqrt{-1} d_2 \frac{-kq_0}{n} + \sqrt{-1} d_2 (1-z_{q_0}) \frac{j}{n} \right) \right| \\ &\leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned} \quad (108)$$

We recall that this holds for any $j \in J$ as long as $[j, j+kq_0] \subset J$, and there is no $P_{j, p_0}(\cdot)$ or m_{j+iq_0} on the LHS. Applying the estimate for $j = j+L$, and using triangle inequality we obtain the following.

Lemma 9.5. *Assume that $[j, j+L+kq_0] \subset J$. We then have*

$$\begin{aligned} &\left| (1-z_{q_0})^k \left| \frac{d_2 L}{n} \right. \right| \\ &\leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2^k \sum_{i=0}^k \|\psi_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned} \quad (109)$$

Now we complete the proof of Proposition 9.1. We will work with $q_0 = lp_0$ so that $kq_0 \leq |J|/2$. Recall that $J = [a, b]$ with length $|J|$ at least $n/T^6 = n^{1-6\tau^*}$, and hence $(a+b)/2 \geq |J|/2 \geq n^{1-6\tau^*}/2$. We divide the treatment into two cases depending on the parameters d_1, d_2 .

Subcase 1. Assume that $|d_2| \geq n^{-\tau^*}$. We first fix $q_0 = lp_0$ such that

$$|q_0| \leq |J|/8k. \quad (110)$$

Choose any $L \in [|J|/8, |J|/4]$, say $L = |J|/8$. With this fixed L , it is clear that $|d_2 L/n| \geq n^{-\tau^*} n^{1-7\tau^*}/n \geq n^{-8\tau^*}$. We will then choose j in the interval $j \in [(a+b)/2 - |J|/8, (a+b)/2 + |J|/8]$ such that the RHS

of (109) is as small as possible. (Clearly with these choices we have $[j, j + L + kq_0] \subset J$.) If we sum of the RHS of (109) for j from the above range, then this sum is bounded by $O_k(\sum_{i \in J} \|\psi_i\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_i\|_{\mathbb{R}/\mathbb{Z}}) = O(\sqrt{nT}) = O(n^{1/2+\tau_*})$ because each term $\|\psi_i\|_{\mathbb{R}/\mathbb{Z}}$ appears a bounded number of times in the total sum. Hence by averaging, there exists $j \in [(a+b)/2 - |J|/8, (a+b)/2 + |J|/8]$ such that the RHS of (109) is bounded by $O(n^{1/2+\tau_*}/(|J|/4)) = O(n^{-1/2+7\tau_*})$.

Putting together, with such choice of L and j , the equation (109) implies that

$$|(1 - z_{q_0})^k| \times n^{-8\tau_*} \leq O(n^{-1/2+7\tau_*}).$$

Thus we have that $|(1 - z_{q_0})^k| \leq n^{-1/2+15\tau_*}$ in this case, and so with sufficiently small τ_*

$$|1 - z_{q_0}| \leq n^{-1/3k}. \quad (111)$$

As of this point, recall that $z_{q_0} = e(q_0 t/n)$. Because (111) holds for any $q_0 = lp_0$ satisfying (110), we thus have for all $1 \leq l \leq n^{1-8\tau_*} \leq |J|/(8kp_0)$

$$|1 - e(lp_0 t/n)| \leq n^{-1/3k}. \quad (112)$$

As of this point we then use the following elementary result to obtain more information on t .

Claim 9.6. *Assume that $z = e(\theta)$, $|\theta| \leq \pi/8$ such that for all $1 \leq l \leq M$ we have $|1 - z^l| \leq 1/32$ for a sufficiently large M . Then $|\theta| = O(1/M)$.*

Proof. By assumption, $|\theta| \leq \pi/8$ and $\|2^m \theta\|_{\mathbb{R}/\mathbb{Z}} \leq \pi/8$ for all $1 \leq m \leq \log M$, and so we can repeatedly estimate $|\theta|$ to obtain $|\theta| = O(1/M)$. \square

Claim 9.6 and (112) then implies that for large enough n ,

$$\|p_0 t/\pi n\|_{\mathbb{R}/\mathbb{Z}} = O(n^{-1+8\tau_*}).$$

However this contradicts Condition 1 because $p_0 \leq A = n^{\tau_*}$ and τ_* is sufficiently small given τ .

Subcase 2. Now we consider the remaining (very degenerate) case that $|d_2| \leq n^{-\tau_*}$. Then $|d_1| \asymp \|\mathbf{D}\|_2 \geq n^{1-4\tau}$. In the case that $d_1|z_{q_0} - 1| \leq n^{4\tau_*}$ then we have

$$|z_{q_0} - 1| \leq n^{-1+4\tau-4\tau_*}.$$

In the other case that $d_1|z_{q_0} - 1| \geq n^{4\tau_*}$, then this term dominates all other terms involving d_2 on the LHS of (108) (because each of which has order $O(1)$ as d_2 is small). So we have

$$|(1 - z_{q_0})^k| d_1/2 \leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} = O(1).$$

Hence,

$$|z_{q_0} - 1| = O(1/d_1^{1/k}) \leq n^{-1/2k}.$$

Thus from both scenarios on the magnitude of $d_1|z_{q_0} - 1|$ we at least have $|z_{q_0} - 1| \leq n^{-1/2k}$. We can then repeat the argument as in the previous case to vary q_0 and use Claim 9.6. It thus also follows that $\|p_0 t/\pi n\|_{\mathbb{R}/\mathbb{Z}} = O(n^{-1+8\tau_*})$, which is again impossible.

Before concluding this section, as our approach to prove Proposition 9.1 starts with (96), by considering subintervals of J when needed (where we note that at least one of such subintervals still has length $\Omega(n/T^6)$), we obtain the following analog of Theorem 2.4.

Theorem 9.7. *Let C_* and ℓ be given positive constants, and t satisfies Condition 1 for some sufficiently small constant τ . The following holds for sufficiently large n and sufficiently small τ_* (depending on C_* , τ and ℓ). For any $n^{5\tau-1/2} \leq \|x\|_2 \leq n^{C_*}$, for any set $I \subset [n]$ of at most ℓ entries we have*

$$|\prod_{i \notin I} \phi_i(x)| \leq \exp(-n^{\tau_*}).$$

10. CHARACTERISTIC FUNCTIONS IN \mathbb{R}^4 , PROOF OF THEOREM 2.5

In this section we continue our “inverse-type” analysis of the characteristic functions, but now in \mathbb{R}^4 . As there are two parameters s, t from Condition 2, the situation is much more complicated, but we will call on the previous sections whenever possible.

First we show the following analog of Claim 9.2.

Claim 10.1. *Assume that τ_* is sufficiently small given τ , and assume that s, t satisfy Condition 2. Let $I = \{a + lq, 0 \leq l \leq L\} \subset [n]$ be any arithmetic progression of length $L = n^{1-6\tau_*}$. Then*

(1) *For all $\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4 \in \{-1, 1\}$ and $\theta \in [-\pi, \pi]$, and for any positive integer $A_0 = O(n^{\tau_*})$ there exists $i \in I$ so that*

$$\varepsilon_1 \sin(iA_0s/n + \theta), \varepsilon_2 \cos(iA_0s/n + \theta), \varepsilon_3 \sin(iA_0t/n + \theta), \varepsilon_4 \cos(iA_0t/n + \theta) > 0.$$

(2) *For any unit vector $\mathbf{e} \in \mathbb{R}^4$ we have*

$$\sum_{i \in I} \langle \mathbf{e}, \mathbf{v}_i \rangle^2 \geq n^{1-\tau} \quad \text{and} \quad \sum_{i \in [n]} \langle \mathbf{e}, \mathbf{v}'_i \rangle^2 \geq n^{1-\tau}. \quad (113)$$

Proof. Note that $q \leq n/L \leq n^{6\tau_*}$, which is much smaller than n^τ if τ_* is chosen sufficiently small. We first show (113) for \mathbf{v}_i , given any unit vector $\mathbf{e} = (x_1, x_2, x_3, x_4)$. First, replacing s, t by qs, qt and rotate \mathbf{e} if need, without loss of generality we assume that $I = [0, L]$. Then

$$\begin{aligned} \sum_{i \in I} \langle \mathbf{e}, \mathbf{v}_i \rangle^2 &= \sum_i (x_1 \cos(it/n) - x_2(i/n) \sin(it/n) + x_3 \cos(is/n) - x_4(i/n) \sin(is/n))^2 \\ &\quad + (x_1 \sin(it/n) + x_2(i/n) \cos(it/n) + x_3 \sin(is/n) + x_4(i/n) \cos(is/n))^2. \end{aligned}$$

Clearly the sum over the diagonal terms is $n \sum_i x_i^2 = L$. We thus need to work with the cross terms

$$A_1 = \sum_{i \in I} \cos(is/n) \cos(it/n), \quad B_1 = \sum_{i \in I} (i/n)^2 \cos(is/n) \cos(it/n)$$

and

$$C_1 = \sum_{i \in I} \sin(is/n) \sin(it/n), \quad D_1 = \sum_{i \in I} (i/n)^2 \sin(is/n) \sin(it/n).$$

We show that under the condition on s, t (and (qs, qt)) from the claim these terms are all of order $o(L)$ (actually we obtain a slightly strong bound). Let's work with B_1 for instance, we have

$$\begin{aligned} 2B_1 &= \sum_{i \in I} (i/n)^2 (\cos(is/n - it/n) + \cos(is/n + it/n)) = \left(-\sum_{i \in I} \cos(ix/n)\right)'' \Big|_{x=s-t} + \left(-\sum_{i \in I} \cos(ix/n)\right)'' \Big|_{x=s+t} \\ &= \left(\frac{\sin((L+1/2)x/n)}{\sin(x/2n)} - 1\right)'' \Big|_{x=s-t} + \left(\frac{\sin((L+1/2)x/n)}{\sin(x/2n)} - 1\right)'' \Big|_{x=s+t} = O\left(\frac{1}{|\sin^3((s-t)/2n)|} + \frac{1}{|\sin^3((s+t)/2n)|}\right. \\ &\quad \left.+ \frac{1}{n} \frac{1}{|\sin^2((s-t)/2n)|} + \frac{1}{n} \frac{1}{|\sin^2((s+t)/2n)|} + \frac{1}{n^2} \frac{1}{|\sin((s-t)/2n)|} + \frac{1}{n^2} \frac{1}{|\sin^2((s+t)/2n)|}\right). \end{aligned}$$

It thus follows that, because $|(s-t)/\pi n|_{\mathbb{R}/\mathbb{Z}} \geq n^{-1+8\tau}$ and $|(s+t)/\pi n|_{\mathbb{R}/\mathbb{Z}} \geq n^{-1+8\tau}$,

$$|B_1| = O(n^{1-8\tau}).$$

The treatments for D_1, A_1, C_1 are somewhat simpler, and hence we omit.

Now we focus on the first part. By the (quantitative) Weyl's equi-distribution criterion on \mathbf{T}^2 (see for instance [62, Proposition 9; Exercises 18, 19]) if the sequence $\{(i(qA_0s/\pi n) + \theta, i(qA_0t/\pi n) + \theta), 1 \leq i \leq L\}$ in the two dimensional torus $(\mathbb{R}/\mathbb{Z})^2$, where $A_0 = O(n^{\tau_*})$ and $q \leq n^{6\tau_*}$, is not δ -equidistributed⁴ (for some fixed

⁴Actually we just need the points to appear in all four quadrants of the plane.

sufficiently small constant δ to guarantee the sign changes) then there exist positive integers $k_0, l_0 = n^{O_{\tau_*}(1)}$ such that

$$\|k_0(A_0qs/\pi n) + l_0(A_0qt/\pi n)\|_{\mathbb{R}/\mathbb{Z}} = O\left(\frac{n^{O_{\tau_*}(1)}}{L}\right) = O\left(\frac{1}{n^{1-8\tau}}\right),$$

provided that τ_* is sufficiently small compared to τ . This contradicts with our condition. \square

Let $\mathbf{D} = (d_1, d_2, d_3, d_4) \in \mathbb{R}^4$ be any non-zero vector and let \mathbf{e} be the unit vector in the direction of \mathbf{D} . For $s, t \in [-n\pi, n\pi]$ we define

$$\psi_i := \langle \mathbf{D}, \mathbf{v}_i \rangle = d_1 \cos(it/n) - d_2 \frac{i}{n} \sin(it/n) + d_3 \cos(is/n) - d_4 \frac{i}{n} \sin(is/n) \quad (114)$$

$$\psi'_i := \langle \mathbf{D}, \mathbf{v}'_i \rangle = d_1 \sin(it/n) + d_2 \frac{i}{n} \cos(it/n) + d_3 \sin(is/n) + d_4 \frac{i}{n} \cos(is/n). \quad (115)$$

Define $T = n^{\tau_*}$. Our key ingredient in the proof of Theorem 2.5 is the following analog of Proposition 9.1.

Proposition 10.2. *Assume that s, t satisfy Condition 2. We have*

$$\sum_j \|\psi_j\|_{\mathbb{R}/\mathbb{Z}}^2 + \sum_j \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}}^2 \geq T.$$

It is clear this result implies Theorem 2.5 via (89), (90) and (91). For the rest of this section we prove Proposition 10.2 by contradiction: assume the opposite that we have

$$\sum_j \|\psi_j\|_{\mathbb{R}/\mathbb{Z}}^2 + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}}^2 \leq T. \quad (116)$$

We will then show that this is impossible as long as s, t satisfy Condition 2. First, argue as in (96), it follows from (116) that there exists an interval $J = [a, b] \subset [n]$ of length n/T^6 so that for $j \in J$

$$\|\psi_j\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}} < 1/T. \quad (117)$$

Differencing. Let A, k be chosen as in (97). We then can find $p_0 \in \mathbb{Z}, p_0 \neq 0$ and s_0, t_0 so that

$$p_0 \frac{s}{2\pi n} - s_0 \in \mathbb{Z}, p_0 \frac{t}{2\pi n} - t_0 \in \mathbb{Z}, 1 \leq |p_0| \leq A, |s_0|^2 + |t_0|^2 \leq \frac{4}{A}. \quad (118)$$

Indeed, consider the sequence of pairs $(\{q(s/2\pi n)\}, \{q(t/2\pi n)\}), 1 \leq q \leq A$ in $[0, 1]^2$. Using Dirichlet's principle, there exists q_1, q_2 such that the distance of the pairs is at most $2/\sqrt{A}$. In other words,

$$|\{q_1(s/2\pi n)\} - \{q_2(s/2\pi n)\}|^2 + |\{q_1(t/2\pi n)\} - \{q_2(t/2\pi n)\}|^2 \leq \frac{4}{A}.$$

This implies that there exists integers p_1, p_2 such that

$$|(q_1 - q_2)(s/2\pi n) - p_1|^2 + |(q_1 - q_2)(t/2\pi n) - p_1|^2 \leq \frac{4}{A}.$$

Set $s_0 = (q_1 - q_2)(s/2\pi n) - p_1$ and $t_0 = (q_1 - q_2)(t/2\pi n) - p_1$ and $p_0 = q_1 - q_2$ we obtain as claimed.

From the approximation we infer that

$$|e(p_0 \frac{t}{n}) - 1| = |e(2\pi t_0) - 1| \leq |2 \sin(\pi t_0)| \leq 4\pi/\sqrt{A} \quad (119)$$

and

$$|e(p_0 \frac{s}{n}) - 1| = |e(2\pi s_0) - 1| \leq |2 \sin(\pi s_0)| \leq 4\pi/\sqrt{A}. \quad (120)$$

Next, consider the differential operation $\Delta^k g_{j, p_p}$ as in the previous section. Let m_j and m'_j be the integers closest to ψ_j and ψ'_j respectively. We then have

Lemma 10.3. *We have*

$$|\Delta^k m_{j,p_0}| + |\Delta^k m'_{j,p_0}| \leq 4\|\mathbf{D}\|_2 \frac{(4\pi)^k}{A^{(k-3)/2}} + 4 \times 2^k \frac{1}{T} \quad (121)$$

provided that $[j, j + kp_0] \subset J$.

Proof. (of Lemma 10.3) Argue similarly as in the proof of Lemma 9.3, by using (119) and (120) we can show that

$$|\Delta^k \psi_j| = \left| \sum_{i=0}^k \binom{k}{i} (-1)^i \psi_{j+ip_0} \right| \leq (|d_1| + |d_3|) \left(\frac{4\pi}{\sqrt{A}} \right)^k + (|d_2| + |d_4|) \frac{(4\pi)^k}{A^{(k-3)/2}} < 4\|\mathbf{D}\|_2 \frac{(4\pi)^k}{A^{(k-3)/2}}.$$

One can also obtain similarly estimates for $|\Delta^k \psi'_j|$. It thus follows that

$$\begin{aligned} |\Delta^k m_j| &\leq |\Delta^k \psi_{j+lp_0}| + |\Delta^k (\psi_{j+lp_0} - m_{j+lp_0})| \\ &\leq 4\|\mathbf{D}\|_2 \frac{(4\pi)^k}{A^{(k-3)/2}} + 4 \times 2^k \frac{1}{T}, \end{aligned}$$

and similarly for $|\Delta^k m'_j|$. □

As of this point, we choose A and k as in (105). Then as long as $\{j, j + p_0, \dots, j + kp_0\} \subset J = [a, b]$ we must have

$$\Delta^k m_{j,p_0} = 0 \text{ and } \Delta^k m'_{j,p_0} = 0. \quad (122)$$

This results into the following analog of Lemma 9.4.

Lemma 10.4. *For given $j \in J$, there exists a polynomial of degree at most k so that for any $0 \leq i \leq (b-j)/p_0$ we have*

$$m_{j+ip_0} = P_{j,p_0}(i).$$

We also have a similarly conclusion for m'_{j+ip_0} .

We again note that this result holds for any j such that $[j, j + kp_0] \subset J$. We next consider the case that $\|\mathbf{D}\|_2$ is small.

Case 1. Assume that $\|\mathbf{D}\|_2 \leq n^{1-4\tau}$. Here our treatment is identical to Case 1. of the previous proof, that we can deduce from here that $m_{j+lp_0} = m_j, \forall j, l \in \mathbb{Z}, [j + lp_0, j + (l+k)p_0] \subset J = [a, b]$ and as over the interval $I = (a-j)/p_0 \leq l \leq (b-j)/p_0 - O(1)$, the condition of s, t in 2 shows that $\psi_{j+lp_0} = d_1 \cos((j + lp_0)t) - d_2 \frac{i}{n} \sin((j + lp_0)t) + d_3 \cos((j + lp_0)s/n) - d_4 \frac{(j+lp_0)}{n} \sin((j + lp_0)s/n)$ changes size and this implies that $m_{j+lp_0} = 0$ for all l in the range I above.

One will then have

$$\begin{aligned} \sum_i \|\langle \mathbf{v}_i, \mathbf{D} \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 + \|\langle \mathbf{v}'_i, \mathbf{D} \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 &\geq \sum_{l \in I} \|\langle \mathbf{u}_{j+lp_0}, \mathbf{D} \rangle\|_{\mathbb{R}/\mathbb{Z}}^2 + \sum_{l \in I} \|\langle \mathbf{v}'_{j+lp_0}, \mathbf{D} \rangle\|_2^2 \\ &= \sum_{l \in I} \|\langle \mathbf{v}_{j+lp_0}, \mathbf{D} \rangle\|_2^2 + \sum_{l \in I} \|\langle \mathbf{v}'_{j+lp_0}, \mathbf{D} \rangle\|_2^2 \\ &= \|\mathbf{D}\|_2^2 \sum_{l \in I} \langle \mathbf{v}_{j+lp_0}, \mathbf{e} \rangle^2 + \langle \mathbf{v}'_{j+lp_0}, \mathbf{e} \rangle^2 \geq r^2 n^{1-8\tau} > n^{2\tau}, \end{aligned}$$

where we used the second point of Claim 10.1. This bound contradicts (116).

Case 2. Assume that $n^{1-4\tau} \leq \|\mathbf{D}\|_2 \leq n^{C_*}$. In this case our “inverse” analysis is more complicated than that of the previous section because there are two unknowns. Nevertheless, our final bounds are almost optimal.

Recall that $\sum_{j \in J} \|\psi_j\|_{\mathbb{R}/\mathbb{Z}}^2 + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}}^2 \leq T$ and by Cauchy-Schwarz we have

$$\sum_{j \in J} \|\psi_j\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}} \leq 2\sqrt{nT}.$$

By reapplying the differential process as in Case 2 of Section 9 (this time for s and t) with $q_0 = lp_0$ for a given positive integer l . Set

$$z_{q_0,t} = e(q_0t/n) \text{ and } z_{q_0,s} = e(q_0s/n). \quad (123)$$

By the polynomial properties of the $m_{j+ip_0} = P(i)$ for a polynomial P of degree at most $k-1$, we obtain the following analog of (108).

$$\begin{aligned} & \left| (1 - z_{q_0,t})^{k-1} e(jt/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j}{n} \right) \right. \\ & \left. + (1 - z_{q_0,s})^{k-1} e(js/n) \left(d_3(1 - z_s) + \sqrt{-1}d_4 \frac{-kq_0}{n} + \sqrt{-1}d_4(1 - z_s) \frac{j}{n} \right) \right| \\ & \leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned} \quad (124)$$

In what follows, for a fixed q_0 , we will take advantage of the above inequality by varying j and eliminate the terms involving $(1 - z_s)$. Let L be a parameter to be chosen, we show the following

Lemma 10.5. *Assume that $[j_1, j_1 + j + L + kq_0] \subset J = [a, b]$. We then have*

$$\begin{aligned} & |(1 - z_{q_0,t})^k| \times (d_2 \frac{j}{n}) |e(L(s-t)/n) - 1|^2 \\ & \leq 2^k \sum_{i=0}^k \|\psi_{j_1+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi_{j_1+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi_{j_1+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\ & + 2^k \sum_{i=0}^k \|\psi'_{j_1+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi'_{j_1+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j_1+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\ & + 2^k \sum_{i=0}^k \|\psi_{j_1+j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi_{j_1+j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi_{j_1+j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\ & + 2^k \sum_{i=0}^k \|\psi'_{j_1+j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi'_{j_1+j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j_1+j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned}$$

We also have the same bound for $|(1 - z_{q_0,s})^k| \times (d_4 \frac{j}{n}) |e(L(s-t)/n) - 1|^2$.

Proof. In what follows we note that if $[j_1, j_1 + j + L + kq_0] \subset J$ then automatically $[j_1, j_1 + kq_0] \cup [j_1 + L, j_1 + L + kq_0] \cup [j_1 + j, j_1 + j + kq_0] \cup [j_1 + j + L, j_1 + j + L + kq_0] \subset J$, and so we can apply the followings as the indices are all in J . First, multiply both side of (124) by $e(Ls/n)$

$$\begin{aligned} & \left| (1 - z_{q_0,t})^{k-1} e(Ls/n) e(jt/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j}{n} \right) \right. \\ & \left. + (1 - z_{q_0,s})^{k-1} e((j+L)s/n) \left(d_3(1 - z_{q_0,s}) + \sqrt{-1}d_4 \frac{-kq_0}{n} + \sqrt{-1}d_4(1 - z_{q_0,s}) \frac{j}{n} \right) \right| \\ & \leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned}$$

One the other hand, (124) applied for j being replaced by $j + L$ shows

$$\begin{aligned} & \left| (1 - z_{q_0,t})^{k-1} e((j+L)t/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j+L}{n} \right) \right. \\ & \left. + (1 - z_{q_0,s})^{k-1} e((j+L)s/n) \left(d_3(1 - z_{q_0,s}) + \sqrt{-1}d_4 \frac{-kq_0}{n} + \sqrt{-1}d_4(1 - z_{q_0,s}) \frac{j+L}{n} \right) \right| \\ & \leq 2^k \sum_{i=0}^k \|\psi_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned}$$

By the triangle inequality, it follows from these two inequalities that

$$\begin{aligned} & \left| (1 - z_{q_0,t})^{k-1} e(Ls/n) e(jt/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j}{n} \right) \right. \\ & - (1 - z_{q_0,t})^{k-1} e((j+L)t/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j+L}{n} \right) \\ & \left. - (1 - z_{q_0,s})^{k-1} e((j+L)s/n) \sqrt{-1}d_4(1 - z_{q_0,s}) \frac{L}{n} \right| \\ & \leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2^k \sum_{i=0}^k \|\psi_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned} \quad (125)$$

Multiply both sides with $e(Ls/n)$ again we obtain

$$\begin{aligned} & \left| (1 - z_{q_0,t})^{k-1} e(2Ls/n) e(jt/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j}{n} \right) \right. \\ & - (1 - z_{q_0,t})^{k-1} e(Ls/n) e((j+L)t/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j+L}{n} \right) \\ & \left. - (1 - z_{q_0,s})^{k-1} e((j+2L)s/n) \sqrt{-1}d_4(1 - z_{q_0,s}) \frac{L}{n} \right| \\ & \leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2^k \sum_{i=0}^k \|\psi_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned}$$

Applying the triangle inequality once more, it follows from this inequality and from (125) applied for j being replaced by $j + L$ we can eliminate $(1 - z_s)^{k-1} e((j+2L)s/n)$ and obtain

$$\begin{aligned} & \left| (1 - z_{q_0,t})^{k-1} e(2Ls/n) e(jt/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j}{n} \right) \right. \\ & - (1 - z_{q_0,t})^{k-1} e(Ls/n) e((j+L)t/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j+L}{n} \right) \\ & - (1 - z_{q_0,t})^{k-1} e(Ls/n) e((j+L)t/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j+L}{n} \right) \\ & \left. + (1 - z_{q_0,t})^{k-1} e((j+2L)t/n) \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j+2L}{n} \right) \right| \\ & \leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2^k \sum_{i=0}^k \|\psi_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\ & + 2^k \sum_{i=0}^k \|\psi_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2^k \sum_{i=0}^k \|\psi_{j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \end{aligned}$$

After pulling out the common factor $(1 - z_{q_0,t})^{k-1}e(jt/n)$ and simplifying we then have

$$\begin{aligned}
& |(1 - z_{q_0,t})^{k-1}| \times \left| \left(d_1(1 - z_{q_0,t}) + \sqrt{-1}d_2 \frac{-kq_0}{n} + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{j}{n} \right) (e(Ls/n) - e(Lt/n))^2 \right. \\
& \quad \left. + \sqrt{-1}d_2(1 - z_{q_0,t}) \frac{2L}{n} (e(2Lt/n) - e(Ls/n)e(Lt/n)) \right| \\
& \leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi_{j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\
& \quad + 2^k \sum_{i=0}^k \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi'_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \tag{126}
\end{aligned}$$

Now if we apply (126) for $j = j_1$ and $j = j_1 + j$, and then use the triangle inequality once more we have

$$\begin{aligned}
& |(1 - z_{q_0,t})^{k-1}| \times \left(d_2(1 - z_{q_0,t}) \frac{j}{n} \right) |(e(L(s-t)/n) - 1)|^2 \\
& \leq 2^k \sum_{i=0}^k \|\psi_{j_1+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi_{j_1+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi_{j_1+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\
& \quad + 2^k \sum_{i=0}^k \|\psi'_{j_1+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi'_{j_1+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j_1+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\
& \quad + 2^k \sum_{i=0}^k \|\psi_{j_1+j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi_{j_1+j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi_{j_1+j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\
& \quad + 2^k \sum_{i=0}^k \|\psi'_{j_1+j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi'_{j_1+j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j_1+j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}}. \tag{127}
\end{aligned}$$

The bound for $|(1 - z_{q_0,s})^{k-1}| \times \left(d_4(1 - z_{q_0,s}) \frac{j}{n} \right) |(e(L(s-t)/n) - 1)|^2$ is identical, and we omit. □

Using the obtained inequalities, we can conclude the section as follows.

Proof. (of Proposition 10.2) Recall the notation $J = [a, b]$. We will work with q_0 so that $kq_0 \leq |J|/2$. We divide the treatment into two cases depending on the parameters d_1, d_2, d_3, d_4 .

Subcase 1. Assume that either $|d_2| \geq n^{-\tau_*}$ or $|d_4| \geq n^{-\tau_*}$. Without loss of generality we assume the first case. Notice that with $j = n^{1-8\tau_*}$, then $|jd_2|/n \geq n^{-9\tau_*}$, while as by Condition 2, $\|(s-t)/n\|_{\mathbb{R}/\mathbb{Z}} \geq n^{-1+8\tau_*}$, so there exists $1 \leq L \leq |J|/2 = n^{1-6\tau_*}/2$ such that $|e(L(s-t)/n) - 1| \geq n^{-\tau_*}$. Let us fix such an L .

We next observe that the RHS of (127), for some j_1 from the interval $a \leq j_1 \leq b - 2L - kq_0$ (there are at least $|J|/3$ such j_1), is at most $O(n^{-1/2+7\tau_*})$. Indeed, this is because the sum of the RHS of (127) for j_1 from the above range is bounded by $O(\sum_{j \in J} \|\psi_j\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_j\|_{\mathbb{R}/\mathbb{Z}}) = O(\sqrt{nT}) = O(n^{1/2+\tau_*})$ (as each term $\|\psi_i\|_{\mathbb{R}/\mathbb{Z}}$ appears a bounded number of times in the total sum).

Putting together, with such choice of L and j_1 , the equation (127) implies that

$$|(1 - z_{q_0,t})^k| \times n^{-9\tau_*} \leq O(n^{-1/2+7\tau_*}).$$

Thus we have that $|(1 - z_{q_0,t})^k| \leq n^{-1/2+16\tau_*}$ in this case, and so with sufficiently small τ_*

$$|1 - z_{q_0,t}| \leq n^{-1/3k}. \tag{128}$$

Recall that $z_{q_0,t} = e(q_0 t/n)$. Because (128) holds for any $q_0 = lp_0$ as long as $kq_0 \leq |J|/2$, we thus have for all $1 \leq l \leq n^{1-8\tau^*}$,

$$|1 - e(lp_0 t/n)| \leq n^{-1/3k}. \quad (129)$$

As a consequence, by Claim 9.6 we then have

$$\|p_0 t/\pi n\|_{\mathbb{R}/\mathbb{Z}} = O(n^{-1+8\tau^*}),$$

which contradicts Condition 2.

Subcase 2. Now we consider the remaining case that $|d_2|, |d_4| \leq n^{-\tau^*}$. Without loss of generality we assume $|d_1| \asymp \|\mathbf{D}\|_2 \geq n^{1-4\tau}$. In the case that $d_1|z_{q_0} - 1| \leq n^{4\tau^*}$ then we have

$$|z_{q_0} - 1| \leq n^{-1+4\tau-4\tau^*}.$$

In the other case that $d_1|z_{q_0} - 1| \geq n^{4\tau^*}$, then with L so that $|e(L(s-t)/n) - 1| \geq n^{-\tau^*}$ as in the previous subcase, the factor $d_1|z_{q_0} - 1|((e(L(s-t)/n) - 1))$ is at least $n^{3\tau^*}$, which clearly dominates all other terms involving d_2 on the LHS of (126) (because each of which has order $O(1)$ as d_2 is small). So by (126) we have

$$\begin{aligned} |(1 - z_{q_0})^k|d_1/2 &\leq 2^k \sum_{i=0}^k \|\psi_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi_{j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\ &\quad + 2^k \sum_{i=0}^k \|\psi'_{j+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + 2\|\psi'_{j+L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} + \|\psi'_{j+2L+iq_0}\|_{\mathbb{R}/\mathbb{Z}} \\ &= O(1). \end{aligned}$$

Hence,

$$|z_{q_0} - 1| = O(1/d_1^{1/k}) = O(n^{-1/2k}).$$

Thus from both cases we always have $|z_{q_0} - 1| = O(n^{-1/2k})$. Varying q_0 and using Claim 9.6 we deduce that $\|p_0 t/\pi n\|_{\mathbb{R}/\mathbb{Z}} = O(n^{-1+8\tau^*})$, a contradiction. \square

Finally, similarly to Theorem 9.7, as our approach to prove Proposition 10.2 starts with (117), by passing to subintervals of J when needed we obtain the following analog of Theorem 2.5.

Theorem 10.6. *Let C_* and ℓ be given positive constants, and s, t satisfy Condition 2 for some sufficiently small constant τ . Then following holds for sufficiently large n and sufficiently small τ_* (depending on C_*, τ and ℓ). For any $n^{5\tau-1/2} \leq \|x\|_2 \leq n^{C_*}$, for any set $I \subset [n]$ of at most ℓ entries we have*

$$\left| \prod_{i \notin I} \phi_i(x) \right| \leq \exp(-n^{\tau_*}).$$

REFERENCES

- [1] R. J. Adler and J. E. Taylor, Random Fields and Geometry, Springer Monographs in Mathematics, (2007).
- [2] J. Angst and G. Poly, A weak Cramér condition and application to Edgeworth expansions, Electron. J. Probab. Volume 22 (2017), paper no. 59, 24 pp.
- [3] J.M. Azaïs and M. Wschebor, Level Sets and Extrema of Random Processes and Fields. Wiley (2009), Hoboken, NJ.
- [4] K. Astala, P. Jones, A. Kupiainen, and E. Saksman, Random conformal weldings, Acta Math. 207 (2011), 203-254.
- [5] J.M. Azaïs, F. Dalmao, J. Leon, CLT for the zeros of classical random trigonometric polynomials. Ann. Inst. Henri-Poincare. 52(2), 804-820 (2016).
- [6] J. M. Azaïs, J. Leon, CLT for crossings of random trigonometric polynomials. Electron. J. Probab. 18 (68), 1-17 (2013).
- [7] V. Bally, L. Caramellino, and G. Poly, Convergence in distribution norms in the CLT for non identical distributed random variables. Electron. J. Probab. 23(45), 1-51 (2018).
- [8] V. Bally, L. Caramellino, and G. Poly, Non universality for the variance of the number of real roots of random trigonometric polynomials. Probab. Theory Relat. Fields (2018). <https://doi.org/10.1007/s00440-018-0869-2>, arXiv:1711.03316.
- [9] R. N. Bhattacharya and R. Rao. Normal approximation and asymptotic expansions. Robert E. Krieger Publishing Co., Inc., Melbourne, FL, 1986. Reprint of the 1976 original. MR-0855460.

- [10] A. T. Bharucha-Reid and M. Sambandham, Random polynomials, Probability and Mathematical Statistics. Academic Press, Inc., Orlando, Fla., 1986.
- [11] P. Bleher and X. Di. Correlations between zeros of a random polynomial. *Journal of Statistical Physics*, 88(1):269:305, (1997) .
- [12] P. Bleher and X. Di. Correlations between zeros of non-gaussian random polynomials. *International Mathematics Research Notices*, 2004(46):2443–2484, 2004.
- [13] A. Bloch and G. Pólya, On the roots of certain algebraic equations, *Proc. London Math. Soc.* 33(1932), 102-114.
- [14] E. Bogomolny, O. Bohigas, and P. Leboeuf, Quantum chaotic dynamics and random polynomials, *J. Statist. Phys.* 85 (1996), nos. 5-6, 639-679.
- [15] T. Bloom and D. Dauvergne. "Asymptotic zero distribution of random orthogonal polynomials." arXiv preprint arXiv:1801.10125 (2018).
- [16] H. Cramér. On the composition of elementary errors. *Scandinavian Actuarial Journal*, 1928(1):13-74, 1928.
- [17] Y. Do, Oanh Nguyen, and Van Vu, Roots of random polynomials with coefficients with polynomial growth. *Annals of probability*, to appear.
- [18] Y. Do, Hoi Nguyen, and Van Vu, Real roots of random polynomials: expectation and repulsion, *Proceedings London Mathematical Society* (2015), Vol. 111 (6), 1231-1260.
- [19] A. Edelman and E. Kostlan, How many zeros of a random polynomial are real?, *Bull. Amer. Math. Soc. (N.S.)* 32 (1995), 1–37. Erratum: *Bull. Amer. Math. Soc. (N.S.)* 33 (1996), 325.
- [20] T. Erdélyi, Extensions of the Bloch-Pólya theorem on the number of real zeroes of polynomials, *J. Théor. Nombres Bordeaux* 20 (2008), no. 2, 281-287.
- [21] P. Erdős and A. C. Offord, On the number of real roots of a random algebraic equation, *Proc. London Math. Soc.* 6 (1956), 139–160.
- [22] F. Y. Edgeworth. The generalised law of error, or law of great numbers. *Journal of the Royal Statistical Society*, 69(3):497-539, 1906.
- [23] C. G. Esseen, On the Kolmogorov-Rogozin inequality for the concentration function, *Z. Wahrsch. Verw. Gebiete* 5 (1966), 210-216.
- [24] K. Farahmand, Topics in random polynomials, Pitman research notes in mathematics, series 393. Longman, Harlow, 1998.
- [25] H. Flasche and Z. Kabluchko. Expected number of real zeros of random taylor series. arXiv preprint arXiv:1709.02937, (2017)
- [26] H. Flasche and Z. Kabluchko. Real zeros of random analytic functions associated with geometries of constant curvature. arXiv preprint arXiv:1802.02390, 2018.
- [27] A. Granville and I. Wigman. The distribution of the zeros of random trigonometric polynomials. *Amer. J. Math.* 133 (2) (2011) 295-357.
- [28] G. Halász, Estimates for the concentration function of combinatorial number theory and probability, *Period. Math. Hungar.* 8 (1977), no. 3-4, 197-211.
- [29] I. A. Ibragimov and N. B. Maslova, The average number of zeros of random polynomials, *Vestnik Leningrad. Univ.* 23 (1968), 171–172.
- [30] I. A. Ibragimov and N. B. Maslova, The mean number of real zeros of random polynomials. I. Coefficients with zero mean, *Theor. Probability Appl.* 16 (1971), 228–248.
- [31] I. A. Ibragimov and N. B. Maslova, The mean number of real zeros of random polynomials. II. Coefficients with a nonzero mean., *Theor. Probability Appl.* 16 (1971), 485–493.
- [32] I. A. Ibragimov and N. B. Maslova, The average number of real roots of random polynomials, *Soviet Math. Dokl.* 12 (1971), 1004–1008.
- [33] I. A. Ibragimov and O. Zeitouni. On roots of random polynomials. *Trans. American Math. Soc.* 349 (1997), 2427–2441.
- [34] A. Iksanov, Z. Kabluchko, and A. Marynych. Local universality for real roots of random trigonometric polynomials. *Electronic Journal of Probability*, 21, 2016.
- [35] Z. Kabluchko and D. Zaporozhets. Universality for zeros of random analytic functions. arXiv preprint, arXiv:1205.5355, 2012.
- [36] Z. Kabluchko and D. Zaporozhets. Asymptotic distribution of complex zeros of random analytic functions. *The Annals of Probability*, 42(4):1374:1395, 2014.
- [37] M. Kac, On the average number of real roots of a random algebraic equation, *Bull. Amer. Math. Soc.* 49 (1943) 314–320.
- [38] M. Kac, On the average number of real roots of a random algebraic equation. II. *Proc. London Math. Soc.* 50, (1949), 390–408.

- [39] M. Kac, Probability and related topics in physical sciences, Lectures in Applied Mathematics, Proceedings of the Summer Seminar, Boulder, Colo., 1957, Vol. I Interscience Publishers, London-New York, 1959.
- [40] E. Kostlan, On the distribution of roots of random polynomials, Chapter 38, From Topology to Computation: Proceeding of the Samefest, edited by M. W. Hirsch, J.E. , Marsden and M. Shub, Springer-Verlag, NY 1993.
- [41] S. V. Konyagin and W. Schlag, Lower bounds for the absolute value of random polynomials on a neighborhood of the unit circle, Transactions AMS 351 (1999), 4963-4980.
- [42] J. E. Littlewood and A. C. Offord, On the distribution of the zeros and a -values of a random integral function. I., J. Lond. Math. Soc., 20 (1945), 120-136.
- [43] J. E. Littlewood and A. C. Offord, On the number of real roots of a random algebraic equation. II. Proc. Cambridge Philos. Soc. 35, (1939), 133-148.
- [44] J. E. Littlewood and A. C. Offord, On the number of real roots of a random algebraic equation. III. Rec. Math. [Mat. Sbornik] N.S. 54, (1943), 277-286.
- [45] J. E. Littlewood and A. C. Offord, On the distribution of the zeros and values of a random integral function. II., Ann. Math. 49 (1948), 885-952. Errata, 50 (1949), 990-991. 976), 35-58.
- [46] B. F. Logan and L. A. Shepp, Real zeros of random polynomials. Proc. London Math. Soc. 18 (1968), 29-35.
- [47] B. F. Logan and L. A. Shepp, Real zeros of random polynomials. II. Proc. London Math. Soc. 18 (1968), 308-314.
- [48] N. B. Maslova, The variance of the number of real roots of random polynomials. Teor. Vero- jatnost. i Primenen. 19 (1974), 36-51.
- [49] N. B. Maslova, The distribution of the number of real roots of random polynomials. Theor. Probability Appl. 19 (1974), 461-473
- [50] H. Nguyen, O. Nguyen and Van Vu, On the number of real roots of random polynomials, Communications in Contemporary Mathematics (2016) Vol. 18, 4, 1550052.
- [51] H. Nguyen and O. Zeitouni, Random trigonometric polynomials: concentration on the number of roots, preprint.
- [52] O. Nguyen, V. Vu, Roots of random functions: A general condition for local universality, arxiv.org/abs/1711.03615.
- [53] O. Nguyen, V. Vu, Random polynomials: central limit theorems for the real roots. [arXivpreprintarXiv:1904.04347](https://arxiv.org/abs/1904.04347).
- [54] I. Pritsker, and K. Ramachandran, Equidistribution of zeros of random polynomials, J. Approx. Theory, 215 106-117, (2017)
- [55] S.O. Rice, Mathematical analysis of random noise, Bell System Tech. J. 23: 282-332.
- [56] I. Rivin, Random space and plane curves, arxiv.org/abs/1607.05239.
- [57] C. Qualls, On the number of zeros of a stationary Gaussian random trigonometric polynomial, J. London Math. Soc. (2) 2 (1970), 216-220.
- [58] D.C. Stevens, The average number of real zeros of a random polynomial. Comm. Pure Appl. Math. 22 (1969), 457-477.
- [59] I. Todhunter, A history of the mathematical theory of probability, Stechert, New York, 1931.
- [60] T. Tao and V. Vu, Random matrices: The Circular Law, Communication in Contemporary Mathematics 10 (2008), 261-307.
- [61] T. Tao and V. Vu, Local universality of zeroes of random polynomials. International Mathematics Research Notices, page rnu084, 2014.
- [62] T. Tao's note, terrytao.wordpress.com/2010/03/28/254b-notes-1-equidistribution-of-polynomial-sequences-in-torii/.
- [63] P. Tchebycheff. Sur deux théoremes relatifs aux probabilités. Acta Math., 14(1):305-315, 1890.
- [64] J. E. Wilkins, An asymptotic expansion for the expected number of real zeros of a random polynomial, Proc. Amer. Math. Soc. 103 (1988), 1249-1258.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF VIRGINIA, 141 CABELL DRIVE, CHARLOTTESVILLE, VA 22904

Email address: yendo@virginia.edu

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, 231 W 18TH AVE, COLUMBUS, OH 43210 USA

Email address: nguyen.1261@math.osu.edu

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, 304 WASHINGTON ROAD, PRINCETON, NJ 08540 USA

Email address: onguyen@math.princeton.edu