

A new approach to Regularity and Singularity

Questions in some PDEs including

3-D Navier-Stokes

Saleh Tanveer

(Ohio State University)

Collaborators: Ovidiu Costin, Guo Luo

Research supported in part by

- **IMS (Imperial College), EPSRC & NSF.**

Regularity and Singularity in PDEs-background

- PDEs modeling physical phenomena typically include some effects while ignoring others.
- **Existence and uniqueness questions of smooth solutions fundamental to relevance of a PDE model, as is blow-up.**
- **Global existence of evolutionary PDE solutions typically rely on "Energy" methods. Control over sufficiently higher order Sobolev norm often necessary.**
- **Numerical discretization not rigorously controllable, generally. Further, numerical resolution becomes an issue in higher dimensions.**

Navier-Stokes existence–background

- **Global Existence of smooth 3-D Navier-Stokes solution is an important open problem.**
- **Deviation from linear stress-strain relation or incompressibility is potentially important if N-S solutions are singular**
- **Globally smooth solutions known only when Reynolds number small**
- **Generally, smooth solutions for smooth data on $[0, T]$ known to exist, for T scaling inversely with initial data/forcing.**
- **Global weak solutions known since Leray, but not known whether they are unique. For unforced problem in \mathbb{T}^3 , such a solution becomes smooth again for $t > T_c$, T_c depends on IC**

Borel Summation—background and main idea

- Borel summation generates a 1-1 correspondence between series and functions that preserve algebraic operations (Ecalte, Costin,..).
- Borel sum can involve large or small variable(s)/ parameter(s).
- Formal expansion for $t \ll 1$: $v(x, t) = v_0(x) + \sum_{m=1}^{\infty} t^m v_m(x)$ obtained algorithmically by plugging into $v_t = \mathcal{N}[v]$, where \mathcal{N} being some differential operator. Series divergent
- Borel Sum of this series gives actual solution, which transcends restriction $t \ll 1$
- For NS or Burger's equation, Borel sum given by:

$$v(x, t) = v_0(x) + \int_0^{\infty} U(x, p) e^{-p/t} dp$$

U satisfies Integral equation obtained by inverse LT of PDE.

Integral Equation corresponding to Burger's equation

Plug in $v = v_0(x) + u(x, t)$ into 1-D Burger's to obtain

$$u_t - u_{xx} = -v_0 u_x - u v_{0,x} - u u_x + v_1(x) , \quad v_1(x) = v_0'' - v_0 v_{0,x}$$

with $u(x, 0) = 0$

Inverse Laplace Transform in $1/t$ and Fourier-Transform in x :

$$p\hat{U}_{pp} + 2U_p + k^2\hat{U} = -ik\hat{v}_0 \hat{*}\hat{U} - ik\hat{U} \hat{*}\hat{U} \equiv \hat{G}(k, p)$$

Inverting left side using $\hat{U}(k, 0) = 0$ gives:

$$\hat{U}(k, p) = \int_0^p \mathcal{K}(p, p'; k) \hat{G}(k, p') dp' + \hat{U}^{(0)}(k, p) \equiv \mathcal{N}[\hat{U}](k, p)$$

$$\mathcal{K}(p, p'; k) = \frac{ik\pi}{z} \{z' Y_1(z') J_1(z) - z' Y_1(z) J_1(z')\}$$

$$z = 2|k|\sqrt{p} , \quad z' = 2|k|\sqrt{p'} , \quad \hat{U}^{(0)}(k, p) = 2 \frac{J_1(z)}{z} \hat{v}_1(k)$$

Solution to integral equation $\hat{U} = \mathcal{N}[\hat{U}]$

$$|\mathcal{K}(p, p'; k)| \leq \frac{C}{\sqrt{p}}, \quad C \text{ a constant}$$

$$\|\hat{F}(\cdot, p) \hat{*} \hat{G}(\cdot, p)\|_{L^1(\mathbb{R}^3)} \leq C \|\hat{F}(\cdot, p)\|_{L^1(\mathbb{R}^3)} \|\hat{G}(\cdot, p)\|_{L^1(\mathbb{R}^3)}$$

Define for functions of $F(p, k)$ the norm:

$$\|F\|^{(\alpha)} = \int_0^\infty e^{-\alpha p} \|F(\cdot, p)\|_{L^1(\mathbb{R}^3)} dp, \text{ then can show}$$

$$\|F \hat{*} G\|^{(\alpha)} \leq C \|F\|^{(\alpha)} \|G\|^{(\alpha)}$$

Using above, can show \mathcal{N} contractive for large α ; implies integral equation has unique solution and so Burger PDE has continuous solution for $\text{Re } \frac{1}{t} > \alpha$ as $v(x, t) = v_0(x) + \int_0^\infty e^{-p/t} U(x, p) dp$

Global PDE solution if $\|\hat{U}(\cdot, p)\|_{L^1(\mathbb{R}^3)}$ does not grow as $p \rightarrow \infty$

Incompressible 3-D Navier-Stokes in Fourier-Space

Consider 3-D N-S in infinite geometry or periodic box. Similar results expected for finite domain with no-slip BC using eigenfunctions of Stokes operator as basis. In Fourier-Space

$$\hat{v}_t + \nu |k|^2 \hat{v} = -ik_j P_k [\hat{v}_j \hat{*} \hat{v}] + \hat{f}(k)$$

$$P_k = \left(I - \frac{k(k \cdot)}{|k|^2} \right), \quad \hat{v}(k, 0) = \hat{v}_0(k)$$

where P_k is the Hodge projection in Fourier space, $\hat{f}(k)$ is the Fourier-Transform of forcing $f(x)$, assumed divergence free and t -independent. Subscript j denotes the j -th component of a vector. $k \in \mathbb{R}^3$ or \mathbb{Z}^3 . Einstein convention for repeated index followed. $\hat{*}$ denotes Fourier convolution.

Decompose $\hat{v} = \hat{v}_0 + \hat{u}(k, t)$, inverse-Laplace Transform in $1/t$ and invert the differential operator on the left side

Integral equation associated with Navier-Stokes

We obtain:

$$\hat{U}(k, p) = \int_0^p \mathcal{K}_j(p, p'; k) \hat{H}_j(k, p') dp' + \hat{U}^{(0)}(k, p) \equiv \mathcal{N} [\hat{U}] (k, p) \quad (1)$$

$$\mathcal{K}_j(p, p'; k) = \frac{ik_j \pi}{z} \{z' Y_1(z') J_1(z) - z' Y_1(z) J_1(z')\}$$

$$z = 2|k| \sqrt{\nu p}, \quad z' = 2|k| \sqrt{\nu p'}, \quad \hat{H}_j = P_k \left\{ \hat{v}_{0,j} \hat{*} \hat{U} + \hat{U}_j \hat{*} \hat{v}_0 + \hat{U}_j \hat{*} \hat{U} \right\}$$

$$\hat{U}^{(0)}(k, p) = 2 \frac{J_1(z)}{z} \hat{v}_1(k), \quad P_k = \left(I - \frac{k(k \cdot)}{|k|^2} \right)$$

$$\hat{v}_1(k) = (-\nu |k|^2 \hat{v}_0 - ik_j \mathcal{P}_k [\hat{v}_{0,j} \hat{*} \hat{v}_0]) + \hat{f}(k),$$

$\hat{*}$, denotes Fourier Convolution, $*$ denotes Laplace convolution, while $\hat{*}$ denotes Fourier followed by Laplace convolution. J_1 and Y_1 are the usual Bessel functions.

Results for Integral equation and Navier-Stokes-1

Theorem: If $\|\hat{v}_0\|_{l^1(\mathbb{Z}^3)}, \|\hat{f}\|_{l^1(\mathbb{Z}^3)} < \infty$ then there exists some α so that integral equation $\hat{U} = \mathcal{N}[\hat{U}]$ has a unique solution for $p \in \mathbb{R}^+$ in the space of functions $\{\hat{U} : \|\hat{U}\|^{(\alpha)} < \infty\}$. Further, $\hat{v}(k, t) = \hat{v}_0(k) + \int_0^\infty \hat{U}(k, p)e^{-p/t}$ solves 3-D Navier-Stokes in Fourier-Space; the corresponding $v(x, t)$ is a classical Navier-Stokes solution for $t \in (0, \alpha^{-1})$.

Remark 1: Local existence results in Theorem 1 already known through classical methods. In the present formulation, global PDE existence is a question of asymptotics of known solution to integral equation in the sense that a sub-exponential growth of \hat{U} as $p \rightarrow \infty$ implies global existence of PDE solution.

More Remarks on Theorem 1 for 3-D Navier-Stokes

Remark 2: Errors in Numerical solutions rigorously controlled.
Discretization in p and Galerkin approximation in k results in:

$$\begin{aligned}\hat{U}_\delta(k, m\delta) &= \delta \sum_{m'=0}^m \mathcal{K}_{m,m'} \mathcal{P}_N \mathcal{H}_\delta(k, m'\delta) + \hat{U}^{(0)}(k, m\delta) \\ &\equiv \mathcal{N}_\delta \left[\hat{U}_\delta \right] \quad \text{for } k_j = -N, \dots, N, \quad j = 1, 2, 3\end{aligned}$$

\mathcal{P}_N is the Galerkin Projection into N -Fourier modes. \mathcal{N}_δ has properties similar to \mathcal{N} . The continuous solution \hat{U} satisfies $\hat{U} = \mathcal{N}_\delta \left[\hat{U} \right] + E$, where E is the truncation error. Thus, $\hat{U} - \hat{U}_\delta$ can be estimated using same tools as in Theorem 1.

Note: Similar control over discretized solutions to PDEs not available since truncation errors involve derivatives of PDE solution which are not known to exist beyond a short-time.

Numerical Solutions to integral equation

We choose the Kida initial conditions and forcing

$$v_0(\mathbf{x}) = (v_1(x_1, x_2, x_3, 0), v_2(x_1, x_2, x_3, 0), v_3(x_1, x_2, x_3, 0))$$

$$v_1(x_1, x_2, x_3, 0) = v_2(x_3, x_1, x_2, 0) = v_3(x_2, x_3, x_1, 0)$$

$$v_1(x_1, x_2, x_3, 0) = \sin x_3 (\cos 3x_2 \cos x_3 - \cos x_2 \cos 3x_3)$$


$$f_1(x_1, x_2, x_3) = \frac{1}{5} v_1(x_1, x_2, x_3, 0)$$

High Degree of Symmetry makes computationally less expensive

**Corresponding Euler problem believed to blow up in finite time;
so good candidate to study viscous effects**


In the plots, "constant forcing" corresponds to $f = (f_1, f_2, f_3)$ as above, while zero forcing refers to $f = 0$. Recall sub-exponential growth in p corresponds to global N-S solution.

Numerical solution to integral equation-plot-1




$\|\hat{U}(\cdot, p)\|_{l^1}$ vs. p for $\nu = 1$, constant forcing.

Numerical solution to integral equation-plot-2




$\|\hat{U}(\cdot, p)\|_{l^1}$ vs. p for $\nu = 1$, no forcing

Numerical solution to integral equation-plot-3




$\|\hat{U}(\cdot, p)\|_{l^1}$ vs. p for $\nu = 0.16$, constant forcing

Numerical solution to integral equation-plot-4




$\|\hat{U}(\cdot, p)\|_{l^1}$ vs. p for $\nu = 0.1$, constant forcing

Numerical solution to integral equation-plot-5



$\hat{U}(k, p)$ vs. p for $k = (1, 1, 17)$, $\nu = 0.1$, no forcing.

Numerical solution to integral equation-plot-6



$\log \|\hat{U}(\cdot, p)\|_{l^1}$ vs. $\log p$ for $\nu = 0.001$, constant forcing

Issues raised by numerical computations

Numerical solutions to integral equation available on finite interval $[0, p_0]$, yet N-S solution requires $[0, \infty)$ interval since

$$\hat{v}(k, t) = \hat{v}_0 + \int_0^\infty e^{-p/t} \hat{U}(k, p) dp$$

Actually, the integral over $\int_0^{p_0}$ gives an approximate N-S solution, with errors that can be bounded for a time interval $[0, T]$, if computed solution to integral equation eventually decreases with p on a sufficiently large interval $[0, p_0]$.

Further, a non-increasing \hat{U} over a sufficiently large interval $[0, p_0]$ gives smaller bounds on growth rate α as $p \rightarrow \infty$.

Therefore, in such cases smooth NS solution exists over a long interval $[0, \alpha^{-1})$.

Recall for unforced problem in \mathbb{T}^3 , even weak solution to NS becomes smooth for $t > T_c$, with T_c estimated from initial data. Hence global existence follows under some conditions.

Extending Navier-Stokes interval of existence

For $\alpha_0 \geq 0$, define

$$\epsilon = \nu^{-1/2} p_0^{-1/2}, \quad a = \|\hat{v}_0\|_{l^1}, \quad c = \int_{p_0}^{\infty} \|\hat{U}^{(0)}(\cdot, p)\|_{l^1} e^{-\alpha_0 p} dp$$

$$\epsilon_1 = \nu^{-1/2} p_0^{-1/2} \left(2 \int_0^{p_0} e^{-\alpha_0 s} \|\hat{U}(\cdot, s)\|_{l^1} ds + \|\hat{v}_0\|_{l^1} \right)$$

$$b = \frac{e^{-\alpha_0 p_0}}{\sqrt{\nu p_0} \alpha} \int_0^{p_0} \|\hat{U}^* \hat{U} + \hat{v}_0 \cdot \hat{U}\|_{l^1} ds$$


Theorem 3: A smooth solution to 3-D Navier-Stokes equation exists on the interval $[0, \alpha^{-1})$, when $\alpha \geq \alpha_0$ is chosen to satisfy

$$\alpha > \epsilon_1 + 2\epsilon c + \sqrt{(\epsilon_1 + 2\epsilon c)^2 + 4b\epsilon} - \epsilon_1^2$$

Remark: If p_0 is chosen large enough, ϵ, ϵ_1 is small when computed solution in $[0, p_0]$ decays with q . Then α can be chosen rather small.

Relation of Optimal α to Navier-Stokes singularities

$$\hat{U}(k, p) = \frac{1}{2\pi i} \int_{c_0 - i\infty}^{c_0 + i\infty} e^{p/t} [\hat{v}(k, t) - \hat{v}_0(k)] d \left[\frac{1}{t} \right]$$



Rightmost singularity(ies) of NS solution $\hat{v}(k, t)$ in the $1/t$ plane determines optimal α . γ gives dominant oscillation frequency.

Laplace-transform and accelerated representation


To get rid of the effect of complex singularity, it is prudent to seek a more general Laplace-transform involves

$$\hat{v}(k, t) = \hat{v}_0(k) + \int_0^\infty e^{-q/t^n} \hat{U}(k, q) dq$$

We have proved that for the unforced problem, if there are complex singularities t_s in the right-half plane, but not on the real axis, then a nonzero lower bound for $|\arg t_s|$ exists. Then, for sufficiently large n , no singularities in the $\tau = t^{-n}$ plane in the right-half plane. Hence, $\hat{U}(k, q)$ will not grow with q

$\hat{U}(k, q)$ satisfies an integral equation similar to the one satisfied by $\hat{U}(k, p)$ and Theorems similar to Theorem 1 follow. In the context of ODEs, change of variable $p \rightarrow q$ is called acceleration (Ecale)

$\|\hat{U}(\cdot, q)\|_{l^1}$ vs. q , $n = 2$, $\nu = 0.1$



Kida I.C. $v_1^{(0)} = \sin x_1 (\cos 3x_2 \cos x_3 - \cos x_2 \cos 3x_3)$

Other components from cyclic relation:

$$v_1^{(0)}(x_1, x_2, x_3) = v_1^{(0)}(x_3, x_1, x_2) = v_3^{(0)}(x_2, x_3, x_1)$$

Extending Navier-Stokes interval of existence

For $\alpha_0 \geq 0$, define

$$\epsilon_1 = \nu^{-1/2} q_0^{-1+1/(2n)}, \quad c = \int_{q_0}^{\infty} \|\hat{U}^{(0)}(\cdot, q)\|_{l^1} e^{-\alpha_0 q} dq$$

$$\epsilon_1 = \nu^{-1/2} q_0^{-1+1/(2n)} \left(2 \int_0^{q_0} e^{-\alpha_0 s} \|\hat{U}(\cdot, s)\|_{l^1} ds + \|\hat{v}_0\|_{l^1} \right)$$

$$b = \frac{e^{-\alpha_0 q_0}}{\sqrt{\nu} q_0^{1-1/(2n)} \alpha} \int_0^{q_0} \|\hat{U}_*^* \hat{U} + \hat{v}_0 \cdot \hat{U}\|_{l^1} ds$$

Theorem 4: A smooth solution to 3-D Navier-Stokes equation exists in the $\|\cdot\|_{l^1}$ space on the interval $[0, \alpha^{-1/n})$, when $\alpha \geq \alpha_0$ is chosen to satisfy

$$\alpha > \epsilon_1 + 2\epsilon c + \sqrt{(\epsilon_1 + 2\epsilon c)^2 + 4b\epsilon - \epsilon_1^2}$$

Remark: If q_0 is chosen large enough, ϵ, ϵ_1 is small when computed solution in $[0, q_0]$ decays with q . Then α can be chosen rather small.

Example problems where approach is applicable

- Navier-Stokes with temperature field (Boussinesq approximation)

- Fourth order Parabolic equations of the type:

$$u_t + \Delta^2 u = N[u, Du, D^2 u, D^3 u]$$

- KDV and related equations.

- Magneto-hydrodynamic equation with certain approximations.

- For some PDE problems with finite-time blow-up, blow-up time related to exponent α of exponential growth of IE solution, provided there is no-oscillation even with $p \rightarrow q$ acceleration.

Conclusions

We have shown how Borel summation methods provides an alternate existence theory for PDE Initial value problems like N-S. With this integral equation (IE) approach, the PDE global existence is implied if known solution to IE has subexponential growth at ∞ .

The solution to integral equation in a finite interval can be computed numerically with rigorously controlled errors.

Integral equation in a suitable accelerated variable q will decay exponentially for unforced N-S equation, unless there is a real time singularity of PDE solution.

The computation over a finite $[0, q_0]$ interval gives a refined bound on exponent α at ∞ , and hence a longer existence time $[0, \alpha^{-1/n})$ to 3-D Navier-Stokes.

Approach should be useful in both regularity and singularity studies of more general PDE initial value problems.